




$$(a + b)^2 = a^2 + 2ab + b^2$$



$$y = \cos x$$

$$\Sigma$$


$$\sqrt[n]{z}$$



$$\int_a^b f(x) dx$$

# 1300

# Math

# Formulas

Ketabton.com

Alex Svirin, Ph.D.

# 1300 Math Formulas

ISBN 9949107741

Copyright © 2004 A.Svirin. All Rights Reserved.

**This page is intentionally left blank.**

## Preface

This handbook is a complete desktop reference for students and engineers. It has everything from high school math to math for advanced undergraduates in engineering, economics, physical sciences, and mathematics. The ebook contains hundreds of formulas, tables, and figures from Number Sets, Algebra, Geometry, Trigonometry, Matrices and Determinants, Vectors, Analytic Geometry, Calculus, Differential Equations, Series, and Probability Theory. The structured table of contents, links, and layout make finding the relevant information quick and painless, so it can be used as an everyday online reference guide.

# Contents

<b>1</b>	<b>NUMBER SETS</b>	
1.1	Set Identities	<b>1</b>
1.2	Sets of Numbers	<b>5</b>
1.3	Basic Identities	<b>7</b>
1.4	Complex Numbers	<b>8</b>
<b>2</b>	<b>ALGEBRA</b>	
2.1	Factoring Formulas	<b>12</b>
2.2	Product Formulas	<b>13</b>
2.3	Powers	<b>14</b>
2.4	Roots	<b>15</b>
2.5	Logarithms	<b>16</b>
2.6	Equations	<b>18</b>
2.7	Inequalities	<b>19</b>
2.8	Compound Interest Formulas	<b>22</b>
<b>3</b>	<b>GEOMETRY</b>	
3.1	Right Triangle	<b>24</b>
3.2	Isosceles Triangle	<b>27</b>
3.3	Equilateral Triangle	<b>28</b>
3.4	Scalene Triangle	<b>29</b>
3.5	Square	<b>33</b>
3.6	Rectangle	<b>34</b>
3.7	Parallelogram	<b>35</b>
3.8	Rhombus	<b>36</b>
3.9	Trapezoid	<b>37</b>
3.10	Isosceles Trapezoid	<b>38</b>
3.11	Isosceles Trapezoid with Inscribed Circle	<b>40</b>
3.12	Trapezoid with Inscribed Circle	<b>41</b>

- 3.13 Kite **42**
- 3.14 Cyclic Quadrilateral **43**
- 3.15 Tangential Quadrilateral **45**
- 3.16 General Quadrilateral **46**
- 3.17 Regular Hexagon **47**
- 3.18 Regular Polygon **48**
- 3.19 Circle **50**
- 3.20 Sector of a Circle **53**
- 3.21 Segment of a Circle **54**
- 3.22 Cube **55**
- 3.23 Rectangular Parallelepiped **56**
- 3.24 Prism **57**
- 3.25 Regular Tetrahedron **58**
- 3.26 Regular Pyramid **59**
- 3.27 Frustum of a Regular Pyramid **61**
- 3.28 Rectangular Right Wedge **62**
- 3.29 Platonic Solids **63**
- 3.30 Right Circular Cylinder **66**
- 3.31 Right Circular Cylinder with an Oblique Plane Face **68**
- 3.32 Right Circular Cone **69**
- 3.33 Frustum of a Right Circular Cone **70**
- 3.34 Sphere **72**
- 3.35 Spherical Cap **72**
- 3.36 Spherical Sector **73**
- 3.37 Spherical Segment **74**
- 3.38 Spherical Wedge **75**
- 3.39 Ellipsoid **76**
- 3.40 Circular Torus **78**

#### **4 TRIGONOMETRY**

- 4.1 Radian and Degree Measures of Angles **80**
- 4.2 Definitions and Graphs of Trigonometric Functions **81**
- 4.3 Signs of Trigonometric Functions **86**
- 4.4 Trigonometric Functions of Common Angles **87**
- 4.5 Most Important Formulas **88**

4.6	Reduction Formulas	89
4.7	Periodicity of Trigonometric Functions	90
4.8	Relations between Trigonometric Functions	90
4.9	Addition and Subtraction Formulas	91
4.10	Double Angle Formulas	92
4.11	Multiple Angle Formulas	93
4.12	Half Angle Formulas	94
4.13	Half Angle Tangent Identities	94
4.14	Transforming of Trigonometric Expressions to Product	95
4.15	Transforming of Trigonometric Expressions to Sum	97
4.16	Powers of Trigonometric Functions	98
4.17	Graphs of Inverse Trigonometric Functions	99
4.18	Principal Values of Inverse Trigonometric Functions	102
4.19	Relations between Inverse Trigonometric Functions	103
4.20	Trigonometric Equations	106
4.21	Relations to Hyperbolic Functions	106
<b>5</b>	<b>MATRICES AND DETERMINANTS</b>	
5.1	Determinants	107
5.2	Properties of Determinants	109
5.3	Matrices	110
5.4	Operations with Matrices	111
5.5	Systems of Linear Equations	114
<b>6</b>	<b>VECTORS</b>	
6.1	Vector Coordinates	118
6.2	Vector Addition	120
6.3	Vector Subtraction	122
6.4	Scaling Vectors	122
6.5	Scalar Product	123
6.6	Vector Product	125
6.7	Triple Product	127
<b>7</b>	<b>ANALYTIC GEOMETRY</b>	
7.1	One -Dimensional Coordinate System	130

7.2	Two -Dimensional Coordinate System	131
7.3	Straight Line in Plane	139
7.4	Circle	149
7.5	Ellipse	152
7.6	Hyperbola	154
7.7	Parabola	158
7.8	Three -Dimensional Coordinate System	161
7.9	Plane	165
7.10	Straight Line in Space	175
7.11	Quadric Surfaces	180
7.12	Sphere	189
<b>8</b>	<b>DIFFERENTIAL CALCULUS</b>	
8.1	Functions and Their Graphs	191
8.2	Limits of Functions	208
8.3	Definition and Properties of the Derivative	209
8.4	Table of Derivatives	211
8.5	Higher Order Derivatives	215
8.6	Applications of Derivative	217
8.7	Differential	221
8.8	Multivariable Functions	222
8.9	Differential Operators	225
<b>9</b>	<b>INTEGRAL CALCULUS</b>	
9.1	Indefinite Integral	227
9.2	Integrals of Rational Functions	228
9.3	Integrals of Irrational Functions	231
9.4	Integrals of Trigonometric Functions	237
9.5	Integrals of Hyperbolic Functions	241
9.6	Integrals of Exponential and Logarithmic Functions	242
9.7	Reduction Formulas	243
9.8	Definite Integral	247
9.9	Improper Integral	253
9.10	Double Integral	257
9.11	Triple Integral	269



- 9.12 Line Integral **275**
- 9.13 Surface Integral **285**

## **10 DIFFERENTIAL EQUATIONS**

- 10.1 First Order Ordinary Differential Equations **295**
- 10.2 Second Order Ordinary Differential Equations **298**
- 10.3 Some Partial Differential Equations **302**

## **11 SERIES**

- 11.1 Arithmetic Series **304**
- 11.2 Geometric Series **305**
- 11.3 Some Finite Series **305**
- 11.4 Infinite Series **307**
- 11.5 Properties of Convergent Series **307**
- 11.6 Convergence Tests **308**
- 11.7 Alternating Series **310**
- 11.8 Power Series **311**
- 11.9 Differentiation and Integration of Power Series **312**
- 11.10 Taylor and Maclaurin Series **313**
- 11.11 Power Series Expansions for Some Functions **314**
- 11.12 Binomial Series **316**
- 11.13 Fourier Series **316**

## **12 PROBABILITY**

- 12.1 Permutations and Combinations **318**
- 12.2 Probability Formulas **319**

This page is intentionally left blank.

# Chapter 1

## Number Sets

### 1.1 Set Identities

Sets: A, B, C

Universal set: I

Complement :  $A'$

Proper subset:  $A \subset B$

Empty set:  $\emptyset$

Union of sets:  $A \cup B$

Intersection of sets:  $A \cap B$

Difference of sets:  $A \setminus B$

1.  $A \subset I$
2.  $A \subset A$
3.  $A = B$  if  $A \subset B$  and  $B \subset A$ .
4. Empty Set  
 $\emptyset \subset A$
5. Union of Sets  
 $C = A \cup B = \{x \mid x \in A \text{ or } x \in B\}$

CHAPTER 1. NUMBER SETS




Figure 1.

6. Commutativity

$$A \cup B = B \cup A$$

7. Associativity

$$A \cup (B \cup C) = (A \cup B) \cup C$$

8. Intersection of Sets

$$C = A \cap B = \{x \mid x \in A \text{ and } x \in B\}$$




Figure 2.

9. Commutativity

$$A \cap B = B \cap A$$

10. Associativity

$$A \cap (B \cap C) = (A \cap B) \cap C$$

CHAPTER 1. NUMBER SETS

11. **Distributivity**  
 $A \cup (B \cap C) = (A \cup B) \cap (A \cup C),$   
 $A \cap (B \cup C) = (A \cap B) \cup (A \cap C).$
12. **Idempotency**  
 $A \cap A = A,$   
 $A \cup A = A$
13. **Domination**  
 $A \cap \emptyset = \emptyset,$   
 $A \cup I = I$
14. **Identity**  
 $A \cup \emptyset = A,$   
 $A \cap I = A$
15. **Complement**  
 $A' = \{x \in I \mid x \notin A\}$
16. **Complement of Intersection and Union**  
 $A \cup A' = I,$   
 $A \cap A' = \emptyset$
17. **De Morgan's Laws**  
 $(A \cup B)' = A' \cap B',$   
 $(A \cap B)' = A' \cup B'$
18. **Difference of Sets**  
 $C = B \setminus A = \{x \mid x \in B \text{ and } x \notin A\}$

CHAPTER 1. NUMBER SETS




Figure 3.

- 19.  $B \setminus A = B \setminus (A \cap B)$
- 20.  $B \setminus A = B \cap A'$
- 21.  $A \setminus A = \emptyset$
- 22.  $A \setminus B = A$  if  $A \cap B = \emptyset$ .




Figure 4.

- 23.  $(A \setminus B) \cap C = (A \cap C) \setminus (B \cap C)$
- 24.  $A' = I \setminus A$
- 25. Cartesian Product  
 $C = A \times B = \{(x, y) \mid x \in A \text{ and } y \in B\}$

## CHAPTER 1. NUMBER SETS

### 1.2 Sets of Numbers

Natural numbers:  $\mathbb{N}$

Whole numbers:  $\mathbb{N}_0$

Integers:  $\mathbb{Z}$

Positive integers:  $\mathbb{Z}^+$

Negative integers:  $\mathbb{Z}^-$

Rational numbers:  $\mathbb{Q}$

Real numbers:  $\mathbb{R}$

Complex numbers:  $\mathbb{C}$

**26.** Natural Numbers

Counting numbers:  $\mathbb{N} = \{1, 2, 3, \dots\}$ .

**27.** Whole Numbers

Counting numbers and zero:  $\mathbb{N}_0 = \{0, 1, 2, 3, \dots\}$ .

**28.** Integers

Whole numbers and their opposites and zero:

$$\mathbb{Z}^+ = \mathbb{N} = \{1, 2, 3, \dots\},$$

$$\mathbb{Z}^- = \{\dots, -3, -2, -1\},$$

$$\mathbb{Z} = \mathbb{Z}^- \cup \{0\} \cup \mathbb{Z}^+ = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}.$$

**29.** Rational Numbers

Repeating or terminating decimals:

$$\mathbb{Q} = \left\{ x \mid x = \frac{a}{b} \text{ and } a \in \mathbb{Z} \text{ and } b \in \mathbb{Z} \text{ and } b \neq 0 \right\}.$$

**30.** Irrational Numbers

Nonrepeating and nonterminating decimals.

CHAPTER 1. NUMBER SETS

- 31. Real Numbers  
Union of rational and irrational numbers:  $\mathbb{R}$ .
- 32. Complex Numbers  
 $C = \{x + iy \mid x \in \mathbb{R} \text{ and } y \in \mathbb{R}\}$ ,  
where  $i$  is the imaginary unit.
- 33.  $\mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R} \subset \mathbb{C}$




Figure 5.



CHAPTER 1. NUMBER SETS

## 1.3 Basic Identities

Real numbers:  $a, b, c$

- 34.** Additive Identity  
 $a + 0 = a$
- 35.** Additive Inverse  
 $a + (-a) = 0$
- 36.** Commutative of Addition  
 $a + b = b + a$
- 37.** Associative of Addition  
 $(a + b) + c = a + (b + c)$
- 38.** Definition of Subtraction  
 $a - b = a + (-b)$
- 39.** Multiplicative Identity  
 $a \cdot 1 = a$
- 40.** Multiplicative Inverse  
 $a \cdot \frac{1}{a} = 1, a \neq 0$
- 41.** Multiplication Times 0  
 $a \cdot 0 = 0$
- 42.** Commutative of Multiplication  
 $a \cdot b = b \cdot a$

CHAPTER 1. NUMBER SETS

43. Associative of Multiplication  
 $(a \cdot b) \cdot c = a \cdot (b \cdot c)$

44. Distributive Law  
 $a(b + c) = ab + ac$

45. Definition of Division  
 $\frac{a}{b} = a \cdot \frac{1}{b}$

## 1.4 Complex Numbers

Natural number:  $n$

Imaginary unit:  $i$

Complex number:  $z$

Real part:  $a, c$

Imaginary part:  $bi, di$

Modulus of a complex number:  $r, r_1, r_2$

Argument of a complex number:  $\varphi, \varphi_1, \varphi_2$

46.

$i^1 = i$	$i^5 = i$	$i^{4n+1} = i$
$i^2 = -1$	$i^6 = -1$	$i^{4n+2} = -1$
$i^3 = -i$	$i^7 = -i$	$i^{4n+3} = -i$
$i^4 = 1$	$i^8 = 1$	$i^{4n} = 1$

47.  $z = a + bi$

48. Complex Plane

CHAPTER 1. NUMBER SETS




Figure 6.

49.  $(a + bi) + (c + di) = (a + c) + (b + d)i$

50.  $(a + bi) - (c + di) = (a - c) + (b - d)i$

51.  $(a + bi)(c + di) = (ac - bd) + (ad + bc)i$

52.  $\frac{a + bi}{c + di} = \frac{ac + bd}{c^2 + d^2} + \frac{bc - ad}{c^2 + d^2} \cdot i$

53. Conjugate Complex Numbers

$$\overline{a + bi} = a - bi$$

54.  $a = r \cos \varphi, b = r \sin \varphi$

## CHAPTER 1. NUMBER SETS




Figure 7.

**55. Polar Presentation of Complex Numbers**

$$a + bi = r(\cos \varphi + i \sin \varphi)$$

**56. Modulus and Argument of a Complex Number**

If  $a + bi$  is a complex number, then

$$r = \sqrt{a^2 + b^2} \text{ (modulus),}$$

$$\varphi = \arctan \frac{b}{a} \text{ (argument).}$$

**57. Product in Polar Representation**

$$\begin{aligned} z_1 \cdot z_2 &= r_1(\cos \varphi_1 + i \sin \varphi_1) \cdot r_2(\cos \varphi_2 + i \sin \varphi_2) \\ &= r_1 r_2 [\cos(\varphi_1 + \varphi_2) + i \sin(\varphi_1 + \varphi_2)] \end{aligned}$$

**58. Conjugate Numbers in Polar Representation**

$$\overline{r(\cos \varphi + i \sin \varphi)} = r[\cos(-\varphi) + i \sin(-\varphi)]$$

**59. Inverse of a Complex Number in Polar Representation**

$$\frac{1}{r(\cos \varphi + i \sin \varphi)} = \frac{1}{r} [\cos(-\varphi) + i \sin(-\varphi)]$$

CHAPTER 1. NUMBER SETS

**60. Quotient in Polar Representation**

$$\frac{z_1}{z_2} = \frac{r_1(\cos \varphi_1 + i \sin \varphi_1)}{r_2(\cos \varphi_2 + i \sin \varphi_2)} = \frac{r_1}{r_2} [\cos(\varphi_1 - \varphi_2) + i \sin(\varphi_1 - \varphi_2)]$$

**61. Power of a Complex Number**

$$z^n = [r(\cos \varphi + i \sin \varphi)]^n = r^n [\cos(n\varphi) + i \sin(n\varphi)]$$

**62. Formula “De Moivre”**

$$(\cos \varphi + i \sin \varphi)^n = \cos(n\varphi) + i \sin(n\varphi)$$

**63. Nth Root of a Complex Number**

$$\sqrt[n]{z} = \sqrt[n]{r(\cos \varphi + i \sin \varphi)} = \sqrt[n]{r} \left( \cos \frac{\varphi + 2\pi k}{n} + i \sin \frac{\varphi + 2\pi k}{n} \right),$$

where

$$k = 0, 1, 2, \dots, n-1.$$

**64. Euler’s Formula**

$$e^{ix} = \cos x + i \sin x$$

## Chapter 2

# Algebra

### 2.1 Factoring Formulas

Real numbers:  $a, b, c$   
Natural number:  $n$

**65.**  $a^2 - b^2 = (a + b)(a - b)$

**66.**  $a^3 - b^3 = (a - b)(a^2 + ab + b^2)$

**67.**  $a^3 + b^3 = (a + b)(a^2 - ab + b^2)$

**68.**  $a^4 - b^4 = (a^2 - b^2)(a^2 + b^2) = (a - b)(a + b)(a^2 + b^2)$

**69.**  $a^5 - b^5 = (a - b)(a^4 + a^3b + a^2b^2 + ab^3 + b^4)$

**70.**  $a^5 + b^5 = (a + b)(a^4 - a^3b + a^2b^2 - ab^3 + b^4)$

**71.** If  $n$  is odd, then  
 $a^n + b^n = (a + b)(a^{n-1} - a^{n-2}b + a^{n-3}b^2 - \dots - ab^{n-2} + b^{n-1}).$

**72.** If  $n$  is even, then  
 $a^n - b^n = (a - b)(a^{n-1} + a^{n-2}b + a^{n-3}b^2 + \dots + ab^{n-2} + b^{n-1}),$

## CHAPTER 2. ALGEBRA

$$a^n + b^n = (a + b)(a^{n-1} - a^{n-2}b + a^{n-3}b^2 - \dots + ab^{n-2} - b^{n-1}).$$

## 2.2 Product Formulas

Real numbers:  $a, b, c$ Whole numbers:  $n, k$ 

$$73. \quad (a - b)^2 = a^2 - 2ab + b^2$$

$$74. \quad (a + b)^2 = a^2 + 2ab + b^2$$

$$75. \quad (a - b)^3 = a^3 - 3a^2b + 3ab^2 - b^3$$

$$76. \quad (a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$$

$$77. \quad (a - b)^4 = a^4 - 4a^3b + 6a^2b^2 - 4ab^3 + b^4$$

$$78. \quad (a + b)^4 = a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4$$

## 79. Binomial Formula

$$(a + b)^n = {}^nC_0 a^n + {}^nC_1 a^{n-1}b + {}^nC_2 a^{n-2}b^2 + \dots + {}^nC_{n-1} ab^{n-1} + {}^nC_n b^n,$$

where  ${}^nC_k = \frac{n!}{k!(n-k)!}$  are the binomial coefficients.

$$80. \quad (a + b + c)^2 = a^2 + b^2 + c^2 + 2ab + 2ac + 2bc$$

$$81. \quad (a + b + c + \dots + u + v)^2 = a^2 + b^2 + c^2 + \dots + u^2 + v^2 + \\ + 2(ab + ac + \dots + au + av + bc + \dots + bu + bv + \dots + uv)$$

## 2.3 Powers

Bases (positive real numbers):  $a, b$

Powers (rational numbers):  $n, m$

$$82. \quad a^m a^n = a^{m+n}$$

$$83. \quad \frac{a^m}{a^n} = a^{m-n}$$

$$84. \quad (ab)^m = a^m b^m$$

$$85. \quad \left(\frac{a}{b}\right)^m = \frac{a^m}{b^m}$$

$$86. \quad (a^m)^n = a^{mn}$$

$$87. \quad a^0 = 1, a \neq 0$$

$$88. \quad a^1 = a$$

$$89. \quad a^{-m} = \frac{1}{a^m}$$

$$90. \quad a^{\frac{m}{n}} = \sqrt[n]{a^m}$$



## 2.4 Roots

Bases:  $a, b$ Powers (rational numbers):  $n, m$  $a, b \geq 0$  for even roots ( $n = 2k, k \in \mathbb{N}$ )

91.  $\sqrt[n]{ab} = \sqrt[n]{a} \sqrt[n]{b}$
92.  $\sqrt[n]{a} \sqrt[m]{b} = \sqrt[nm]{a^m b^n}$
93.  $\sqrt[n]{\frac{a}{b}} = \frac{\sqrt[n]{a}}{\sqrt[n]{b}}, b \neq 0$
94.  $\frac{\sqrt[n]{a}}{\sqrt[m]{b}} = \frac{\sqrt[nm]{a^m}}{\sqrt[nm]{b^n}} = \sqrt[nm]{\frac{a^m}{b^n}}, b \neq 0.$
95.  $(\sqrt[n]{a^m})^p = \sqrt[n]{a^{mp}}$
96.  $(\sqrt[n]{a})^n = a$
97.  $\sqrt[n]{a^m} = \sqrt[np]{a^{mp}}$
98.  $\sqrt[n]{a^m} = a^{\frac{m}{n}}$
99.  $\sqrt[m]{\sqrt[n]{a}} = \sqrt[mn]{a}$
100.  $(\sqrt[n]{a})^m = \sqrt[n]{a^m}$

CHAPTER 2. ALGEBRA

$$101. \frac{1}{\sqrt[n]{a}} = \frac{\sqrt[n]{a^{n-1}}}{a}, a \neq 0.$$

$$102. \sqrt{a \pm \sqrt{b}} = \sqrt{\frac{a + \sqrt{a^2 - b}}{2}} \pm \sqrt{\frac{a - \sqrt{a^2 - b}}{2}}$$

$$103. \frac{1}{\sqrt{a \pm \sqrt{b}}} = \frac{\sqrt{a \mp \sqrt{b}}}{a - b}$$

## 2.5 Logarithms

Positive real numbers:  $x, y, a, c, k$

Natural number:  $n$

### 104. Definition of Logarithm

$y = \log_a x$  if and only if  $x = a^y$ ,  $a > 0$ ,  $a \neq 1$ .

$$105. \log_a 1 = 0$$

$$106. \log_a a = 1$$

$$107. \log_a 0 = \begin{cases} -\infty & \text{if } a > 1 \\ +\infty & \text{if } a < 1 \end{cases}$$

$$108. \log_a(xy) = \log_a x + \log_a y$$

$$109. \log_a \frac{x}{y} = \log_a x - \log_a y$$

CHAPTER 2. ALGEBRA

**110.**  $\log_a(x^n) = n \log_a x$

**111.**  $\log_a \sqrt[n]{x} = \frac{1}{n} \log_a x$

**112.**  $\log_a x = \frac{\log_c x}{\log_c a} = \log_c x \cdot \log_a c, c > 0, c \neq 1.$

**113.**  $\log_a c = \frac{1}{\log_c a}$

**114.**  $x = a^{\log_a x}$

**115.** Logarithm to Base 10  
 $\log_{10} x = \log x$

**116.** Natural Logarithm  
 $\log_e x = \ln x,$

where  $e = \lim_{k \rightarrow \infty} \left(1 + \frac{1}{k}\right)^k = 2.718281828\dots$

**117.**  $\log x = \frac{1}{\ln 10} \ln x = 0.434294 \ln x$

**118.**  $\ln x = \frac{1}{\log e} \log x = 2.302585 \log x$

## 2.6 Equations

Real numbers:  $a, b, c, p, q, u, v$

Solutions:  $x_1, x_2, y_1, y_2, y_3$

**119.** Linear Equation in One Variable

$$ax + b = 0, x = -\frac{b}{a}.$$

**120.** Quadratic Equation

$$ax^2 + bx + c = 0, x_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

**121.** Discriminant

$$D = b^2 - 4ac$$

**122.** Viète's Formulas

If  $x^2 + px + q = 0$ , then

$$\begin{cases} x_1 + x_2 = -p \\ x_1 x_2 = q \end{cases}.$$

**123.**  $ax^2 + bx = 0, x_1 = 0, x_2 = -\frac{b}{a}.$

**124.**  $ax^2 + c = 0, x_{1,2} = \pm \sqrt{-\frac{c}{a}}.$

**125.** Cubic Equation. Cardano's Formula.

$$y^3 + py + q = 0,$$

## CHAPTER 2. ALGEBRA

$$y_1 = \mathbf{u} + \mathbf{v}, \quad y_{2,3} = -\frac{1}{2}(\mathbf{u} + \mathbf{v}) \pm \frac{\sqrt{3}}{2}(\mathbf{u} + \mathbf{v})\mathbf{i},$$

where

$$\mathbf{u} = \sqrt[3]{-\frac{\mathbf{q}}{2} + \sqrt{\left(\frac{\mathbf{q}}{2}\right)^2 + \left(\frac{\mathbf{p}}{3}\right)^2}}, \quad \mathbf{v} = \sqrt[3]{-\frac{\mathbf{q}}{2} - \sqrt{\left(\frac{\mathbf{q}}{2}\right)^2 + \left(\frac{\mathbf{p}}{3}\right)^2}}.$$

## 2.7 Inequalities

Variables:  $x, y, z$

Real numbers:  $\begin{cases} a, b, c, d \\ a_1, a_2, a_3, \dots, a_n \end{cases}, m, n$

Determinants:  $D, D_x, D_y, D_z$

### 126. Inequalities, Interval Notations and Graphs

Inequality	Interval Notation	Graph
$a \leq x \leq b$	$[a, b]$	
$a < x \leq b$	$(a, b]$	
$a \leq x < b$	$[a, b)$	
$a < x < b$	$(a, b)$	
$-\infty < x \leq b,$ $x \leq b$	$(-\infty, b]$	
$-\infty < x < b,$ $x < b$	$(-\infty, b)$	
$a \leq x < \infty,$ $x \geq a$	$[a, \infty)$	
$a < x < \infty,$ $x > a$	$(a, \infty)$	

CHAPTER 2. ALGEBRA

- 127.** If  $a > b$ , then  $b < a$ .
- 128.** If  $a > b$ , then  $a - b > 0$  or  $b - a < 0$ .
- 129.** If  $a > b$ , then  $a + c > b + c$ .
- 130.** If  $a > b$ , then  $a - c > b - c$ .
- 131.** If  $a > b$  and  $c > d$ , then  $a + c > b + d$ .
- 132.** If  $a > b$  and  $c > d$ , then  $a - d > b - c$ .
- 133.** If  $a > b$  and  $m > 0$ , then  $ma > mb$ .
- 134.** If  $a > b$  and  $m > 0$ , then  $\frac{a}{m} > \frac{b}{m}$ .
- 135.** If  $a > b$  and  $m < 0$ , then  $ma < mb$ .
- 136.** If  $a > b$  and  $m < 0$ , then  $\frac{a}{m} < \frac{b}{m}$ .
- 137.** If  $0 < a < b$  and  $n > 0$ , then  $a^n < b^n$ .
- 138.** If  $0 < a < b$  and  $n < 0$ , then  $a^n > b^n$ .
- 139.** If  $0 < a < b$ , then  $\sqrt[n]{a} < \sqrt[n]{b}$ .
- 140.**  $\sqrt{ab} \leq \frac{a+b}{2}$ ,  
where  $a > 0$ ,  $b > 0$ ; an equality is valid only if  $a = b$ .
- 141.**  $a + \frac{1}{a} \geq 2$ , where  $a > 0$ ; an equality takes place only at  $a = 1$ .







## CHAPTER 2. ALGEBRA

142.  $\sqrt[n]{a_1 a_2 \dots a_n} \leq \frac{a_1 + a_2 + \dots + a_n}{n}$ , where  $a_1, a_2, \dots, a_n > 0$ .

143. If  $ax + b > 0$  and  $a > 0$ , then  $x > -\frac{b}{a}$ .

144. If  $ax + b > 0$  and  $a < 0$ , then  $x < -\frac{b}{a}$ .

145.  $ax^2 + bx + c > 0$

	$a > 0$	$a < 0$
$D > 0$	 <p><math>x &lt; x_1, x &gt; x_2</math></p>	 <p><math>x_1 &lt; x &lt; x_2</math></p>
$D = 0$	 <p><math>x_1 &lt; x, x &gt; x_1</math></p>	 <p><math>x \in \emptyset</math></p>
$D < 0$	 <p><math>-\infty &lt; x &lt; \infty</math></p>	 <p><math>x \in \emptyset</math></p>

CHAPTER 2. ALGEBRA

146.  $|a + b| \leq |a| + |b|$
147. If  $|x| < a$ , then  $-a < x < a$ , where  $a > 0$ .
148. If  $|x| > a$ , then  $x < -a$  and  $x > a$ , where  $a > 0$ .
149. If  $x^2 < a$ , then  $|x| < \sqrt{a}$ , where  $a > 0$ .
150. If  $x^2 > a$ , then  $|x| > \sqrt{a}$ , where  $a > 0$ .
151. If  $\frac{f(x)}{g(x)} > 0$ , then  $\begin{cases} f(x) \cdot g(x) > 0 \\ g(x) \neq 0 \end{cases}$ .
152.  $\frac{f(x)}{g(x)} < 0$ , then  $\begin{cases} f(x) \cdot g(x) < 0 \\ g(x) \neq 0 \end{cases}$ .

## 2.8 Compound Interest Formulas

Future value: A  
Initial deposit: C  
Annual rate of interest: r  
Number of years invested: t  
Number of times compounded per year: n

153. General Compound Interest Formula

$$A = C \left( 1 + \frac{r}{n} \right)^{nt}$$



CHAPTER 2. ALGEBRA

**154.** Simplified Compound Interest Formula

If interest is compounded once per year, then the previous formula simplifies to:

$$A = C(1+r)^t.$$

**155.** Continuous Compound Interest

If interest is compounded continually ( $n \rightarrow \infty$ ), then

$$A = Ce^{rt}.$$

## Chapter 3

# Geometry

### 3.1 Right Triangle

Legs of a right triangle:  $a, b$

Hypotenuse:  $c$

Altitude:  $h$

Medians:  $m_a, m_b, m_c$

Angles:  $\alpha, \beta$

Radius of circumscribed circle:  $R$

Radius of inscribed circle:  $r$

Area:  $S$




Figure 8.

**156.**  $\alpha + \beta = 90^\circ$

CHAPTER 3. GEOMETRY

157.  $\sin \alpha = \frac{a}{c} = \cos \beta$

158.  $\cos \alpha = \frac{b}{c} = \sin \beta$

159.  $\tan \alpha = \frac{a}{b} = \cot \beta$

160.  $\cot \alpha = \frac{b}{a} = \tan \beta$

161.  $\sec \alpha = \frac{c}{b} = \operatorname{cosec} \beta$

162.  $\operatorname{cosec} \alpha = \frac{c}{a} = \sec \beta$

163. Pythagorean Theorem  
 $a^2 + b^2 = c^2$

164.  $a^2 = fc$ ,  $b^2 = gc$ ,  
where  $f$  and  $c$  are projections of the legs  $a$  and  $b$ , respectively, onto the hypotenuse  $c$ .




Figure 9.

CHAPTER 3. GEOMETRY

165.  $h^2 = fg$ ,  
where  $h$  is the altitude from the right angle.

166.  $m_a^2 = b^2 - \frac{a^2}{4}$ ,  $m_b^2 = a^2 - \frac{b^2}{4}$ ,  
where  $m_a$  and  $m_b$  are the medians to the legs  $a$  and  $b$ .




Figure 10.

167.  $m_c = \frac{c}{2}$ ,  
where  $m_c$  is the median to the hypotenuse  $c$ .

168.  $R = \frac{c}{2} = m_c$

169.  $r = \frac{a + b - c}{2} = \frac{ab}{a + b + c}$

170.  $ab = ch$

CHAPTER 3. GEOMETRY

171.  $S = \frac{ab}{2} = \frac{ch}{2}$

### 3.2 Isosceles Triangle

Base:  $a$   
Legs:  $b$   
Base angle:  $\beta$   
Vertex angle:  $\alpha$   
Altitude to the base:  $h$   
Perimeter:  $L$   
Area:  $S$




Figure 11.

172.  $\beta = 90^\circ - \frac{\alpha}{2}$

173.  $h^2 = b^2 - \frac{a^2}{4}$

CHAPTER 3. GEOMETRY

174.  $L = a + 2b$

175.  $S = \frac{ah}{2} = \frac{b^2}{2} \sin \alpha$

### 3.3 Equilateral Triangle

Side of a equilateral triangle:  $a$   
Altitude:  $h$   
Radius of circumscribed circle:  $R$   
Radius of inscribed circle:  $r$   
Perimeter:  $L$   
Area:  $S$




Figure 12.

176.  $h = \frac{a\sqrt{3}}{2}$

CHAPTER 3. GEOMETRY

$$177. R = \frac{2}{3}h = \frac{a\sqrt{3}}{3}$$

$$178. r = \frac{1}{3}h = \frac{a\sqrt{3}}{6} = \frac{R}{2}$$

$$179. L = 3a$$

$$180. S = \frac{ah}{2} = \frac{a^2\sqrt{3}}{4}$$

### 3.4 Scalene Triangle

(A triangle with no two sides equal)

Sides of a triangle:  $a, b, c$

Semiperimeter:  $p = \frac{a+b+c}{2}$

Angles of a triangle:  $\alpha, \beta, \gamma$

Altitudes to the sides  $a, b, c$ :  $h_a, h_b, h_c$

Medians to the sides  $a, b, c$ :  $m_a, m_b, m_c$

Bisectors of the angles  $\alpha, \beta, \gamma$ :  $t_a, t_b, t_c$

Radius of circumscribed circle:  $R$

Radius of inscribed circle:  $r$

Area:  $S$

CHAPTER 3. GEOMETRY




Figure 13.

181.  $\alpha + \beta + \gamma = 180^\circ$

182.  $a + b > c$ ,  
 $b + c > a$ ,  
 $a + c > b$ .

183.  $|a - b| < c$ ,  
 $|b - c| < a$ ,  
 $|a - c| < b$ .

184. Midline  
 $q = \frac{a}{2}$ ,  $q \parallel a$ .




Figure 14.



## CHAPTER 3. GEOMETRY

**185.** Law of Cosines

$$a^2 = b^2 + c^2 - 2bc \cos \alpha,$$

$$b^2 = a^2 + c^2 - 2ac \cos \beta,$$

$$c^2 = a^2 + b^2 - 2ab \cos \gamma.$$

**186.** Law of Sines

$$\frac{a}{\sin \alpha} = \frac{b}{\sin \beta} = \frac{c}{\sin \gamma} = 2R,$$

where  $R$  is the radius of the circumscribed circle.

$$**187.** \quad R = \frac{a}{2 \sin \alpha} = \frac{b}{2 \sin \beta} = \frac{c}{2 \sin \gamma} = \frac{bc}{2h_a} = \frac{ac}{2h_b} = \frac{ab}{2h_c} = \frac{abc}{4S}$$

$$**188.** \quad r^2 = \frac{(p-a)(p-b)(p-c)}{p},$$

$$\frac{1}{r} = \frac{1}{h_a} + \frac{1}{h_b} + \frac{1}{h_c}.$$

$$**189.** \quad \sin \frac{\alpha}{2} = \sqrt{\frac{(p-b)(p-c)}{bc}},$$

$$\cos \frac{\alpha}{2} = \sqrt{\frac{p(p-a)}{bc}},$$

$$\tan \frac{\alpha}{2} = \sqrt{\frac{(p-b)(p-c)}{p(p-a)}}.$$

$$**190.** \quad h_a = \frac{2}{a} \sqrt{p(p-a)(p-b)(p-c)},$$

$$h_b = \frac{2}{b} \sqrt{p(p-a)(p-b)(p-c)},$$

$$h_c = \frac{2}{c} \sqrt{p(p-a)(p-b)(p-c)}.$$

## CHAPTER 3. GEOMETRY

$$\begin{aligned}
 191. \quad h_a &= b \sin \gamma = c \sin \beta, \\
 h_b &= a \sin \gamma = c \sin \alpha, \\
 h_c &= a \sin \beta = b \sin \alpha.
 \end{aligned}$$

$$\begin{aligned}
 192. \quad m_a^2 &= \frac{b^2 + c^2}{2} - \frac{a^2}{4}, \\
 m_b^2 &= \frac{a^2 + c^2}{2} - \frac{b^2}{4}, \\
 m_c^2 &= \frac{a^2 + b^2}{2} - \frac{c^2}{4}.
 \end{aligned}$$




Figure 15.

$$193. \quad AM = \frac{2}{3}m_a, \quad BM = \frac{2}{3}m_b, \quad CM = \frac{2}{3}m_c \quad (\text{Fig.15}).$$

$$\begin{aligned}
 194. \quad t_a^2 &= \frac{4bcp(p-a)}{(b+c)^2}, \\
 t_b^2 &= \frac{4acp(p-b)}{(a+c)^2}, \\
 t_c^2 &= \frac{4abp(p-c)}{(a+b)^2}.
 \end{aligned}$$

CHAPTER 3. GEOMETRY

195.  $S = \frac{ah_a}{2} = \frac{bh_b}{2} = \frac{ch_c}{2},$   
 $S = \frac{ab \sin \gamma}{2} = \frac{ac \sin \beta}{2} = \frac{bc \sin \alpha}{2},$   
 $S = \sqrt{p(p-a)(p-b)(p-c)}$  (Heron's Formula),  
 $S = pr,$   
 $S = \frac{abc}{4R},$   
 $S = 2R^2 \sin \alpha \sin \beta \sin \gamma,$   
 $S = p^2 \tan \frac{\alpha}{2} \tan \frac{\beta}{2} \tan \frac{\gamma}{2}.$

### 3.5 Square

Side of a square:  $a$   
Diagonal:  $d$   
Radius of circumscribed circle:  $R$   
Radius of inscribed circle:  $r$   
Perimeter:  $L$   
Area:  $S$




Figure 16.

CHAPTER 3. GEOMETRY

196.  $d = a\sqrt{2}$

197.  $R = \frac{d}{2} = \frac{a\sqrt{2}}{2}$

198.  $r = \frac{a}{2}$

199.  $L = 4a$

200.  $S = a^2$

### 3.6 Rectangle

Sides of a rectangle:  $a, b$

Diagonal:  $d$

Radius of circumscribed circle:  $R$

Perimeter:  $L$

Area:  $S$




Figure 17.

201.  $d = \sqrt{a^2 + b^2}$

CHAPTER 3. GEOMETRY

202.  $R = \frac{d}{2}$

203.  $L = 2(a + b)$

204.  $S = ab$

### 3.7 Parallelogram

Sides of a parallelogram:  $a, b$

Diagonals:  $d_1, d_2$

Consecutive angles:  $\alpha, \beta$

Angle between the diagonals:  $\varphi$

Altitude:  $h$

Perimeter:  $L$

Area:  $S$




Figure 18.

205.  $\alpha + \beta = 180^\circ$

206.  $d_1^2 + d_2^2 = 2(a^2 + b^2)$

CHAPTER 3. GEOMETRY

207.  $h = b \sin \alpha = b \sin \beta$

208.  $L = 2(a + b)$

209.  $S = ah = ab \sin \alpha$  ,  
 $S = \frac{1}{2}d_1d_2 \sin \varphi$  .

### 3.8 Rhombus

Side of a rhombus:  $a$

Diagonals:  $d_1, d_2$

Consecutive angles:  $\alpha, \beta$

Altitude:  $H$

Radius of inscribed circle:  $r$

Perimeter:  $L$

Area:  $S$




Figure 19.

CHAPTER 3. GEOMETRY

**210.**  $\alpha + \beta = 180^\circ$

**211.**  $d_1^2 + d_2^2 = 4a^2$

**212.**  $h = a \sin \alpha = \frac{d_1 d_2}{2a}$

**213.**  $r = \frac{h}{2} = \frac{d_1 d_2}{4a} = \frac{a \sin \alpha}{2}$

**214.**  $L = 4a$

**215.**  $S = ah = a^2 \sin \alpha ,$   
 $S = \frac{1}{2} d_1 d_2 .$

### 3.9 Trapezoid

Bases of a trapezoid: a, b

Midline: q

Altitude: h

Area: S

CHAPTER 3. GEOMETRY




Figure 20.

216.  $q = \frac{a+b}{2}$

217.  $S = \frac{a+b}{2} \cdot h = qh$

### 3.10 Isosceles Trapezoid

Bases of a trapezoid:  $a, b$   
Leg:  $c$   
Midline:  $q$   
Altitude:  $h$   
Diagonal:  $d$   
Radius of circumscribed circle:  $R$   
Area:  $S$



CHAPTER 3. GEOMETRY




Figure 21.

218.  $q = \frac{a+b}{2}$

219.  $d = \sqrt{ab+c^2}$

220.  $h = \sqrt{c^2 - \frac{1}{4}(b-a)^2}$

221.  $R = \frac{c\sqrt{ab+c^2}}{\sqrt{(2c-a+b)(2c+a-b)}}$

222.  $S = \frac{a+b}{2} \cdot h = qh$

CHAPTER 3. GEOMETRY

### 3.11 Isosceles Trapezoid with Inscribed Circle

Bases of a trapezoid:  $a, b$   
Leg:  $c$   
Midline:  $q$   
Altitude:  $h$   
Diagonal:  $d$   
Radius of inscribed circle:  $R$   
Radius of circumscribed circle:  $r$   
Perimeter:  $L$   
Area:  $S$




Figure 22.

**223.**  $a + b = 2c$

**224.**  $q = \frac{a+b}{2} = c$

**225.**  $d^2 = h^2 + c^2$

CHAPTER 3. GEOMETRY

$$226. \quad r = \frac{h}{2} = \frac{\sqrt{ab}}{2}$$

$$227. \quad R = \frac{cd}{2h} = \frac{cd}{4r} = \frac{c}{2} \sqrt{1 + \frac{c^2}{ab}} = \frac{c}{2h} \sqrt{h^2 + c^2} = \frac{a+b}{8} \sqrt{\frac{a}{b} + 6 + \frac{b}{a}}$$

$$228. \quad L = 2(a+b) = 4c$$

$$229. \quad S = \frac{a+b}{2} \cdot h = \frac{(a+b)\sqrt{ab}}{2} = qh = ch = \frac{Lr}{2}$$

### 3.12 Trapezoid with Inscribed Circle

Bases of a trapezoid:  $a, b$

Lateral sides:  $c, d$

Midline:  $q$

Altitude:  $h$

Diagonals:  $d_1, d_2$

Angle between the diagonals:  $\varphi$

Radius of inscribed circle:  $r$

Radius of circumscribed circle:  $R$

Perimeter:  $L$

Area:  $S$

CHAPTER 3. GEOMETRY




Figure 23.

230.  $a + b = c + d$

231.  $q = \frac{a + b}{2} = \frac{c + d}{2}$

232.  $L = 2(a + b) = 2(c + d)$

233.  $S = \frac{a + b}{2} \cdot h = \frac{c + d}{2} \cdot h = qh,$   
 $S = \frac{1}{2}d_1d_2 \sin \varphi.$

### 3.13 Kite

Sides of a kite:  $a, b$

Diagonals:  $d_1, d_2$

Angles:  $\alpha, \beta, \gamma$

Perimeter:  $L$

Area:  $S$

CHAPTER 3. GEOMETRY




Figure 24.

234.  $\alpha + \beta + 2\gamma = 360^\circ$

235.  $L = 2(a + b)$

236.  $S = \frac{d_1 d_2}{2}$

### 3.14 Cyclic Quadrilateral

Sides of a quadrilateral:  $a, b, c, d$

Diagonals:  $d_1, d_2$

Angle between the diagonals:  $\varphi$

Internal angles:  $\alpha, \beta, \gamma, \delta$

Radius of circumscribed circle:  $R$

Perimeter:  $L$

Semiperimeter:  $p$

Area:  $S$

## CHAPTER 3. GEOMETRY




Figure 25.

237.  $\alpha + \gamma = \beta + \delta = 180^\circ$

238. Ptolemy's Theorem  
 $ac + bd = d_1 d_2$

239.  $L = a + b + c + d$

240. 
$$R = \frac{1}{4} \sqrt{\frac{(ac + bd)(ad + bc)(ab + cd)}{(p - a)(p - b)(p - c)(p - d)}}$$
,

where  $p = \frac{L}{2}$ .

241.  $S = \frac{1}{2} d_1 d_2 \sin \varphi$ ,

$$S = \sqrt{(p - a)(p - b)(p - c)(p - d)},$$

where  $p = \frac{L}{2}$ .

### 3.15 Tangential Quadrilateral

Sides of a quadrilateral:  $a, b, c, d$

Diagonals:  $d_1, d_2$

Angle between the diagonals:  $\varphi$

Radius of inscribed circle:  $r$

Perimeter:  $L$

Semiperimeter:  $p$

Area:  $S$




Figure 26.

242.  $a + c = b + d$

243.  $L = a + b + c + d = 2(a + c) = 2(b + d)$

244. 
$$r = \frac{\sqrt{d_1^2 d_2^2 - (a - b)^2 (a + b - p)^2}}{2p},$$

where  $p = \frac{L}{2}$ .

CHAPTER 3. GEOMETRY

245.  $S = pr = \frac{1}{2}d_1d_2 \sin \varphi$

### 3.16 General Quadrilateral

Sides of a quadrilateral:  $a, b, c, d$

Diagonals:  $d_1, d_2$

Angle between the diagonals:  $\varphi$

Internal angles:  $\alpha, \beta, \gamma, \delta$

Perimeter:  $L$

Area:  $S$




Figure 27.

246.  $\alpha + \beta + \gamma + \delta = 360^\circ$

247.  $L = a + b + c + d$



CHAPTER 3. GEOMETRY

248.  $S = \frac{1}{2}d_1d_2 \sin \varphi$

### 3.17 Regular Hexagon

Side:  $a$   
Internal angle:  $\alpha$   
Slant height:  $m$   
Radius of inscribed circle:  $r$   
Radius of circumscribed circle:  $R$   
Perimeter:  $L$   
Semiperimeter:  $p$   
Area:  $S$




Figure 28.

249.  $\alpha = 120^\circ$

250.  $r = m = \frac{a\sqrt{3}}{2}$

CHAPTER 3. GEOMETRY

251.  $R = a$

252.  $L = 6a$

253.  $S = pr = \frac{a^2 3\sqrt{3}}{2}$ ,  
where  $p = \frac{L}{2}$ .

### 3.18 Regular Polygon

Side:  $a$

Number of sides:  $n$

Internal angle:  $\alpha$

Slant height:  $m$

Radius of inscribed circle:  $r$

Radius of circumscribed circle:  $R$

Perimeter:  $L$

Semiperimeter:  $p$

Area:  $S$

CHAPTER 3. GEOMETRY




Figure 29.

254.  $\alpha = \frac{n-2}{2} \cdot 180^\circ$

255.  $\alpha = \frac{n-2}{2} \cdot 180^\circ$

256.  $R = \frac{a}{2 \sin \frac{\pi}{n}}$

257.  $r = m = \frac{a}{2 \tan \frac{\pi}{n}} = \sqrt{R^2 - \frac{a^2}{4}}$

258.  $L = na$

259.  $S = \frac{nR^2}{2} \sin \frac{2\pi}{n},$   
 $S = pr = p \sqrt{R^2 - \frac{a^2}{4}},$

CHAPTER 3. GEOMETRY

where  $p = \frac{L}{2}$ .

### 3.19 Circle

Radius:  $R$   
Diameter:  $d$   
Chord:  $a$   
Secant segments:  $e, f$   
Tangent segment:  $g$   
Central angle:  $\alpha$   
Inscribed angle:  $\beta$   
Perimeter:  $L$   
Area:  $S$

**260.**  $a = 2R \sin \frac{\alpha}{2}$




Figure 30.

CHAPTER 3. GEOMETRY

261.  $a_1a_2 = b_1b_2$




Figure 31.

262.  $ee_1 = ff_1$




Figure 32.

263.  $g^2 = ff_1$

CHAPTER 3. GEOMETRY




Figure 33.

264.  $\beta = \frac{\alpha}{2}$




Figure 34.

265.  $L = 2\pi R = \pi d$

266.  $S = \pi R^2 = \frac{\pi d^2}{4} = \frac{LR}{2}$

### 3.20 Sector of a Circle

Radius of a circle:  $R$

Arc length:  $s$

Central angle (in radians):  $x$

Central angle (in degrees):  $\alpha$

Perimeter:  $L$

Area:  $S$




Figure 35.

267.  $s = Rx$

268.  $s = \frac{\pi R \alpha}{180^\circ}$

269.  $L = s + 2R$

270.  $S = \frac{Rs}{2} = \frac{R^2 x}{2} = \frac{\pi R^2 \alpha}{360^\circ}$

### 3.21 Segment of a Circle

Radius of a circle:  $R$

Arc length:  $s$

Chord:  $a$

Central angle (in radians):  $x$

Central angle (in degrees):  $\alpha$

Height of the segment:  $h$

Perimeter:  $L$

Area:  $S$




Figure 36.

271.  $a = 2\sqrt{2hR - h^2}$

272.  $h = R - \frac{1}{2}\sqrt{4R^2 - a^2}$ ,  $h < R$

273.  $L = s + a$



CHAPTER 3. GEOMETRY

$$274. S = \frac{1}{2}[sR - a(R - h)] = \frac{R^2}{2} \left( \frac{\alpha\pi}{180^\circ} - \sin \alpha \right) = \frac{R^2}{2} (x - \sin x),$$
$$S \approx \frac{2}{3} ha.$$

### 3.22 Cube

Edge:  $a$

Diagonal:  $d$

Radius of inscribed sphere:  $r$

Radius of circumscribed sphere:  $r$

Surface area:  $S$

Volume:  $V$




Figure 37.

$$275. d = a\sqrt{3}$$

$$276. r = \frac{a}{2}$$

CHAPTER 3. GEOMETRY

277.  $R = \frac{a\sqrt{3}}{2}$

278.  $S = 6a^2$

279.  $V = a^3$

### 3.23 Rectangular Parallelepiped

Edges:  $a, b, c$   
Diagonal:  $d$   
Surface area:  $S$   
Volume:  $V$




Figure 38.

280.  $d = \sqrt{a^2 + b^2 + c^2}$

281.  $S = 2(ab + ac + bc)$

282.  $V = abc$

### 3.24 Prism

Lateral edge:  $l$   
Height:  $h$   
Lateral area:  $S_L$   
Area of base:  $S_B$   
Total surface area:  $S$   
Volume:  $V$




Figure 39.

**283.**  $S = S_L + 2S_B.$

**284.** Lateral Area of a Right Prism  
 $S_L = (a_1 + a_2 + a_3 + \dots + a_n)l$

**285.** Lateral Area of an Oblique Prism  
 $S_L = pl,$   
where  $p$  is the perimeter of the cross section.

CHAPTER 3. GEOMETRY

286.  $V = S_B h$

287. Cavalieri's Principle

Given two solids included between parallel planes. If every plane cross section parallel to the given planes has the same area in both solids, then the volumes of the solids are equal.

### 3.25 Regular Tetrahedron

Triangle side length:  $a$

Height:  $h$

Area of base:  $S_B$

Surface area:  $S$

Volume:  $V$




Figure 40.

288.  $h = \sqrt{\frac{2}{3}} a$

CHAPTER 3. GEOMETRY

289.  $S_B = \frac{\sqrt{3}a^2}{4}$

290.  $S = \sqrt{3}a^2$

291.  $V = \frac{1}{3}S_B h = \frac{a^3}{6\sqrt{2}}$ .

### 3.26 Regular Pyramid

Side of base:  $a$

Lateral edge:  $b$

Height:  $h$

Slant height:  $m$

Number of sides:  $n$

Semiperimeter of base:  $p$

Radius of inscribed sphere of base:  $r$

Area of base:  $S_B$

Lateral surface area:  $S_L$

Total surface area:  $S$

Volume:  $V$

CHAPTER 3. GEOMETRY




Figure 41.

$$292. \quad m = \sqrt{b^2 - \frac{a^2}{4}}$$

$$293. \quad h = \frac{\sqrt{4b^2 \sin^2 \frac{\pi}{n} - a^2}}{2 \sin \frac{\pi}{n}}$$

$$294. \quad S_L = \frac{1}{2} nam = \frac{1}{4} na \sqrt{4b^2 - a^2} = pm$$

$$295. \quad S_B = pr$$

$$296. \quad S = S_B + S_L$$

$$297. \quad V = \frac{1}{3} S_B h = \frac{1}{3} prh$$

CHAPTER 3. GEOMETRY

### 3.27 Frustum of a Regular Pyramid

Base and top side lengths:  $\begin{cases} a_1, a_2, a_3, \dots, a_n \\ b_1, b_2, b_3, \dots, b_n \end{cases}$

Height:  $h$

Slant height:  $m$

Area of bases:  $S_1, S_2$

Lateral surface area:  $S_L$

Perimeter of bases:  $P_1, P_2$

Scale factor:  $k$

Total surface area:  $S$

Volume:  $V$




Figure 42.

298.  $\frac{b_1}{a_1} = \frac{b_2}{a_2} = \frac{b_3}{a_3} = \dots = \frac{b_n}{a_n} = \frac{b}{a} = k$

CHAPTER 3. GEOMETRY

$$299. \frac{S_2}{S_1} = k^2$$

$$300. S_L = \frac{m(P_1 + P_2)}{2}$$

$$301. S = S_L + S_1 + S_2$$

$$302. V = \frac{h}{3}(S_1 + \sqrt{S_1 S_2} + S_2)$$

$$303. V = \frac{hS_1}{3} \left[ 1 + \frac{b}{a} + \left( \frac{b}{a} \right)^2 \right] = \frac{hS_1}{3} [1 + k + k^2]$$

### 3.28 Rectangular Right Wedge

Sides of base: a, b

Top edge: c

Height: h

Lateral surface area:  $S_L$

Area of base:  $S_B$

Total surface area: S

Volume: V



CHAPTER 3. GEOMETRY




Figure 43.

304.  $S_L = \frac{1}{2}(a + c)\sqrt{4h^2 + b^2} + b\sqrt{h^2 + (a - c)^2}$

305.  $S_B = ab$

306.  $S = S_B + S_L$

307.  $V = \frac{bh}{6}(2a + c)$

### 3.29 Platonic Solids

Edge:  $a$

Radius of inscribed circle:  $r$

Radius of circumscribed circle:  $R$

Surface area:  $S$

Volume:  $V$

CHAPTER 3. GEOMETRY

**308.** Five Platonic Solids

The platonic solids are convex polyhedra with equivalent faces composed of congruent convex regular polygons.

Solid	Number of Vertices	Number of Edges	Number of Faces	Section
Tetrahedron	4	6	4	3.25
Cube	8	12	6	3.22
Octahedron	6	12	8	3.27
Icosahedron	12	30	20	3.27
Dodecahedron	20	30	12	3.27

Octahedron




Figure 44.

**309.**  $r = \frac{a\sqrt{6}}{6}$

**310.**  $R = \frac{a\sqrt{2}}{2}$

CHAPTER 3. GEOMETRY

311.  $S = 2a^2\sqrt{3}$

312.  $V = \frac{a^3\sqrt{2}}{3}$

Icosahedron




Figure 45.

313.  $r = \frac{a\sqrt{3}(3+\sqrt{5})}{12}$

314.  $R = \frac{a}{4}\sqrt{2(5+\sqrt{5})}$

315.  $S = 5a^2\sqrt{3}$

316.  $V = \frac{5a^3(3+\sqrt{5})}{12}$

CHAPTER 3. GEOMETRY

## Dodecahedron




Figure 46.

$$317. \quad r = \frac{a\sqrt{10(25+11\sqrt{5})}}{2}$$

$$318. \quad R = \frac{a\sqrt{3}(1+\sqrt{5})}{4}$$

$$319. \quad S = 3a^2\sqrt{5(5+2\sqrt{5})}$$

$$320. \quad V = \frac{a^3(15+7\sqrt{5})}{4}$$

### 3.30 Right Circular Cylinder

Radius of base:  $R$   
Diameter of base:  $d$

CHAPTER 3. GEOMETRY

Height:  $H$   
Lateral surface area:  $S_L$   
Area of base:  $S_B$   
Total surface area:  $S$   
Volume:  $V$




Figure 47.

**321.**  $S_L = 2\pi RH$

**322.**  $S = S_L + 2S_B = 2\pi R(H + R) = \pi d \left( H + \frac{d}{2} \right)$

**323.**  $V = S_B H = \pi R^2 H$

CHAPTER 3. GEOMETRY

### 3.31 Right Circular Cylinder with an Oblique Plane Face

Radius of base:  $R$

The greatest height of a side:  $h_1$

The shortest height of a side:  $h_2$

Lateral surface area:  $S_L$

Area of plane end faces:  $S_B$

Total surface area:  $S$

Volume:  $V$




Figure 48.

**324.**  $S_L = \pi R(h_1 + h_2)$

**325.**  $S_B = \pi R^2 + \pi R \sqrt{R^2 + \left(\frac{h_1 - h_2}{2}\right)^2}$

CHAPTER 3. GEOMETRY

$$326. S = S_L + S_B = \pi R \left[ h_1 + h_2 + R + \sqrt{R^2 + \left( \frac{h_1 - h_2}{2} \right)^2} \right]$$

$$327. V = \frac{\pi R^2}{2} (h_1 + h_2)$$

### 3.32 Right Circular Cone

Radius of base:  $R$   
Diameter of base:  $d$   
Height:  $H$   
Slant height:  $m$   
Lateral surface area:  $S_L$   
Area of base:  $S_B$   
Total surface area:  $S$   
Volume:  $V$




Figure 49.

CHAPTER 3. GEOMETRY

$$328. H = \sqrt{m^2 - R^2}$$

$$329. S_L = \pi R m = \frac{\pi m d}{2}$$

$$330. S_B = \pi R^2$$

$$331. S = S_L + S_B = \pi R(m + R) = \frac{1}{2} \pi d \left( m + \frac{d}{2} \right)$$

$$332. V = \frac{1}{3} S_B H = \frac{1}{3} \pi R^2 H$$

### 3.33 Frustum of a Right Circular Cone

Radius of bases:  $R, r$

Height:  $H$

Slant height:  $m$

Scale factor:  $k$

Area of bases:  $S_1, S_2$

Lateral surface area:  $S_L$

Total surface area:  $S$

Volume:  $V$



CHAPTER 3. GEOMETRY




Figure 50.

333.  $H = \sqrt{m^2 - (R - r)^2}$

334.  $\frac{R}{r} = k$

335.  $\frac{S_2}{S_1} = \frac{R^2}{r^2} = k^2$

336.  $S_L = \pi m(R + r)$

337.  $S = S_1 + S_2 + S_L = \pi[R^2 + r^2 + m(R + r)]$

338.  $V = \frac{h}{3}(S_1 + \sqrt{S_1 S_2} + S_2)$

339.  $V = \frac{hS_1}{3} \left[ 1 + \frac{R}{r} + \left( \frac{R}{r} \right)^2 \right] = \frac{hS_1}{3} [1 + k + k^2]$

### 3.34 Sphere

Radius:  $R$   
Diameter:  $d$   
Surface area:  $S$   
Volume:  $V$




Figure 51.

**340.**  $S = 4\pi R^2$

**341.**  $V = \frac{4}{3}\pi R^3 = \frac{1}{6}\pi d^3 = \frac{1}{3}SR$

### 3.35 Spherical Cap

Radius of sphere:  $R$   
Radius of base:  $r$   
Height:  $h$   
Area of plane face:  $S_B$   
Area of spherical cap:  $S_C$   
Total surface area:  $S$   
Volume:  $V$

CHAPTER 3. GEOMETRY




Figure 52.

342.  $R = \frac{r^2 + h^2}{2h}$

343.  $S_B = \pi r^2$

344.  $S_C = \pi(h^2 + r^2)$

345.  $S = S_B + S_C = \pi(h^2 + 2r^2) = \pi(2Rh + r^2)$

346.  $V = \frac{\pi}{6}h^2(3R - h) = \frac{\pi}{6}h(3r^2 + h^2)$

### 3.36 Spherical Sector

Radius of sphere:  $R$

Radius of base of spherical cap:  $r$

Height:  $h$

Total surface area:  $S$

Volume:  $V$

CHAPTER 3. GEOMETRY




Figure 53.

347.  $S = \pi R(2h + r)$

348.  $V = \frac{2}{3}\pi R^2 h$

Note: The given formulas are correct both for “open” and “closed” spherical sector.

### 3.37 Spherical Segment

Radius of sphere:  $R$

Radius of bases:  $r_1, r_2$

Height:  $h$

Area of spherical surface:  $S_s$

Area of plane end faces:  $S_1, S_2$

Total surface area:  $S$

Volume:  $V$

CHAPTER 3. GEOMETRY




Figure 54.

349.  $S_s = 2\pi Rh$

350.  $S = S_s + S_1 + S_2 = \pi(2Rh + r_1^2 + r_2^2)$

351.  $V = \frac{1}{6}\pi h(3r_1^2 + 3r_2^2 + h^2)$

### 3.38 Spherical Wedge

Radius:  $R$

Dihedral angle in degrees:  $x$

Dihedral angle in radians:  $\alpha$

Area of spherical lune:  $S_l$

Total surface area:  $S$

Volume:  $V$

CHAPTER 3. GEOMETRY




Figure 55.

$$352. S_L = \frac{\pi R^2}{90} \alpha = 2R^2 x$$

$$353. S = \pi R^2 + \frac{\pi R^2}{90} \alpha = \pi R^2 + 2R^2 x$$

$$354. V = \frac{\pi R^3}{270} \alpha = \frac{2}{3} R^3 x$$

### 3.39 Ellipsoid

Semi-axes:  $a, b, c$

Volume:  $V$

CHAPTER 3. GEOMETRY




Figure 56.

**355.**  $V = \frac{4}{3}\pi abc$

### Prolate Spheroid

Semi-axes:  $a, b, b$  ( $a > b$ )

Surface area:  $S$

Volume:  $V$

**356.**  $S = 2\pi b \left( b + \frac{a \arcsin e}{e} \right),$

where  $e = \frac{\sqrt{a^2 - b^2}}{a}.$

**357.**  $V = \frac{4}{3}\pi b^2 a$

CHAPTER 3. GEOMETRY

## Oblate Spheroid

Semi-axes:  $a, b, b$  ( $a < b$ )

Surface area:  $S$

Volume:  $V$

$$358. \quad S = 2\pi b \left( b + \frac{a \operatorname{arcsinh} \left( \frac{be}{a} \right)}{be/a} \right),$$

$$\text{where } e = \frac{\sqrt{b^2 - a^2}}{b}.$$

$$359. \quad V = \frac{4}{3}\pi b^2 a$$

### 3.40 Circular Torus

Major radius:  $R$


Minor radius:  $r$

Surface area:  $S$

Volume:  $V$



CHAPTER 3. GEOMETRY



Picture 57.

**360.**  $S = 4\pi^2 Rr$

**361.**  $V = 2\pi^2 Rr^2$

## Chapter 4

# Trigonometry

Angles:  $\alpha, \beta$

Real numbers (coordinates of a point):  $x, y$

Whole number:  $k$

### 4.1 Radian and Degree Measures of Angles

**362.**  $1 \text{ rad} = \frac{180^\circ}{\pi} \approx 57^\circ 17' 45''$

**363.**  $1^\circ = \frac{\pi}{180} \text{ rad} \approx 0.017453 \text{ rad}$

**364.**  $1' = \frac{\pi}{180 \cdot 60} \text{ rad} \approx 0.000291 \text{ rad}$

**365.**  $1'' = \frac{\pi}{180 \cdot 3600} \text{ rad} \approx 0.000005 \text{ rad}$

**366.**

Angle (degrees)	0	30	45	60	90	180	270	360
Angle (radians)	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\pi$	$\frac{3\pi}{2}$	$2\pi$

CHAPTER 4. TRIGONOMETRY

## 4.2 Definitions and Graphs of Trigonometric Functions




Figure 58.

**367.**  $\sin \alpha = \frac{y}{r}$

**368.**  $\cos \alpha = \frac{x}{r}$

**369.**  $\tan \alpha = \frac{y}{x}$

**370.**  $\cot \alpha = \frac{x}{y}$

CHAPTER 4. TRIGONOMETRY

371.  $\sec \alpha = \frac{r}{x}$

372.  $\operatorname{cosec} \alpha = \frac{r}{y}$

373. Sine Function  
 $y = \sin x, -1 \leq \sin x \leq 1.$




Figure 59.

374. Cosine Function  
 $y = \cos x, -1 \leq \cos x \leq 1.$

CHAPTER 4. TRIGONOMETRY




Figure 60.

**375.** Tangent Function

$$y = \tan x, \quad x \neq (2k+1)\frac{\pi}{2}, \quad -\infty \leq \tan x \leq \infty.$$




Figure 61.

CHAPTER 4. TRIGONOMETRY

**376.** Cotangent Function

$$y = \cot x, \quad x \neq k\pi, \quad -\infty \leq \cot x \leq \infty.$$




Figure 62.

**377.** Secant Function

$$y = \sec x, \quad x \neq (2k+1)\frac{\pi}{2}.$$

CHAPTER 4. TRIGONOMETRY




Figure 63.

**378.** Cosecant Function  
 $y = \operatorname{cosec} x, x \neq k\pi.$




Figure 64.

CHAPTER 4. TRIGONOMETRY

### 4.3. Signs of Trigonometric Functions

379.

Quadrant	Sin $\alpha$	Cos $\alpha$	Tan $\alpha$	Cot $\alpha$	Sec $\alpha$	Cosec $\alpha$
I	+	+	+	+	+	+
II	+					+
III			+	+		
IV		+			+	

380.




Figure 65.



## CHAPTER 4. TRIGONOMETRY

## 4.4 Trigonometric Functions of Common Angles

381.

$\alpha^\circ$	$\alpha$ rad	$\sin \alpha$	$\cos \alpha$	$\tan \alpha$	$\cot \alpha$	$\sec \alpha$	$\operatorname{cosec} \alpha$
0	0	0	1	0	$\infty$	1	$\infty$
30	$\frac{\pi}{6}$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{3}}$	$\sqrt{3}$	$\frac{2}{\sqrt{3}}$	2
45	$\frac{\pi}{4}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1	1	$\sqrt{2}$	$\sqrt{2}$
60	$\frac{\pi}{3}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$	$\frac{1}{\sqrt{3}}$	2	$\frac{2}{\sqrt{3}}$
90	$\frac{\pi}{2}$	1	0	$\infty$	0	$\infty$	1
120	$\frac{2\pi}{3}$	$\frac{\sqrt{3}}{2}$	$-\frac{1}{2}$	$-\sqrt{3}$	$-\frac{1}{\sqrt{3}}$	-2	$\frac{2}{\sqrt{3}}$
180	$\pi$	0	-1	0	$\infty$	-1	$\infty$
270	$\frac{3\pi}{2}$	-1	0	$\infty$	0	$\infty$	-1
360	$2\pi$	0	1	0	$\infty$	1	$\infty$

## CHAPTER 4. TRIGONOMETRY

**382.**

$\alpha^\circ$	$\alpha$ rad	$\sin \alpha$	$\cos \alpha$	$\tan \alpha$	$\cot \alpha$
15	$\frac{\pi}{12}$	$\frac{\sqrt{6}-\sqrt{2}}{4}$	$\frac{\sqrt{6}+\sqrt{2}}{4}$	$2-\sqrt{3}$	$2+\sqrt{3}$
18	$\frac{\pi}{10}$	$\frac{\sqrt{5}-1}{4}$	$\frac{\sqrt{10+2\sqrt{5}}}{4}$	$\sqrt{\frac{5-2\sqrt{5}}{5}}$	$\sqrt{5+2\sqrt{5}}$
36	$\frac{\pi}{5}$	$\frac{\sqrt{10-2\sqrt{5}}}{4}$	$\frac{\sqrt{5}+1}{4}$	$\frac{\sqrt{10-2\sqrt{5}}}{\sqrt{5}+1}$	$\frac{\sqrt{5}+1}{\sqrt{10-2\sqrt{5}}}$
54	$\frac{3\pi}{10}$	$\frac{\sqrt{5}+1}{4}$	$\frac{\sqrt{10-2\sqrt{5}}}{4}$	$\frac{\sqrt{5}+1}{\sqrt{10-2\sqrt{5}}}$	$\frac{\sqrt{10-2\sqrt{5}}}{\sqrt{5}+1}$
72	$\frac{2\pi}{5}$	$\frac{\sqrt{10+2\sqrt{5}}}{4}$	$\frac{\sqrt{5}-1}{4}$	$\sqrt{5+2\sqrt{5}}$	$\sqrt{\frac{5-2\sqrt{5}}{5}}$
75	$\frac{5\pi}{12}$	$\frac{\sqrt{6}+\sqrt{2}}{4}$	$\frac{\sqrt{6}-\sqrt{2}}{4}$	$2+\sqrt{3}$	$2-\sqrt{3}$

## 4.5 Most Important Formulas

**383.**  $\sin^2 \alpha + \cos^2 \alpha = 1$

**384.**  $\sec^2 \alpha - \tan^2 \alpha = 1$

**385.**  $\csc^2 \alpha - \cot^2 \alpha = 1$

**386.**  $\tan \alpha = \frac{\sin \alpha}{\cos \alpha}$

CHAPTER 4. TRIGONOMETRY

387.  $\cot \alpha = \frac{\cos \alpha}{\sin \alpha}$

388.  $\tan \alpha \cdot \cot \alpha = 1$

389.  $\sec \alpha = \frac{1}{\cos \alpha}$

390.  $\operatorname{cosec} \alpha = \frac{1}{\sin \alpha}$

### 4.6 Reduction Formulas

391.

$\beta$	$\sin \beta$	$\cos \beta$	$\tan \beta$	$\cot \beta$
$-\alpha$	$-\sin \alpha$	$+\cos \alpha$	$-\tan \alpha$	$-\cot \alpha$
$90^\circ - \alpha$	$+\cos \alpha$	$+\sin \alpha$	$+\cot \alpha$	$+\tan \alpha$
$90^\circ + \alpha$	$+\cos \alpha$	$-\sin \alpha$	$-\cot \alpha$	$-\tan \alpha$
$180^\circ - \alpha$	$+\sin \alpha$	$-\cos \alpha$	$-\tan \alpha$	$-\cot \alpha$
$180^\circ + \alpha$	$-\sin \alpha$	$-\cos \alpha$	$+\tan \alpha$	$+\cot \alpha$
$270^\circ - \alpha$	$-\cos \alpha$	$-\sin \alpha$	$+\cot \alpha$	$+\tan \alpha$
$270^\circ + \alpha$	$-\cos \alpha$	$+\sin \alpha$	$-\cot \alpha$	$-\tan \alpha$
$360^\circ - \alpha$	$-\sin \alpha$	$+\cos \alpha$	$-\tan \alpha$	$-\cot \alpha$
$360^\circ + \alpha$	$+\sin \alpha$	$+\cos \alpha$	$+\tan \alpha$	$+\cot \alpha$

## CHAPTER 4. TRIGONOMETRY

## 4.7 Periodicity of Trigonometric Functions

$$392. \quad \sin(\alpha \pm 2\pi n) = \sin \alpha, \text{ period } 2\pi \text{ or } 360^\circ.$$

$$393. \quad \cos(\alpha \pm 2\pi n) = \cos \alpha, \text{ period } 2\pi \text{ or } 360^\circ.$$

$$394. \quad \tan(\alpha \pm \pi n) = \tan \alpha, \text{ period } \pi \text{ or } 180^\circ.$$

$$395. \quad \cot(\alpha \pm \pi n) = \cot \alpha, \text{ period } \pi \text{ or } 180^\circ.$$

## 4.8 Relations between Trigonometric Functions

$$396. \quad \sin \alpha = \pm \sqrt{1 - \cos^2 \alpha} = \pm \sqrt{\frac{1}{2}(1 - \cos 2\alpha)} = 2 \cos^2 \left( \frac{\alpha}{2} - \frac{\pi}{4} \right) - 1$$

$$= \frac{2 \tan \frac{\alpha}{2}}{1 + \tan^2 \frac{\alpha}{2}}$$

$$397. \quad \cos \alpha = \pm \sqrt{1 - \sin^2 \alpha} = \pm \sqrt{\frac{1}{2}(1 + \cos 2\alpha)} = 2 \cos^2 \frac{\alpha}{2} - 1$$

$$= \frac{1 - \tan^2 \frac{\alpha}{2}}{1 + \tan^2 \frac{\alpha}{2}}$$

$$398. \quad \tan \alpha = \frac{\sin \alpha}{\cos \alpha} = \pm \sqrt{\sec^2 \alpha - 1} = \frac{\sin 2\alpha}{1 + \cos 2\alpha} = \frac{1 - \cos 2\alpha}{\sin 2\alpha}$$

## CHAPTER 4. TRIGONOMETRY

$$= \pm \sqrt{\frac{1 - \cos 2\alpha}{1 + \cos 2\alpha}} = \frac{2 \tan \frac{\alpha}{2}}{1 + \tan^2 \frac{\alpha}{2}}$$

$$\begin{aligned} 399. \quad \cot \alpha &= \frac{\cos \alpha}{\sin \alpha} = \pm \sqrt{\csc^2 \alpha - 1} = \frac{1 + \cos 2\alpha}{\sin 2\alpha} = \frac{\sin 2\alpha}{1 - \cos 2\alpha} \\ &= \pm \sqrt{\frac{1 + \cos 2\alpha}{1 - \cos 2\alpha}} = \frac{1 - \tan^2 \frac{\alpha}{2}}{2 \tan \frac{\alpha}{2}} \end{aligned}$$

$$400. \quad \sec \alpha = \frac{1}{\cos \alpha} = \pm \sqrt{1 + \tan^2 \alpha} = \frac{1 + \tan^2 \frac{\alpha}{2}}{1 - \tan^2 \frac{\alpha}{2}}$$

$$401. \quad \csc \alpha = \frac{1}{\sin \alpha} = \pm \sqrt{1 + \cot^2 \alpha} = \frac{1 + \tan^2 \frac{\alpha}{2}}{2 \tan \frac{\alpha}{2}}$$

## 4.9 Addition and Subtraction Formulas

$$402. \quad \sin(\alpha + \beta) = \sin \alpha \cos \beta + \sin \beta \cos \alpha$$

$$403. \quad \sin(\alpha - \beta) = \sin \alpha \cos \beta - \sin \beta \cos \alpha$$

$$404. \quad \cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$405. \quad \cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

CHAPTER 4. TRIGONOMETRY

$$406. \quad \tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta}$$

$$407. \quad \tan(\alpha - \beta) = \frac{\tan \alpha - \tan \beta}{1 + \tan \alpha \tan \beta}$$

$$408. \quad \cot(\alpha + \beta) = \frac{1 - \tan \alpha \tan \beta}{\tan \alpha + \tan \beta}$$

$$409. \quad \cot(\alpha - \beta) = \frac{1 + \tan \alpha \tan \beta}{\tan \alpha - \tan \beta}$$

## 4.10 Double Angle Formulas

$$410. \quad \sin 2\alpha = 2 \sin \alpha \cdot \cos \alpha$$

$$411. \quad \cos 2\alpha = \cos^2 \alpha - \sin^2 \alpha = 1 - 2\sin^2 \alpha = 2\cos^2 \alpha - 1$$

$$412. \quad \tan 2\alpha = \frac{2 \tan \alpha}{1 - \tan^2 \alpha} = \frac{2}{\cot \alpha - \tan \alpha}$$

$$413. \quad \cot 2\alpha = \frac{\cot^2 \alpha - 1}{2 \cot \alpha} = \frac{\cot \alpha - \tan \alpha}{2}$$

CHAPTER 4. TRIGONOMETRY

## 4.11 Multiple Angle Formulas

$$414. \quad \sin 3\alpha = 3\sin \alpha - 4\sin^3 \alpha = 3\cos^2 \alpha \cdot \sin \alpha - \sin^3 \alpha$$

$$415. \quad \sin 4\alpha = 4\sin \alpha \cdot \cos \alpha - 8\sin^3 \alpha \cdot \cos \alpha$$

$$416. \quad \sin 5\alpha = 5\sin \alpha - 20\sin^3 \alpha + 16\sin^5 \alpha$$

$$417. \quad \cos 3\alpha = 4\cos^3 \alpha - 3\cos \alpha = \cos^3 \alpha - 3\cos \alpha \cdot \sin^2 \alpha$$

$$418. \quad \cos 4\alpha = 8\cos^4 \alpha - 8\cos^2 \alpha + 1$$

$$419. \quad \cos 5\alpha = 16\cos^5 \alpha - 20\cos^3 \alpha + 5\cos \alpha$$

$$420. \quad \tan 3\alpha = \frac{3\tan \alpha - \tan^3 \alpha}{1 - 3\tan^2 \alpha}$$

$$421. \quad \tan 4\alpha = \frac{4\tan \alpha - 4\tan^3 \alpha}{1 - 6\tan^2 \alpha + \tan^4 \alpha}$$

$$422. \quad \tan 5\alpha = \frac{\tan^5 \alpha - 10\tan^3 \alpha + 5\tan \alpha}{1 - 10\tan^2 \alpha + 5\tan^4 \alpha}$$

$$423. \quad \cot 3\alpha = \frac{\cot^3 \alpha - 3\cot \alpha}{3\cot^2 \alpha - 1}$$

$$424. \quad \cot 4\alpha = \frac{1 - 6\tan^2 \alpha + \tan^4 \alpha}{4\tan \alpha - 4\tan^3 \alpha}$$

CHAPTER 4. TRIGONOMETRY

$$425. \cot 5\alpha = \frac{1 - 10\tan^2 \alpha + 5\tan^4 \alpha}{\tan^5 \alpha - 10\tan^3 \alpha + 5\tan \alpha}$$

### 4.12 Half Angle Formulas

$$426. \sin \frac{\alpha}{2} = \pm \sqrt{\frac{1 - \cos \alpha}{2}}$$

$$427. \cos \frac{\alpha}{2} = \pm \sqrt{\frac{1 + \cos \alpha}{2}}$$

$$428. \tan \frac{\alpha}{2} = \pm \sqrt{\frac{1 - \cos \alpha}{1 + \cos \alpha}} = \frac{\sin \alpha}{1 + \cos \alpha} = \frac{1 - \cos \alpha}{\sin \alpha} = \csc \alpha - \cot \alpha$$

$$429. \cot \frac{\alpha}{2} = \pm \sqrt{\frac{1 + \cos \alpha}{1 - \cos \alpha}} = \frac{\sin \alpha}{1 - \cos \alpha} = \frac{1 + \cos \alpha}{\sin \alpha} = \csc \alpha + \cot \alpha$$

### 4.13 Half Angle Tangent Identities

$$430. \sin \alpha = \frac{2 \tan \frac{\alpha}{2}}{1 + \tan^2 \frac{\alpha}{2}}$$



CHAPTER 4. TRIGONOMETRY

$$431. \quad \cos \alpha = \frac{1 - \tan^2 \frac{\alpha}{2}}{1 + \tan^2 \frac{\alpha}{2}}$$

$$432. \quad \tan \alpha = \frac{2 \tan \frac{\alpha}{2}}{1 - \tan^2 \frac{\alpha}{2}}$$

$$433. \quad \cot \alpha = \frac{1 - \tan^2 \frac{\alpha}{2}}{2 \tan \frac{\alpha}{2}}$$

#### 4.14 Transforming of Trigonometric Expressions to Product

$$434. \quad \sin \alpha + \sin \beta = 2 \sin \frac{\alpha + \beta}{2} \cos \frac{\alpha - \beta}{2}$$

$$435. \quad \sin \alpha - \sin \beta = 2 \cos \frac{\alpha + \beta}{2} \sin \frac{\alpha - \beta}{2}$$

$$436. \quad \cos \alpha + \cos \beta = 2 \cos \frac{\alpha + \beta}{2} \cos \frac{\alpha - \beta}{2}$$

$$437. \quad \cos \alpha - \cos \beta = -2 \sin \frac{\alpha + \beta}{2} \sin \frac{\alpha - \beta}{2}$$

CHAPTER 4. TRIGONOMETRY

$$438. \quad \tan \alpha + \tan \beta = \frac{\sin(\alpha + \beta)}{\cos \alpha \cdot \cos \beta}$$

$$439. \quad \tan \alpha - \tan \beta = \frac{\sin(\alpha - \beta)}{\cos \alpha \cdot \cos \beta}$$

$$440. \quad \cot \alpha + \cot \beta = \frac{\sin(\beta + \alpha)}{\sin \alpha \cdot \sin \beta}$$

$$441. \quad \cot \alpha - \cot \beta = \frac{\sin(\beta - \alpha)}{\sin \alpha \cdot \sin \beta}$$

$$442. \quad \cos \alpha + \sin \alpha = \sqrt{2} \cos\left(\frac{\pi}{4} - \alpha\right) = \sqrt{2} \sin\left(\frac{\pi}{4} + \alpha\right)$$

$$443. \quad \cos \alpha - \sin \alpha = \sqrt{2} \sin\left(\frac{\pi}{4} - \alpha\right) = \sqrt{2} \cos\left(\frac{\pi}{4} + \alpha\right)$$

$$444. \quad \tan \alpha + \cot \beta = \frac{\cos(\alpha - \beta)}{\cos \alpha \cdot \sin \beta}$$

$$445. \quad \tan \alpha - \cot \beta = -\frac{\cos(\alpha + \beta)}{\cos \alpha \cdot \sin \beta}$$

$$446. \quad 1 + \cos \alpha = 2 \cos^2 \frac{\alpha}{2}$$

$$447. \quad 1 - \cos \alpha = 2 \sin^2 \frac{\alpha}{2}$$

CHAPTER 4. TRIGONOMETRY

$$448. \quad 1 + \sin \alpha = 2 \cos^2 \left( \frac{\pi}{4} - \frac{\alpha}{2} \right)$$

$$449. \quad 1 - \sin \alpha = 2 \sin^2 \left( \frac{\pi}{4} - \frac{\alpha}{2} \right)$$

### 4.15 Transforming of Trigonometric Expressions to Sum

$$450. \quad \sin \alpha \cdot \sin \beta = \frac{\cos(\alpha - \beta) - \cos(\alpha + \beta)}{2}$$

$$451. \quad \cos \alpha \cdot \cos \beta = \frac{\cos(\alpha - \beta) + \cos(\alpha + \beta)}{2}$$

$$452. \quad \sin \alpha \cdot \cos \beta = \frac{\sin(\alpha - \beta) + \sin(\alpha + \beta)}{2}$$

$$453. \quad \tan \alpha \cdot \tan \beta = \frac{\tan \alpha + \tan \beta}{\cot \alpha + \cot \beta}$$

$$454. \quad \cot \alpha \cdot \cot \beta = \frac{\cot \alpha + \cot \beta}{\tan \alpha + \tan \beta}$$

$$455. \quad \tan \alpha \cdot \cot \beta = \frac{\tan \alpha + \cot \beta}{\cot \alpha + \tan \beta}$$

CHAPTER 4. TRIGONOMETRY

### 4.16 Powers of Trigonometric Functions

$$456. \sin^2 \alpha = \frac{1 - \cos 2\alpha}{2}$$

$$457. \sin^3 \alpha = \frac{3 \sin \alpha - \sin 3\alpha}{4}$$

$$458. \sin^4 \alpha = \frac{\cos 4\alpha - 4 \cos 2\alpha + 3}{8}$$

$$459. \sin^5 \alpha = \frac{10 \sin \alpha - 5 \sin 3\alpha + \sin 5\alpha}{16}$$

$$460. \sin^6 \alpha = \frac{10 - 15 \cos 2\alpha + 6 \cos 4\alpha - \cos 6\alpha}{32}$$

$$461. \cos^2 \alpha = \frac{1 + \cos 2\alpha}{2}$$

$$462. \cos^3 \alpha = \frac{3 \cos \alpha + \cos 3\alpha}{4}$$

$$463. \cos^4 \alpha = \frac{\cos 4\alpha + 4 \cos 2\alpha + 3}{8}$$

$$464. \cos^5 \alpha = \frac{10 \cos \alpha + 5 \sin 3\alpha + \cos 5\alpha}{16}$$

$$465. \cos^6 \alpha = \frac{10 + 15 \cos 2\alpha + 6 \cos 4\alpha + \cos 6\alpha}{32}$$

CHAPTER 4. TRIGONOMETRY

## 4.17 Graphs of Inverse Trigonometric Functions

### 466. Inverse Sine Function

$$y = \arcsin x, \quad -1 \leq x \leq 1, \quad -\frac{\pi}{2} \leq \arcsin x \leq \frac{\pi}{2}.$$




Figure 66.

### 467. Inverse Cosine Function

$$y = \arccos x, \quad -1 \leq x \leq 1, \quad 0 \leq \arccos x \leq \pi.$$

CHAPTER 4. TRIGONOMETRY




Figure 67.

**468.** Inverse Tangent Function

$$y = \arctan x, \quad -\infty \leq x \leq \infty, \quad -\frac{\pi}{2} < \arctan x < \frac{\pi}{2}.$$




Figure 68.

CHAPTER 4. TRIGONOMETRY

**469.** Inverse Cotangent Function

$$y = \operatorname{arccot} x, \quad -\infty \leq x \leq \infty, \quad 0 < \operatorname{arccot} x < \pi.$$




Figure 69.

**470.** Inverse Secant Function

$$y = \operatorname{arcsec} x, \quad x \in (-\infty, -1] \cup [1, \infty), \quad \operatorname{arcsec} x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right].$$




Figure 70.

CHAPTER 4. TRIGONOMETRY

471. Inverse Cosecant Function

$$y = \operatorname{arccsc} x, \quad x \in (-\infty, -1] \cup [1, \infty), \quad \operatorname{arccsc} x \in \left[-\frac{\pi}{2}, 0\right) \cup \left(0, \frac{\pi}{2}\right].$$




Figure 71.

4.18 Principal Values of Inverse Trigonometric Functions

472.

x	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1
arcsin x	0°	30°	45°	60°	90°
arccos x	90°	60°	45°	30°	0°
x	$-\frac{1}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{3}}{2}$	-1	
arcsin x	-30°	-45°	-60°	-90°	
arccos x	120°	135°	150°	180°	



## CHAPTER 4. TRIGONOMETRY

**473.**

$x$	0	$\frac{\sqrt{3}}{3}$	1	$\sqrt{3}$	$-\frac{\sqrt{3}}{3}$	-1	$-\sqrt{3}$
$\arctan x$	$0^\circ$	$30^\circ$	$45^\circ$	$60^\circ$	$-30^\circ$	$-45^\circ$	$-60^\circ$
$\operatorname{arccot} x$	$90^\circ$	$60^\circ$	$45^\circ$	$30^\circ$	$120^\circ$	$135^\circ$	$150^\circ$

## 4.19 Relations between Inverse Trigonometric Functions

**474.**  $\arcsin(-x) = -\arcsin x$

**475.**  $\arcsin x = \frac{\pi}{2} - \arccos x$

**476.**  $\arcsin x = \arccos \sqrt{1-x^2}$ ,  $0 \leq x \leq 1$ .

**477.**  $\arcsin x = -\arccos \sqrt{1-x^2}$ ,  $-1 \leq x \leq 0$ .

**478.**  $\arcsin x = \arctan \frac{x}{\sqrt{1-x^2}}$ ,  $x^2 < 1$ .

**479.**  $\arcsin x = \operatorname{arccot} \frac{\sqrt{1-x^2}}{x}$ ,  $0 < x \leq 1$ .

**480.**  $\arcsin x = \operatorname{arccot} \frac{\sqrt{1-x^2}}{x} - \pi$ ,  $-1 \leq x < 0$ .

**481.**  $\arccos(-x) = \pi - \arccos x$

CHAPTER 4. TRIGONOMETRY

$$482. \arccos x = \frac{\pi}{2} - \arcsin x$$

$$483. \arccos x = \arcsin \sqrt{1-x^2}, 0 \leq x \leq 1.$$

$$484. \arccos x = \pi - \arcsin \sqrt{1-x^2}, -1 \leq x \leq 0.$$

$$485. \arccos x = \arctan \frac{\sqrt{1-x^2}}{x}, 0 < x \leq 1.$$

$$486. \arccos x = \pi + \arctan \frac{\sqrt{1-x^2}}{x}, -1 \leq x < 0.$$

$$487. \arccos x = \operatorname{arccot} \frac{x}{\sqrt{1-x^2}}, -1 \leq x \leq 1.$$

$$488. \arctan(-x) = -\arctan x$$

$$489. \arctan x = \frac{\pi}{2} - \operatorname{arccot} x$$

$$490. \arctan x = \arcsin \frac{x}{\sqrt{1+x^2}}$$

$$491. \arctan x = \arccos \frac{1}{\sqrt{1+x^2}}, x \geq 0.$$

$$492. \arctan x = -\arccos \frac{1}{\sqrt{1+x^2}}, x \leq 0.$$

CHAPTER 4. TRIGONOMETRY

$$493. \arctan x = \frac{\pi}{2} - \arctan \frac{1}{x}, x > 0.$$

$$494. \arctan x = -\frac{\pi}{2} - \arctan \frac{1}{x}, x < 0.$$

$$495. \arctan x = \operatorname{arccot} \frac{1}{x}, x > 0.$$

$$496. \arctan x = \operatorname{arccot} \frac{1}{x} - \pi, x < 0.$$

$$497. \operatorname{arccot}(-x) = \pi - \operatorname{arccot} x$$

$$498. \operatorname{arccot} x = \frac{\pi}{2} - \arctan x$$

$$499. \operatorname{arccot} x = \arcsin \frac{1}{\sqrt{1+x^2}}, x > 0.$$

$$500. \operatorname{arccot} x = \pi - \arcsin \frac{1}{\sqrt{1+x^2}}, x < 0.$$

$$501. \operatorname{arccot} x = \arccos \frac{x}{\sqrt{1+x^2}}$$

$$502. \operatorname{arccot} x = \arctan \frac{1}{x}, x > 0.$$

$$503. \operatorname{arccot} x = \pi + \arctan \frac{1}{x}, x < 0.$$

CHAPTER 4. TRIGONOMETRY

## 4.20 Trigonometric Equations

Whole number:  $n$

**504.**  $\sin x = a$ ,  $x = (-1)^n \arcsin a + \pi n$

**505.**  $\cos x = a$ ,  $x = \pm \arccos a + 2\pi n$

**506.**  $\tan x = a$ ,  $x = \arctan a + \pi n$

**507.**  $\cot x = a$ ,  $x = \operatorname{arc cot} a + \pi n$

## 4.21 Relations to Hyperbolic Functions

Imaginary unit:  $i$

**508.**  $\sin(ix) = i \sinh x$

**509.**  $\tan(ix) = i \tanh x$

**510.**  $\cot(ix) = -i \coth x$

**511.**  $\sec(ix) = \operatorname{sech} x$

**512.**  $\csc(ix) = -i \operatorname{csch} x$

## **Chapter 5**

# **Matrices and Determinants**

Matrices:  $A, B, C$

Elements of a matrix:  $a_i, b_i, a_{ij}, b_{ij}, c_{ij}$

Determinant of a matrix:  $\det A$

Minor of an element  $a_{ij}$ :  $M_{ij}$

Cofactor of an element  $a_{ij}$ :  $C_{ij}$

Transpose of a matrix:  $A^T, \tilde{A}$

Adjoint of a matrix:  $\text{adj } A$

Trace of a matrix:  $\text{tr } A$

Inverse of a matrix:  $A^{-1}$

Real number:  $k$

Real variables:  $x_i$

Natural numbers:  $m, n$

### 5.1 Determinants


#### **513.** Second Order Determinant

$$\det A = \begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix} = a_1 b_2 - a_2 b_1$$

## CHAPTER 5. MATRICES AND DETERMINANTS

**514.** Third Order Determinant

$$\det A = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} - \\ - a_{11}a_{23}a_{32} - a_{12}a_{21}a_{33} - a_{13}a_{22}a_{31}$$

**515.** Sarrus Rule (Arrow Rule)**Figure 72.****516.** N-th Order Determinant

$$\det A = \begin{vmatrix} a_{11} & a_{12} & \cdots & a_{1j} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2j} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ a_{i1} & a_{i2} & \cdots & a_{ij} & \cdots & a_{in} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ a_{n1} & a_{n2} & \cdots & a_{nj} & \cdots & a_{nn} \end{vmatrix}$$

**517.** Minor

The minor  $M_{ij}$  associated with the element  $a_{ij}$  of  $n$ -th order matrix  $A$  is the  $(n-1)$ -th order determinant derived from the matrix  $A$  by deletion of its  $i$ -th row and  $j$ -th column.

## CHAPTER 5. MATRICES AND DETERMINANTS

**518.** Cofactor

$$C_{ij} = (-1)^{i+j} M_{ij}$$

**519.** Laplace Expansion of n-th Order Determinant

Laplace expansion by elements of the i-th row

$$\det A = \sum_{j=1}^n a_{ij} C_{ij}, \quad i = 1, 2, \dots, n.$$

Laplace expansion by elements of the j-th column

$$\det A = \sum_{i=1}^n a_{ij} C_{ij}, \quad j = 1, 2, \dots, n.$$

**5.2 Properties of Determinants****520.** The value of a determinant remains unchanged if rows are changed to columns and columns to rows.

$$\begin{vmatrix} a_1 & a_2 \\ b_1 & b_2 \end{vmatrix} = \begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix}$$

**521.** If two rows (or two columns) are interchanged, the sign of the determinant is changed.

$$\begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix} = - \begin{vmatrix} a_2 & b_2 \\ a_1 & b_1 \end{vmatrix}$$

**522.** If two rows (or two columns) are identical, the value of the determinant is zero.

$$\begin{vmatrix} a_1 & a_1 \\ a_2 & a_2 \end{vmatrix} = 0$$

## CHAPTER 5. MATRICES AND DETERMINANTS

- 523.** If the elements of any row (or column) are multiplied by a common factor, the determinant is multiplied by that factor.

$$\begin{vmatrix} ka_1 & kb_1 \\ a_2 & b_2 \end{vmatrix} = k \begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix}$$

- 524.** If the elements of any row (or column) are increased (or decreased) by equal multiples of the corresponding elements of any other row (or column), the value of the determinant is unchanged.

$$\begin{vmatrix} a_1 + kb_1 & b_1 \\ a_2 + kb_2 & b_2 \end{vmatrix} = \begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix}$$

### 5.3 Matrices

- 525.** Definition

An  $m \times n$  matrix  $A$  is a rectangular array of elements (numbers or functions) with  $m$  rows and  $n$  columns.

$$A = [a_{ij}] = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- 526.** **Square matrix** is a matrix of order  $n \times n$ .
- 527.** A square matrix  $[a_{ij}]$  is **symmetric** if  $a_{ij} = a_{ji}$ , i.e. it is symmetric about the leading diagonal.
- 528.** A square matrix  $[a_{ij}]$  is **skew-symmetric** if  $a_{ij} = -a_{ji}$ .



## CHAPTER 5. MATRICES AND DETERMINANTS

- 529.** **Diagonal matrix** is a square matrix with all elements zero except those on the leading diagonal.
- 530.** **Unit matrix** is a diagonal matrix in which the elements on the leading diagonal are all unity. The unit matrix is denoted by I.
- 531.** A **null matrix** is one whose elements are all zero.

## 5.4 Operations with Matrices

- 532.** Two matrices A and B are equal if, and only if, they are both of the same shape  $m \times n$  and corresponding elements are equal.
- 533.** Two matrices A and B can be added (or subtracted) of, and only if, they have the same shape  $m \times n$ . If

$$A = [a_{ij}] = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix},$$

$$B = [b_{ij}] = \begin{bmatrix} b_{11} & b_{12} & \cdots & b_{1n} \\ b_{21} & b_{22} & \cdots & b_{2n} \\ \vdots & \vdots & & \vdots \\ b_{m1} & b_{m2} & \cdots & b_{mn} \end{bmatrix},$$

## CHAPTER 5. MATRICES AND DETERMINANTS

then

$$A + B = \begin{bmatrix} a_{11} + b_{11} & a_{12} + b_{12} & \dots & a_{1n} + b_{1n} \\ a_{21} + b_{21} & a_{22} + b_{22} & \dots & a_{2n} + b_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} + b_{m1} & a_{m2} + b_{m2} & \dots & a_{mn} + b_{mn} \end{bmatrix}.$$

**534.** If  $k$  is a scalar, and  $A = [a_{ij}]$  is a matrix, then

$$kA = [ka_{ij}] = \begin{bmatrix} ka_{11} & ka_{12} & \dots & ka_{1n} \\ ka_{21} & ka_{22} & \dots & ka_{2n} \\ \vdots & \vdots & & \vdots \\ ka_{m1} & ka_{m2} & \dots & ka_{mn} \end{bmatrix}.$$

**535.** Multiplication of Two Matrices

Two matrices can be multiplied together only when the number of columns in the first is equal to the number of rows in the second.

If

$$A = [a_{ij}] = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix},$$

$$B = [b_{ij}] = \begin{bmatrix} b_{11} & b_{12} & \dots & b_{1k} \\ b_{21} & b_{22} & \dots & b_{2k} \\ \vdots & \vdots & & \vdots \\ b_{n1} & b_{n2} & \dots & b_{nk} \end{bmatrix},$$

## CHAPTER 5. MATRICES AND DETERMINANTS

then

$$AB = C = \begin{bmatrix} c_{11} & c_{12} & \cdots & c_{1k} \\ c_{21} & c_{22} & \cdots & c_{2k} \\ \vdots & \vdots & & \vdots \\ b_{m1} & c_{m2} & \cdots & c_{mk} \end{bmatrix},$$

where

$$c_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{in}b_{nj} = \sum_{\lambda=1}^n a_{i\lambda}b_{\lambda j}$$

( $i = 1, 2, \dots, m; j = 1, 2, \dots, k$ ).

Thus if

$$A = [a_{ij}] = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{bmatrix}, \quad B = [b_i] = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix},$$

then

$$AB = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{bmatrix} \cdot \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix} = \begin{bmatrix} a_{11}b_1 & a_{12}b_2 & a_{13}b_3 \\ a_{21}b_1 & a_{22}b_2 & a_{23}b_3 \end{bmatrix}.$$

**536.** Transpose of a Matrix

If the rows and columns of a matrix are interchanged, then the new matrix is called the **transpose** of the original matrix.

If  $A$  is the original matrix, its transpose is denoted  $A^T$  or  $\tilde{A}$ .

**537.** The matrix  $A$  is **orthogonal** if  $AA^T = I$ .

**538.** If the matrix product  $AB$  is defined, then

$$(AB)^T = B^T A^T.$$

## CHAPTER 5. MATRICES AND DETERMINANTS

### 539. Adjoint of Matrix

If  $A$  is a square  $n \times n$  matrix, its **adjoint**, denoted by  $\text{adj } A$ , is the transpose of the matrix of cofactors  $C_{ij}$  of  $A$ :

$$\text{adj } A = [C_{ij}]^T.$$

### 540. Trace of a Matrix

If  $A$  is a square  $n \times n$  matrix, its **trace**, denoted by  $\text{tr } A$ , is defined to be the sum of the terms on the leading diagonal:

$$\text{tr } A = a_{11} + a_{22} + \dots + a_{nn}.$$

### 541. Inverse of a Matrix

If  $A$  is a square  $n \times n$  matrix with a nonsingular determinant  $\det A$ , then its **inverse**  $A^{-1}$  is given by

$$A^{-1} = \frac{\text{adj } A}{\det A}.$$

### 542. If the matrix product $AB$ is defined, then

$$(AB)^{-1} = B^{-1}A^{-1}.$$

### 543. If $A$ is a square $n \times n$ matrix, the **eigenvectors** $X$ satisfy the equation

$$AX = \lambda X,$$

while the **eigenvalues**  $\lambda$  satisfy the characteristic equation

$$|A - \lambda I| = 0.$$

## 5.5 Systems of Linear Equations

Variables:  $x, y, z, x_1, x_2, \dots$

Real numbers:  $a_1, a_2, a_3, b_1, a_{11}, a_{12}, \dots$

## CHAPTER 5. MATRICES AND DETERMINANTS

Determinants:  $D, D_x, D_y, D_z$

Matrices:  $A, B, X$

$$544. \begin{cases} a_1x + b_1y = d_1 \\ a_2x + b_2y = d_2 \end{cases},$$

$$x = \frac{D_x}{D}, y = \frac{D_y}{D} \text{ (Cramer's rule),}$$

where

$$D = \begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix} = a_1b_2 - a_2b_1,$$

$$D_x = \begin{vmatrix} d_1 & b_1 \\ d_2 & b_2 \end{vmatrix} = d_1b_2 - d_2b_1,$$

$$D_y = \begin{vmatrix} a_1 & d_1 \\ a_2 & d_2 \end{vmatrix} = a_1d_2 - a_2d_1.$$

545. If  $D \neq 0$ , then the system has a single solution:

$$x = \frac{D_x}{D}, y = \frac{D_y}{D}.$$

If  $D = 0$  and  $D_x \neq 0$  (or  $D_y \neq 0$ ), then the system has no solution.

If  $D = D_x = D_y = 0$ , then the system has infinitely many solutions.

$$546. \begin{cases} a_1x + b_1y + c_1z = d_1 \\ a_2x + b_2y + c_2z = d_2, \\ a_3x + b_3y + c_3z = d_3 \end{cases}$$

$$x = \frac{D_x}{D}, y = \frac{D_y}{D}, z = \frac{D_z}{D} \text{ (Cramer's rule),}$$

CHAPTER 5. MATRICES AND DETERMINANTS

where

$$D = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}, \quad D_x = \begin{vmatrix} d_1 & b_1 & c_1 \\ d_2 & b_2 & c_2 \\ d_3 & b_3 & c_3 \end{vmatrix},$$

$$D_y = \begin{vmatrix} a_1 & d_1 & c_1 \\ a_2 & d_2 & c_2 \\ a_3 & d_3 & c_3 \end{vmatrix}, \quad D_z = \begin{vmatrix} a_1 & b_1 & d_1 \\ a_2 & b_2 & d_2 \\ a_3 & b_3 & d_3 \end{vmatrix}.$$

547. If  $D \neq 0$ , then the system has a single solution:

$$x = \frac{D_x}{D}, \quad y = \frac{D_y}{D}, \quad z = \frac{D_z}{D}.$$

If  $D = 0$  and  $D_x \neq 0$  (or  $D_y \neq 0$  or  $D_z \neq 0$ ), then the system has no solution.

If  $D = D_x = D_y = D_z = 0$ , then the system has infinitely many solutions.

548. Matrix Form of a System of n Linear Equations in n Unknowns

The set of linear equations

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2 \\ \dots \dots \dots \\ a_{n1}x_1 + a_{n2}x_2 + \dots + a_{nn}x_n = b_n \end{cases}$$

can be written in matrix form

$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{pmatrix} \cdot \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix},$$

i.e.  
 $A \cdot X = B$ ,

CHAPTER 5. MATRICES AND DETERMINANTS

where

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{pmatrix}, \mathbf{X} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}, \mathbf{B} = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix}.$$

**549.** Solution of a Set of Linear Equations  $n \times n$

$$\mathbf{X} = \mathbf{A}^{-1} \cdot \mathbf{B},$$

where  $\mathbf{A}^{-1}$  is the inverse of  $\mathbf{A}$ .

## Chapter 6

# Vectors

Vectors:  $\vec{u}$ ,  $\vec{v}$ ,  $\vec{w}$ ,  $\vec{r}$ ,  $\vec{AB}$ , ...

Vector length:  $|\vec{u}|$ ,  $|\vec{v}|$ , ...

Unit vectors:  $\vec{i}$ ,  $\vec{j}$ ,  $\vec{k}$

Null vector:  $\vec{0}$

Coordinates of vector  $\vec{u}$ :  $X_1, Y_1, Z_1$

Coordinates of vector  $\vec{v}$ :  $X_2, Y_2, Z_2$

Scalars:  $\lambda, \mu$

Direction cosines:  $\cos \alpha$ ,  $\cos \beta$ ,  $\cos \gamma$

Angle between two vectors:  $\theta$

### 6.1 Vector Coordinates

**550.** Unit Vectors

$$\vec{i} = (1, 0, 0),$$

$$\vec{j} = (0, 1, 0),$$

$$\vec{k} = (0, 0, 1),$$

$$|\vec{i}| = |\vec{j}| = |\vec{k}| = 1.$$

**551.**  $\vec{r} = \vec{AB} = (x_1 - x_0)\vec{i} + (y_1 - y_0)\vec{j} + (z_1 - z_0)\vec{k}$



CHAPTER 6. VECTORS




Figure 73.

552.  $|\vec{r}| = |\vec{AB}| = \sqrt{(x_1 - x_0)^2 + (y_1 - y_0)^2 + (z_1 - z_0)^2}$

553. If  $\vec{AB} = \vec{r}$ , then  $\vec{BA} = -\vec{r}$ .




Figure 74.

554.  $X = |\vec{r}| \cos \alpha,$   
 $Y = |\vec{r}| \cos \beta,$   
 $Z = |\vec{r}| \cos \gamma.$

CHAPTER 6. VECTORS




Figure 75.

555. If  $\vec{r}(X, Y, Z) = \vec{r}_1(X_1, Y_1, Z_1)$ , then  $X = X_1, Y = Y_1, Z = Z_1$ .

## 6.2 Vector Addition

556.  $\vec{w} = \vec{u} + \vec{v}$




Figure 76.

CHAPTER 6. VECTORS




Figure 77.

557.  $\vec{w} = \vec{u}_1 + \vec{u}_2 + \vec{u}_3 + \dots + \vec{u}_n$




Figure 78.

558. Commutative Law  
 $\vec{u} + \vec{v} = \vec{v} + \vec{u}$

559. Associative Law  
 $(\vec{u} + \vec{v}) + \vec{w} = \vec{u} + (\vec{v} + \vec{w})$

560.  $\vec{u} + \vec{v} = (X_1 + X_2, Y_1 + Y_2, Z_1 + Z_2)$

CHAPTER 6. VECTORS

### 6.3 Vector Subtraction

**561.**  $\vec{w} = \vec{u} - \vec{v}$  if  $\vec{v} + \vec{w} = \vec{u}$ .




Figure 79.




Figure 80.

**562.**  $\vec{u} - \vec{v} = \vec{u} + (-\vec{v})$

**563.**  $\vec{u} - \vec{u} = \vec{0} = (0, 0, 0)$

**564.**  $|\vec{0}| = 0$

**565.**  $\vec{u} - \vec{v} = (X_1 - X_2, Y_1 - Y_2, Z_1 - Z_2),$

### 6.4 Scaling Vectors

**566.**  $\vec{w} = \lambda \vec{u}$

## CHAPTER 6. VECTORS




Figure 81.

567.  $|\vec{w}| = |\lambda| \cdot |\vec{u}|$
568.  $\lambda \vec{u} = (\lambda X, \lambda Y, \lambda Z)$
569.  $\lambda \vec{u} = \vec{u} \lambda$
570.  $(\lambda + \mu) \vec{u} = \lambda \vec{u} + \mu \vec{u}$
571.  $\lambda(\mu \vec{u}) = \mu(\lambda \vec{u}) = (\lambda \mu) \vec{u}$
572.  $\lambda(\vec{u} + \vec{v}) = \lambda \vec{u} + \lambda \vec{v}$

## 6.5 Scalar Product

573. Scalar Product of Vectors  $\vec{u}$  and  $\vec{v}$   
 $\vec{u} \cdot \vec{v} = |\vec{u}| \cdot |\vec{v}| \cdot \cos \theta$ ,  
 where  $\theta$  is the angle between vectors  $\vec{u}$  and  $\vec{v}$ .

## CHAPTER 6. VECTORS




Figure 82.

**574. Scalar Product in Coordinate Form**

If  $\vec{u} = (X_1, Y_1, Z_1)$ ,  $\vec{v} = (X_2, Y_2, Z_2)$ , then

$$\vec{u} \cdot \vec{v} = X_1X_2 + Y_1Y_2 + Z_1Z_2.$$

**575. Angle Between Two Vectors**

If  $\vec{u} = (X_1, Y_1, Z_1)$ ,  $\vec{v} = (X_2, Y_2, Z_2)$ , then

$$\cos \theta = \frac{X_1X_2 + Y_1Y_2 + Z_1Z_2}{\sqrt{X_1^2 + Y_1^2 + Z_1^2} \sqrt{X_2^2 + Y_2^2 + Z_2^2}}.$$

**576. Commutative Property**

$$\vec{u} \cdot \vec{v} = \vec{v} \cdot \vec{u}$$

**577. Associative Property**

$$(\lambda \vec{u}) \cdot (\mu \vec{v}) = \lambda \mu \vec{u} \cdot \vec{v}$$

**578. Distributive Property**

$$\vec{u} \cdot (\vec{v} + \vec{w}) = \vec{u} \cdot \vec{v} + \vec{u} \cdot \vec{w}$$

**579.**  $\vec{u} \cdot \vec{v} = 0$  if  $\vec{u}, \vec{v}$  are orthogonal ( $\theta = \frac{\pi}{2}$ ).

**580.**  $\vec{u} \cdot \vec{v} > 0$  if  $0 < \theta < \frac{\pi}{2}$ .

## CHAPTER 6. VECTORS

- 581.**  $\vec{u} \cdot \vec{v} < 0$  if  $\frac{\pi}{2} < \theta < \pi$ .
- 582.**  $\vec{u} \cdot \vec{v} \leq |\vec{u}| \cdot |\vec{v}|$
- 583.**  $\vec{u} \cdot \vec{v} = |\vec{u}| \cdot |\vec{v}|$  if  $\vec{u}, \vec{v}$  are parallel ( $\theta = 0$ ).
- 584.** If  $\vec{u} = (X_1, Y_1, Z_1)$ , then  
 $\vec{u} \cdot \vec{u} = \vec{u}^2 = |\vec{u}|^2 = X_1^2 + Y_1^2 + Z_1^2$ .
- 585.**  $\vec{i} \cdot \vec{i} = \vec{j} \cdot \vec{j} = \vec{k} \cdot \vec{k} = 1$
- 586.**  $\vec{i} \cdot \vec{j} = \vec{j} \cdot \vec{k} = \vec{k} \cdot \vec{i} = 0$

### 6.6 Vector Product

- 587.** Vector Product of Vectors  $\vec{u}$  and  $\vec{v}$   
 $\vec{u} \times \vec{v} = \vec{w}$ , where
- $|\vec{w}| = |\vec{u}| \cdot |\vec{v}| \cdot \sin \theta$ , where  $0 \leq \theta \leq \frac{\pi}{2}$ ;
  - $\vec{w} \perp \vec{u}$  and  $\vec{w} \perp \vec{v}$ ;
  - Vectors  $\vec{u}, \vec{v}, \vec{w}$  form a right-handed screw.

## CHAPTER 6. VECTORS




Figure 83.

$$588. \quad \vec{w} = \vec{u} \times \vec{v} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ X_1 & Y_1 & Z_1 \\ X_2 & Y_2 & Z_2 \end{vmatrix}$$

$$589. \quad \vec{w} = \vec{u} \times \vec{v} = \left( \begin{vmatrix} Y_1 & Z_1 \\ Y_2 & Z_2 \end{vmatrix}, -\begin{vmatrix} X_1 & Z_1 \\ X_2 & Z_2 \end{vmatrix}, \begin{vmatrix} X_1 & Y_1 \\ X_2 & Y_2 \end{vmatrix} \right)$$

$$590. \quad S = |\vec{u} \times \vec{v}| = |\vec{u}| \cdot |\vec{v}| \cdot \sin \theta \quad (\text{Fig.83})$$

591. Angle Between Two Vectors (Fig.83)

$$\sin \theta = \frac{|\vec{u} \times \vec{v}|}{|\vec{u}| \cdot |\vec{v}|}$$

592. Noncommutative Property

$$\vec{u} \times \vec{v} = -(\vec{v} \times \vec{u})$$

593. Associative Property

$$(\lambda \vec{u}) \times (\mu \vec{v}) = \lambda \mu \vec{u} \times \vec{v}$$



## CHAPTER 6. VECTORS

**594. Distributive Property**

$$\vec{u} \times (\vec{v} + \vec{w}) = \vec{u} \times \vec{v} + \vec{u} \times \vec{w}$$

**595.**  $\vec{u} \times \vec{v} = \vec{0}$  if  $\vec{u}$  and  $\vec{v}$  are parallel ( $\theta = 0$ ).**596.**  $\vec{i} \times \vec{i} = \vec{j} \times \vec{j} = \vec{k} \times \vec{k} = \vec{0}$ **597.**  $\vec{i} \times \vec{j} = \vec{k}$ ,  $\vec{j} \times \vec{k} = \vec{i}$ ,  $\vec{k} \times \vec{i} = \vec{j}$ 

## 6.7 Triple Product

**598. Scalar Triple Product**

$$[\vec{u}\vec{v}\vec{w}] = \vec{u} \cdot (\vec{v} \times \vec{w}) = \vec{v} \cdot (\vec{w} \times \vec{u}) = \vec{w} \cdot (\vec{u} \times \vec{v})$$

**599.**  $[\vec{u}\vec{v}\vec{w}] = [\vec{w}\vec{u}\vec{v}] = [\vec{v}\vec{w}\vec{u}] = -[\vec{v}\vec{u}\vec{w}] = -[\vec{w}\vec{v}\vec{u}] = -[\vec{u}\vec{w}\vec{v}]$ **600.**  $k\vec{u} \cdot (\vec{v} \times \vec{w}) = k[\vec{u}\vec{v}\vec{w}]$ **601. Scalar Triple Product in Coordinate Form**

$$\vec{u} \cdot (\vec{v} \times \vec{w}) = \begin{vmatrix} X_1 & Y_1 & Z_1 \\ X_2 & Y_2 & Z_2 \\ X_3 & Y_3 & Z_3 \end{vmatrix},$$

where

$$\vec{u} = (X_1, Y_1, Z_1), \vec{v} = (X_2, Y_2, Z_2), \vec{w} = (X_3, Y_3, Z_3).$$

**602. Volume of Parallelepiped**

$$V = |[\vec{u} \cdot (\vec{v} \times \vec{w})]|$$

CHAPTER 6. VECTORS




Figure 84.

603. Volume of Pyramid

$$V = \frac{1}{6} |\vec{u} \cdot (\vec{v} \times \vec{w})|$$




Figure 85.

604. If  $\vec{u} \cdot (\vec{v} \times \vec{w}) = 0$ , then the vectors  $\vec{u}$ ,  $\vec{v}$ , and  $\vec{w}$  are **linearly dependent**, so  $\vec{w} = \lambda \vec{u} + \mu \vec{v}$  for some scalars  $\lambda$  and  $\mu$ .

605. If  $\vec{u} \cdot (\vec{v} \times \vec{w}) \neq 0$ , then the vectors  $\vec{u}$ ,  $\vec{v}$ , and  $\vec{w}$  are **linearly independent**.

CHAPTER 6. VECTORS

**606.** Vector Triple Product

$$\vec{u} \times (\vec{v} \times \vec{w}) = (\vec{u} \cdot \vec{w})\vec{v} - (\vec{u} \cdot \vec{v})\vec{w}$$

## Chapter 7

# Analytic Geometry

### 7.1 One-Dimensional Coordinate System

Point coordinates:  $x_0, x_1, x_2, y_0, y_1, y_2$

Real number:  $\lambda$

Distance between two points:  $d$

**607.** Distance Between Two Points

$$d = AB = |x_2 - x_1| = |x_1 - x_2|$$




Figure 86.

**608.** Dividing a Line Segment in the Ratio  $\lambda$

$$x_0 = \frac{x_1 + \lambda x_2}{1 + \lambda}, \quad \lambda = \frac{AC}{CB}, \quad \lambda \neq -1.$$




Figure 87.

CHAPTER 7. ANALYTIC GEOMETRY

**609.** Midpoint of a Line Segment

$$x_0 = \frac{x_1 + x_2}{2}, \lambda = 1.$$

## 7.2 Two-Dimensional Coordinate System

Point coordinates:  $x_0, x_1, x_2, y_0, y_1, y_2$

Polar coordinates:  $r, \varphi$

Real number:  $\lambda$

Positive real numbers:  $a, b, c,$

Distance between two points:  $d$

Area:  $S$

**610.** Distance Between Two Points

$$d = AB = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$





Figure 88.

CHAPTER 7. ANALYTIC GEOMETRY

**611.** Dividing a Line Segment in the Ratio  $\lambda$

$$x_0 = \frac{x_1 + \lambda x_2}{1 + \lambda}, \quad y_0 = \frac{y_1 + \lambda y_2}{1 + \lambda},$$

$$\lambda = \frac{AC}{CB}, \quad \lambda \neq -1.$$



**Figure 89.**

CHAPTER 7. ANALYTIC GEOMETRY




Figure 90.

**612.** Midpoint of a Line Segment

$$x_0 = \frac{x_1 + x_2}{2}, y_0 = \frac{y_1 + y_2}{2}, \lambda = 1.$$

**613.** Centroid (Intersection of Medians) of a Triangle

$$x_0 = \frac{x_1 + x_2 + x_3}{3}, y_0 = \frac{y_1 + y_2 + y_3}{3},$$

where  $A(x_1, y_1)$ ,  $B(x_2, y_2)$ , and  $C(x_3, y_3)$  are vertices of the triangle ABC.

CHAPTER 7. ANALYTIC GEOMETRY




Figure 91.

**614.** Incenter (Intersection of Angle Bisectors) of a Triangle

$$x_0 = \frac{ax_1 + bx_2 + cx_3}{a + b + c}, \quad y_0 = \frac{ay_1 + by_2 + cy_3}{a + b + c},$$

where  $a = BC$ ,  $b = CA$ ,  $c = AB$ .




Figure 92.



CHAPTER 7. ANALYTIC GEOMETRY

**615.** Circumcenter (Intersection of the Side Perpendicular Bisectors) of a Triangle

$$x_0 = \frac{\begin{vmatrix} x_1^2 + y_1^2 & y_1 & 1 \\ x_2^2 + y_2^2 & y_2 & 1 \\ x_3^2 + y_3^2 & y_3 & 1 \end{vmatrix}}{\begin{vmatrix} x_1 & y_1 & 1 \\ 2x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}}, \quad y_0 = \frac{\begin{vmatrix} x_1 & x_1^2 + y_1^2 & 1 \\ x_2 & x_2^2 + y_2^2 & 1 \\ x_3 & x_3^2 + y_3^2 & 1 \end{vmatrix}}{\begin{vmatrix} x_1 & y_1 & 1 \\ 2x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}}$$





Figure 93.

## CHAPTER 7. ANALYTIC GEOMETRY

**616.** Orthocenter (Intersection of Altitudes) of a Triangle

$$x_0 = \frac{\begin{vmatrix} y_1 & x_2x_3 + y_1^2 & 1 \\ y_2 & x_3x_1 + y_2^2 & 1 \\ y_3 & x_1x_2 + y_3^2 & 1 \end{vmatrix}}{\begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}}, \quad y_0 = \frac{\begin{vmatrix} x_1^2 + y_2y_3 & x_1 & 1 \\ x_2^2 + y_3y_1 & x_2 & 1 \\ x_3^2 + y_1y_2 & x_3 & 1 \end{vmatrix}}{\begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}}$$

**Figure 94.****617.** Area of a Triangle

$$S = (\pm) \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} = (\pm) \frac{1}{2} \begin{vmatrix} x_2 - x_1 & y_2 - y_1 \\ x_3 - x_1 & y_3 - y_1 \end{vmatrix}$$

CHAPTER 7. ANALYTIC GEOMETRY

**618.** Area of a Quadrilateral

$$S = (\pm) \frac{1}{2} [(x_1 - x_2)(y_1 + y_2) + (x_2 - x_3)(y_2 + y_3) + (x_3 - x_4)(y_3 + y_4) + (x_4 - x_1)(y_4 + y_1)]$$




Figure 95.

Note: In formulas 617, 618 we choose the sign (+) or (-) so that to get a positive answer for area.

**619.** Distance Between Two Points in Polar Coordinates

$$d = AB = \sqrt{r_1^2 + r_2^2 - 2r_1r_2 \cos(\varphi_2 - \varphi_1)}$$

CHAPTER 7. ANALYTIC GEOMETRY




Figure 96.

- 620.** Converting Rectangular Coordinates to Polar Coordinates  
 $x = r \cos \varphi$ ,  $y = r \sin \varphi$ .




Figure 97.

- 621.** Converting Polar Coordinates to Rectangular Coordinates  
 $r = \sqrt{x^2 + y^2}$ ,  $\tan \varphi = \frac{y}{x}$ .

CHAPTER 7. ANALYTIC GEOMETRY

### 7.3 Straight Line in Plane

Point coordinates:  $X, Y, x, x_0, x_1, y_0, y_1, a_1, a_2, \dots$

Real numbers:  $k, a, b, p, t, A, B, C, A_1, A_2, \dots$

Angles:  $\alpha, \beta$

Angle between two lines:  $\varphi$

Normal vector:  $\vec{n}$

Position vectors:  $\vec{r}, \vec{a}, \vec{b}$

**622.** General Equation of a Straight Line

$$Ax + By + C = 0$$

**623.** Normal Vector to a Straight Line

The vector  $\vec{n}(A, B)$  is normal to the line  $Ax + By + C = 0$ .




Figure 98.

**624.** Explicit Equation of a Straight Line (Slope-Intercept Form)

$$y = kx + b.$$

CHAPTER 7. ANALYTIC GEOMETRY

The gradient of the line is  $k = \tan \alpha$ .




Figure 99.

**625.** Gradient of a Line

$$k = \tan \alpha = \frac{y_2 - y_1}{x_2 - x_1}$$




Figure 100.

CHAPTER 7. ANALYTIC GEOMETRY

- 626.** Equation of a Line Given a Point and the Gradient  
 $y = y_0 + k(x - x_0)$ ,  
where  $k$  is the gradient,  $P(x_0, y_0)$  is a point on the line.




Figure 101.

- 627.** Equation of a Line That Passes Through Two Points

$$\frac{y - y_1}{y_2 - y_1} = \frac{x - x_1}{x_2 - x_1}$$

or

$$\begin{vmatrix} x & y & 1 \\ x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \end{vmatrix} = 0.$$

CHAPTER 7. ANALYTIC GEOMETRY




Figure 102.

**628.** Intercept Form

$$\frac{x}{a} + \frac{y}{b} = 1$$




Figure 103.



CHAPTER 7. ANALYTIC GEOMETRY

- 629.** Normal Form  
 $x \cos \beta + y \sin \beta - p = 0$




Figure 104.

- 630.** Point Direction Form

$$\frac{x - x_1}{X} = \frac{y - y_1}{Y},$$

where  $(X, Y)$  is the direction of the line and  $P_1(x_1, y_1)$  lies on the line.

CHAPTER 7. ANALYTIC GEOMETRY




Figure 105.

**631.** Vertical Line

$$x = a$$

**632.** Horizontal Line

$$y = b$$

**633.** Vector Equation of a Straight Line

$$\vec{r} = \vec{a} + t\vec{b},$$

where

O is the origin of the coordinates,

X is any variable point on the line,

$\vec{a}$  is the position vector of a known point A on the line ,

$\vec{b}$  is a known vector of direction, parallel to the line,

t is a parameter,

$\vec{r} = \vec{OX}$  is the position vector of any point X on the line.

CHAPTER 7. ANALYTIC GEOMETRY




Figure 106.

**634.** Straight Line in Parametric Form

$$\begin{cases} x = a_1 + tb_1 \\ y = a_2 + tb_2 \end{cases},$$

where

$(x, y)$  are the coordinates of any unknown point on the line,

$(a_1, a_2)$  are the coordinates of a known point on the line,

$(b_1, b_2)$  are the coordinates of a vector parallel to the line,

$t$  is a parameter.

CHAPTER 7. ANALYTIC GEOMETRY




Figure 107.

- 635.** Distance From a Point To a Line  
The distance from the point  $P(a, b)$  to the line

$Ax + By + C = 0$  is

$$d = \frac{|Aa + Bb + C|}{\sqrt{A^2 + B^2}}.$$




Figure 108.

## CHAPTER 7. ANALYTIC GEOMETRY

**636. Parallel Lines**

Two lines  $y = k_1x + b_1$  and  $y = k_2x + b_2$  are parallel if

$$k_1 = k_2.$$

Two lines  $A_1x + B_1y + C_1 = 0$  and  $A_2x + B_2y + C_2 = 0$  are parallel if

$$\frac{A_1}{A_2} = \frac{B_1}{B_2}.$$




Figure 109.

**637. Perpendicular Lines**

Two lines  $y = k_1x + b_1$  and  $y = k_2x + b_2$  are perpendicular if

$$k_2 = -\frac{1}{k_1} \text{ or, equivalently, } k_1k_2 = -1.$$

Two lines  $A_1x + B_1y + C_1 = 0$  and  $A_2x + B_2y + C_2 = 0$  are perpendicular if

$$A_1A_2 + B_1B_2 = 0.$$

CHAPTER 7. ANALYTIC GEOMETRY




Figure 110.

**638.** Angle Between Two Lines

$$\tan \varphi = \frac{k_2 - k_1}{1 + k_1 k_2},$$

$$\cos \varphi = \frac{A_1 A_2 + B_1 B_2}{\sqrt{A_1^2 + B_1^2} \cdot \sqrt{A_2^2 + B_2^2}}.$$

## CHAPTER 7. ANALYTIC GEOMETRY




Figure 111.

**639.** Intersection of Two Lines

If two lines  $A_1x + B_1y + C_1 = 0$  and  $A_2x + B_2y + C_2 = 0$  intersect, the intersection point has coordinates

$$x_0 = \frac{-C_1B_2 + C_2B_1}{A_1B_2 - A_2B_1}, \quad y_0 = \frac{-A_1C_2 + A_2C_1}{A_1B_2 - A_2B_1}.$$

## 7.4 Circle

Radius:  $R$

Center of circle:  $(a, b)$

Point coordinates:  $x, y, x_1, y_1, \dots$

Real numbers:  $A, B, C, D, E, F, t$

CHAPTER 7. ANALYTIC GEOMETRY

- 640.** Equation of a Circle Centered at the Origin (Standard Form)

$$x^2 + y^2 = R^2$$




Figure 112.

- 641.** Equation of a Circle Centered at Any Point (a, b)

$$(x - a)^2 + (y - b)^2 = R^2$$




Figure 113.



## CHAPTER 7. ANALYTIC GEOMETRY

**642. Three Point Form**

$$\begin{vmatrix} x^2 + y^2 & x & y & 1 \\ x_1^2 + y_1^2 & x_1 & y_1 & 1 \\ x_2^2 + y_2^2 & x_2 & y_2 & 1 \\ x_3^2 + y_3^2 & x_3 & y_3 & 1 \end{vmatrix} = 0$$




Figure 114.

**643. Parametric Form**

$$\begin{cases} x = R \cos t \\ y = R \sin t \end{cases}, \quad 0 \leq t \leq 2\pi.$$

**644. General Form**

$$Ax^2 + Ay^2 + Dx + Ey + F = 0 \quad (A \text{ nonzero}, D^2 + E^2 > 4AF).$$

The center of the circle has coordinates  $(a, b)$ , where

$$a = -\frac{D}{2A}, \quad b = -\frac{E}{2A}.$$

The radius of the circle is

CHAPTER 7. ANALYTIC GEOMETRY

$$R = \sqrt{\frac{D^2 + E^2 - 4AF}{2|A|}}.$$

## 7.5 Ellipse

Semimajor axis:  $a$

Semiminor axis:  $b$

Foci:  $F_1(-c, 0)$ ,  $F_2(c, 0)$

Distance between the foci:  $2c$

Eccentricity:  $e$

Real numbers:  $A, B, C, D, E, F, t$

Perimeter:  $L$

Area:  $S$

**645.** Equation of an Ellipse (Standard Form)

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$




Figure 115.

CHAPTER 7. ANALYTIC GEOMETRY

- 646.**  $r_1 + r_2 = 2a$ ,  
where  $r_1$ ,  $r_2$  are distances from any point  $P(x, y)$  on the ellipse to the two foci.




Figure 116.

- 647.**  $a^2 = b^2 + c^2$
- 648.** Eccentricity  
$$e = \frac{c}{a} < 1$$
- 649.** Equations of Directrices  
$$x = \pm \frac{a}{e} = \pm \frac{a^2}{c}$$
- 650.** Parametric Form  
$$\begin{cases} x = a \cos t \\ y = b \sin t \end{cases}, 0 \leq t \leq 2\pi.$$

## CHAPTER 7. ANALYTIC GEOMETRY

**651.** General Form

$$Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0,$$

where  $B^2 - 4AC < 0$ .

**652.** General Form with Axes Parallel to the Coordinate Axes

$$Ax^2 + Cy^2 + Dx + Ey + F = 0,$$

where  $AC > 0$ .

**653.** Circumference

$$L = 4aE(e),$$

where the function  $E$  is the complete elliptic integral of the second kind.

**654.** Approximate Formulas of the Circumference

$$L = \pi (1.5(a + b) - \sqrt{ab}),$$

$$L = \pi \sqrt{2(a^2 + b^2)}.$$

**655.**  $S = \pi ab$

## 7.6 Hyperbola

Transverse axis:  $a$

Conjugate axis:  $b$

Foci:  $F_1(-c, 0)$ ,  $F_2(c, 0)$

Distance between the foci:  $2c$

Eccentricity:  $e$


Asymptotes:  $s, t$

Real numbers:  $A, B, C, D, E, F, t, k$

CHAPTER 7. ANALYTIC GEOMETRY

**656.** Equation of a Hyperbola (Standard Form)

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$$



**Figure 117.**

- 657.**  $|r_1 - r_2| = 2a$ ,  
where  $r_1$ ,  $r_2$  are distances from any point  $P(x,y)$  on  
the hyperbola to the two foci.

CHAPTER 7. ANALYTIC GEOMETRY




Figure 118.

**658.** Equations of Asymptotes

$$y = \pm \frac{b}{a}x$$

**659.**  $c^2 = a^2 + b^2$

**660.** Eccentricity

$$e = \frac{c}{a} > 1$$

**661.** Equations of Directrices

$$x = \pm \frac{a}{e} = \pm \frac{a^2}{c}$$

CHAPTER 7. ANALYTIC GEOMETRY

**662.** Parametric Equations of the Right Branch of a Hyperbola

$$\begin{cases} x = a \cosh t \\ y = b \sinh t \end{cases}, 0 \leq t \leq 2\pi.$$

**663.** General Form

$$Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0,$$

where  $B^2 - 4AC > 0$ .

**664.** General Form with Axes Parallel to the Coordinate Axes

$$Ax^2 + Cy^2 + Dx + Ey + F = 0,$$

where  $AC < 0$ .

**665.** Asymptotic Form

$$xy = \frac{e^2}{4},$$

or

$$y = \frac{k}{x}, \text{ where } k = \frac{e^2}{4}.$$

In this case, the asymptotes have equations  $x=0$  and  $y=0$ .

CHAPTER 7. ANALYTIC GEOMETRY




Figure 119.

## 7.7 Parabola

Focal parameter:  $p$

Focus:  $F$

Vertex:  $M(x_0, y_0)$

Real numbers:  $A, B, C, D, E, F, p, a, b, c$

**666.** Equation of a Parabola (Standard Form)

$$y^2 = 2px$$



CHAPTER 7. ANALYTIC GEOMETRY




Figure 120.

Equation of the directrix

$$x = -\frac{p}{2},$$

Coordinates of the focus

$$F\left(\frac{p}{2}, 0\right),$$

Coordinates of the vertex

$$M(0, 0).$$

**667.** General Form

$$Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0,$$

where  $B^2 - 4AC = 0$ .

**668.**  $y = ax^2$ ,  $p = \frac{1}{2a}$ .

Equation of the directrix

CHAPTER 7. ANALYTIC GEOMETRY

$$y = -\frac{p}{2},$$

Coordinates of the focus

$$F\left(0, \frac{p}{2}\right),$$

Coordinates of the vertex

$$M(0, 0).$$




Figure 121.

**669.** General Form, Axis Parallel to the y-axis

$$Ax^2 + Dx + Ey + F = 0 \quad (A, E \text{ nonzero}),$$

$$y = ax^2 + bx + c, \quad p = \frac{1}{2a}.$$

Equation of the directrix

$$y = y_0 - \frac{p}{2},$$

Coordinates of the focus

CHAPTER 7. ANALYTIC GEOMETRY

$$F\left(x_0, y_0 + \frac{p}{2}\right),$$

Coordinates of the vertex

$$x_0 = -\frac{b}{2a}, y_0 = ax_0^2 + bx_0 + c = \frac{4ac - b^2}{4a}.$$




Figure 122.

## 7.8 Three-Dimensional Coordinate System

Point coordinates:  $x_0, y_0, z_0, x_1, y_1, z_1, \dots$

Real number:  $\lambda$

Distance between two points:  $d$


Area:  $S$

Volume:  $V$

CHAPTER 7. ANALYTIC GEOMETRY

**670.** Distance Between Two Points

$$d = AB = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$



**Figure 123.**

**671.** Dividing a Line Segment in the Ratio  $\lambda$

$$x_0 = \frac{x_1 + \lambda x_2}{1 + \lambda}, \quad y_0 = \frac{y_1 + \lambda y_2}{1 + \lambda}, \quad z_0 = \frac{z_1 + \lambda z_2}{1 + \lambda},$$

where

$$\lambda = \frac{AC}{CB}, \quad \lambda \neq -1.$$

CHAPTER 7. ANALYTIC GEOMETRY




Figure 124.




Figure 125.

## CHAPTER 7. ANALYTIC GEOMETRY

**672.** Midpoint of a Line Segment

$$x_0 = \frac{x_1 + x_2}{2}, y_0 = \frac{y_1 + y_2}{2}, z_0 = \frac{z_1 + z_2}{2}, \lambda = 1.$$

**673.** Area of a Triangle

The area of a triangle with vertices  $P_1(x_1, y_1, z_1)$ ,  $P_2(x_2, y_2, z_2)$ , and  $P_3(x_3, y_3, z_3)$  is given by

$$S = \frac{1}{2} \sqrt{\begin{vmatrix} y_1 & z_1 & 1 \\ y_2 & z_2 & 1 \\ y_3 & z_3 & 1 \end{vmatrix}^2 + \begin{vmatrix} z_1 & x_1 & 1 \\ z_2 & x_2 & 1 \\ z_3 & x_3 & 1 \end{vmatrix}^2 + \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}^2}.$$

**674.** Volume of a Tetrahedron

The volume of a tetrahedron with vertices  $P_1(x_1, y_1, z_1)$ ,  $P_2(x_2, y_2, z_2)$ ,  $P_3(x_3, y_3, z_3)$ , and  $P_4(x_4, y_4, z_4)$  is given by

$$V = \pm \frac{1}{6} \begin{vmatrix} x_1 & y_1 & z_1 & 1 \\ x_2 & y_2 & z_2 & 1 \\ x_3 & y_3 & z_3 & 1 \\ x_4 & y_4 & z_4 & 1 \end{vmatrix},$$

or

$$V = \pm \frac{1}{6} \begin{vmatrix} x_1 - x_4 & y_1 - y_4 & z_1 - z_4 \\ x_2 - x_4 & y_2 - y_4 & z_2 - z_4 \\ x_3 - x_4 & y_3 - y_4 & z_3 - z_4 \end{vmatrix}.$$

Note: We choose the sign (+) or (-) so that to get a positive answer for volume.

CHAPTER 7. ANALYTIC GEOMETRY




Figure 126.

## 7.9 Plane

Point coordinates:  $x, y, z, x_0, y_0, z_0, x_1, y_1, z_1, \dots$

Real numbers:  $A, B, C, D, A_1, A_2, a, b, c, a_1, a_2, \lambda, p, t, \dots$

Normal vectors:  $\vec{n}, \vec{n}_1, \vec{n}_2$

Direction cosines:  $\cos \alpha, \cos \beta, \cos \gamma$

Distance from point to plane:  $d$

### 675. General Equation of a Plane

$$Ax + By + Cz + D = 0$$

CHAPTER 7. ANALYTIC GEOMETRY

- 676.** Normal Vector to a Plane  
The vector  $\vec{n}(A, B, C)$  is normal to the plane  
 $Ax + By + Cz + D = 0$ .




Figure 127.

- 677.** Particular Cases of the Equation of a Plane  
 $Ax + By + Cz + D = 0$

If  $A = 0$ , the plane is parallel to the x-axis.

If  $B = 0$ , the plane is parallel to the y-axis.

If  $C = 0$ , the plane is parallel to the z-axis.

If  $D = 0$ , the plane lies on the origin.

If  $A = B = 0$ , the plane is parallel to the xy-plane.

If  $B = C = 0$ , the plane is parallel to the yz-plane.

If  $A = C = 0$ , the plane is parallel to the xz-plane.



CHAPTER 7. ANALYTIC GEOMETRY

**678.** Point Direction Form

$$A(x - x_0) + B(y - y_0) + C(z - z_0) = 0,$$

where the point  $P(x_0, y_0, z_0)$  lies in the plane, and the vector  $(A, B, C)$  is normal to the plane.




Figure 128.

**679.** Intercept Form

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

CHAPTER 7. ANALYTIC GEOMETRY




Figure 129.

**680. Three Point Form**

$$\begin{vmatrix} x-x_3 & y-y_3 & z-z_3 \\ x_1-x_3 & y_1-y_3 & z_1-z_3 \\ x_2-x_3 & y_2-y_3 & z_2-z_3 \end{vmatrix} = 0,$$

or

$$\begin{vmatrix} x & y & z & 1 \\ x_1 & y_1 & z_1 & 1 \\ x_2 & y_2 & z_2 & 1 \\ x_3 & y_3 & z_3 & 1 \end{vmatrix} = 0.$$

CHAPTER 7. ANALYTIC GEOMETRY




Figure 130.

**681.** Normal Form

$$x \cos \alpha + y \cos \beta + z \cos \gamma - p = 0,$$

where  $p$  is the perpendicular distance from the origin to the plane, and  $\cos \alpha$ ,  $\cos \beta$ ,  $\cos \gamma$  are the direction cosines of any line normal to the plane.

CHAPTER 7. ANALYTIC GEOMETRY




Figure 131.

**682.** Parametric Form

$$\begin{cases} x = x_1 + a_1s + a_2t \\ y = y_1 + b_1s + b_2t, \\ z = z_1 + c_1s + c_2t \end{cases}$$

where  $(x, y, z)$  are the coordinates of any unknown point on the line, the point  $P(x_1, y_1, z_1)$  lies in the plane, the vectors  $(a_1, b_1, c_1)$  and  $(a_2, b_2, c_2)$  are parallel to the plane.

## CHAPTER 7. ANALYTIC GEOMETRY




Figure 132.

**683.** Dihedral Angle Between Two Planes

If the planes are given by

$$A_1x + B_1y + C_1z + D_1 = 0,$$

$$A_2x + B_2y + C_2z + D_2 = 0,$$

then the dihedral angle between them is

$$\cos \varphi = \frac{\vec{n}_1 \cdot \vec{n}_2}{|\vec{n}_1| |\vec{n}_2|} = \frac{A_1A_2 + B_1B_2 + C_1C_2}{\sqrt{A_1^2 + B_1^2 + C_1^2} \cdot \sqrt{A_2^2 + B_2^2 + C_2^2}}.$$

CHAPTER 7. ANALYTIC GEOMETRY




Figure 133.

**684.** Parallel Planes

Two planes  $A_1x + B_1y + C_1z + D_1 = 0$  and  $A_2x + B_2y + C_2z + D_2 = 0$  are parallel if

$$\frac{A_1}{A_2} = \frac{B_1}{B_2} = \frac{C_1}{C_2}.$$

**685.** Perpendicular Planes

Two planes  $A_1x + B_1y + C_1z + D_1 = 0$  and  $A_2x + B_2y + C_2z + D_2 = 0$  are perpendicular if  $A_1A_2 + B_1B_2 + C_1C_2 = 0$ .


**686.** Equation of a Plane Through  $P(x_1, y_1, z_1)$  and Parallel To the Vectors  $(a_1, b_1, c_1)$  and  $(a_2, b_2, c_2)$  (Fig.132)

## CHAPTER 7. ANALYTIC GEOMETRY

$$\begin{vmatrix} x-x_1 & y-y_1 & z-z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

- 687.** Equation of a Plane Through  $P_1(x_1, y_1, z_1)$  and  $P_2(x_2, y_2, z_2)$ , and Parallel To the Vector  $(a, b, c)$

$$\begin{vmatrix} x-x_1 & y-y_1 & z-z_1 \\ x_2-x_1 & y_2-y_1 & z_2-z_1 \\ a & b & c \end{vmatrix} = 0$$



**Figure 134.**

- 688.** Distance From a Point To a Plane  
The distance from the point  $P_1(x_1, y_1, z_1)$  to the plane  $Ax + By + Cz + D = 0$  is

CHAPTER 7. ANALYTIC GEOMETRY

$$d = \left| \frac{Ax_1 + By_1 + Cz_1 + D}{\sqrt{A^2 + B^2 + C^2}} \right|.$$




Figure 135.

**689.** Intersection of Two Planes

If two planes  $A_1x + B_1y + C_1z + D_1 = 0$  and

$A_2x + B_2y + C_2z + D_2 = 0$  intersect, the intersection straight

line is given by

$$\begin{cases} x = x_1 + at \\ y = y_1 + bt, \\ z = z_1 + ct \end{cases}$$

or

$$\frac{x - x_1}{a} = \frac{y - y_1}{b} = \frac{z - z_1}{c},$$

where



## CHAPTER 7. ANALYTIC GEOMETRY

$$\begin{aligned}
 \mathbf{a} &= \begin{vmatrix} B_1 & C_1 \\ B_2 & C_2 \end{vmatrix}, \quad \mathbf{b} = \begin{vmatrix} C_1 & A_1 \\ C_2 & A_2 \end{vmatrix}, \quad \mathbf{c} = \begin{vmatrix} A_1 & B_1 \\ A_2 & B_2 \end{vmatrix}, \\
 x_1 &= \frac{\mathbf{b} \begin{vmatrix} D_1 & C_1 \\ D_2 & C_2 \end{vmatrix} - \mathbf{c} \begin{vmatrix} D_1 & B_1 \\ D_2 & B_2 \end{vmatrix}}{a^2 + b^2 + c^2}, \\
 y_1 &= \frac{\mathbf{c} \begin{vmatrix} D_1 & A_1 \\ D_2 & A_2 \end{vmatrix} - \mathbf{a} \begin{vmatrix} D_1 & C_1 \\ D_2 & C_2 \end{vmatrix}}{a^2 + b^2 + c^2}, \\
 z_1 &= \frac{\mathbf{a} \begin{vmatrix} D_1 & B_1 \\ D_2 & B_2 \end{vmatrix} - \mathbf{b} \begin{vmatrix} D_1 & A_1 \\ D_2 & A_2 \end{vmatrix}}{a^2 + b^2 + c^2}.
 \end{aligned}$$

## 7.10 Straight Line in Space

Point coordinates:  $x, y, z, x_1, y_1, z_1, \dots$

Direction cosines:  $\cos \alpha, \cos \beta, \cos \gamma$

Real numbers:  $A, B, C, D, a, b, c, a_1, a_2, t, \dots$

Direction vectors of a line:  $\vec{s}, \vec{s}_1, \vec{s}_2$

Normal vector to a plane:  $\vec{n}$

Angle between two lines:  $\varphi$

**690.** Point Direction Form of the Equation of a Line

$$\frac{x - x_1}{a} = \frac{y - y_1}{b} = \frac{z - z_1}{c},$$

where the point  $P_1(x_1, y_1, z_1)$  lies on the line, and  $(a, b, c)$  is the direction vector of the line.

CHAPTER 7. ANALYTIC GEOMETRY




Figure 136.

691. Two Point Form

$$\frac{x-x_1}{x_2-x_1} = \frac{y-y_1}{y_2-y_1} = \frac{z-z_1}{z_2-z_1}$$




Figure 137.

## CHAPTER 7. ANALYTIC GEOMETRY

**692. Parametric Form**

$$\begin{cases} x = x_1 + t \cos \alpha \\ y = y_1 + t \cos \beta \\ z = z_1 + t \cos \gamma \end{cases}$$

where the point  $P_1(x_1, y_1, z_1)$  lies on the straight line,  $\cos \alpha$ ,  $\cos \beta$ ,  $\cos \gamma$  are the direction cosines of the direction vector of the line, the parameter  $t$  is any real number.




Figure 138.

**693. Angle Between Two Straight Lines**

$$\cos \varphi = \frac{\vec{s}_1 \cdot \vec{s}_2}{|\vec{s}_1| \cdot |\vec{s}_2|} = \frac{a_1 a_2 + b_1 b_2 + c_1 c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \cdot \sqrt{a_2^2 + b_2^2 + c_2^2}}$$

CHAPTER 7. ANALYTIC GEOMETRY




Figure 139.

**694.** Parallel Lines

Two lines are parallel if

$$\vec{s}_1 \parallel \vec{s}_2,$$

or

$$\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}.$$

**695.** Perpendicular Lines

Two lines are perpendicular if

$$\vec{s}_1 \cdot \vec{s}_2 = 0,$$

or

$$a_1a_2 + b_1b_2 + c_1c_2 = 0.$$

**696.** Intersection of Two Lines

Two lines  $\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1}$  and

## CHAPTER 7. ANALYTIC GEOMETRY

$\frac{x-x_2}{a_2} = \frac{y-y_2}{b_2} = \frac{z-z_2}{c_2}$  intersect if

$$\begin{vmatrix} x_2-x_1 & y_2-y_1 & z_2-z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0.$$

**697.** Parallel Line and Plane

The straight line  $\frac{x-x_1}{a} = \frac{y-y_1}{b} = \frac{z-z_1}{c}$  and the plane

$Ax + By + Cz + D = 0$  are parallel if

$$\vec{n} \cdot \vec{s} = 0,$$

or

$$Aa + Bb + Cc = 0.$$




Figure 140.

CHAPTER 7. ANALYTIC GEOMETRY

**698.** Perpendicular Line and Plane

The straight line  $\frac{x-x_1}{a} = \frac{y-y_1}{b} = \frac{z-z_1}{c}$  and the plane

$Ax + By + Cz + D = 0$  are perpendicular if

$\vec{n} \parallel \vec{s}$ ,

or

$$\frac{A}{a} = \frac{B}{b} = \frac{C}{c}.$$




Figure 141.

## 7.11 Quadric Surfaces

Point coordinates of the quadric surfaces:  $x, y, z$

Real numbers:  $A, B, C, a, b, c, k_1, k_2, k_3, \dots$

## CHAPTER 7. ANALYTIC GEOMETRY

**699.** General Quadratic Equation

$$Ax^2 + By^2 + Cz^2 + 2Fyz + 2Gzx + 2Hxy + 2Px + 2Qy + 2Rz + D = 0$$

**700.** Classification of Quadric Surfaces

Case	Rank(e)	Rank(E)	$\Delta$	k signs	Type of Surface
1	3	4	$< 0$	Same	Real Ellipsoid
2	3	4	$> 0$	Same	Imaginary Ellipsoid
3	3	4	$> 0$	Different	Hyperboloid of 1 Sheet
4	3	4	$< 0$	Different	Hyperboloid of 2 Sheets
5	3	3		Different	Real Quadric Cone
6	3	3		Same	Imaginary Quadric Cone
7	2	4	$< 0$	Same	Elliptic Paraboloid
8	2	4	$> 0$	Different	Hyperbolic Paraboloid
9	2	3		Same	Real Elliptic Cylinder
10	2	3		Same	Imaginary Elliptic Cylinder
11	2	3		Different	Hyperbolic Cylinder
12	2	2		Different	Real Intersecting Planes
13	2	2		Same	Imaginary Intersecting Planes
14	1	3			Parabolic Cylinder
15	1	2			Real Parallel Planes
16	1	2			Imaginary Parallel Planes
17	1	1			Coincident Planes

Here

$$e = \begin{pmatrix} A & H & G \\ H & B & F \\ G & F & C \end{pmatrix}, E = \begin{pmatrix} A & H & Q & P \\ H & B & F & Q \\ G & F & C & R \\ P & Q & R & D \end{pmatrix}, \Delta = \det(E),$$


$k_1, k_2, k_3$  are the roots of the equation,

$$\begin{vmatrix} A-x & H & G \\ H & B-x & F \\ G & F & C-x \end{vmatrix} = 0.$$

CHAPTER 7. ANALYTIC GEOMETRY

**701.** Real Ellipsoid (Case 1)

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$$



**Figure 142.**

**702.** Imaginary Ellipsoid (Case 2)

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = -1$$

**703.** Hyperboloid of 1 Sheet (Case 3)

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1$$



CHAPTER 7. ANALYTIC GEOMETRY




Figure 143.

704. Hyperboloid of 2 Sheets (Case 4)

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = -1$$





Figure 144.

CHAPTER 7. ANALYTIC GEOMETRY

**705.** Real Quadric Cone (Case 5)

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 0$$



**Figure 145.**

**706.** Imaginary Quadric Cone (Case 6)

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 0$$

**707.** Elliptic Paraboloid (Case 7)

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - z = 0$$

CHAPTER 7. ANALYTIC GEOMETRY




Figure 146.

708. Hyperbolic Paraboloid (Case 8)

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} - z = 0$$





Figure 147.

CHAPTER 7. ANALYTIC GEOMETRY

**709.** Real Elliptic Cylinder (Case 9)

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$



**Figure 148.**

**710.** Imaginary Elliptic Cylinder (Case 10)

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = -1$$

**711.** Hyperbolic Cylinder (Case 11)

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$$

CHAPTER 7. ANALYTIC GEOMETRY




Figure 149.

**712.** Real Intersecting Planes (Case 12)

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 0$$

**713.** Imaginary Intersecting Planes (Case 13)

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 0$$

**714.** Parabolic Cylinder (Case 14)

$$\frac{x^2}{a^2} - y = 0$$

CHAPTER 7. ANALYTIC GEOMETRY




Figure 150.

**715.** Real Parallel Planes (Case 15)

$$\frac{x^2}{a^2} = 1$$

**716.** Imaginary Parallel Planes (Case 16)

$$\frac{x^2}{a^2} = -1$$

**717.** Coincident Planes (Case 17)

$$x^2 = 0$$

CHAPTER 7. ANALYTIC GEOMETRY

## 7.12 Sphere

Radius of a sphere:  $R$


Point coordinates:  $x, y, z, x_1, y_1, z_1, \dots$

Center of a sphere:  $(a, b, c)$

Real numbers:  $A, D, E, F, M$

- 718.** Equation of a Sphere Centered at the Origin (Standard Form)

$$x^2 + y^2 + z^2 = R^2$$



**Figure 151.**

- 719.** Equation of a Circle Centered at Any Point  $(a, b, c)$

$$(x - a)^2 + (y - b)^2 + (z - c)^2 = R^2$$

- 720.** Diameter Form

$$(x - x_1)(x - x_2) + (y - y_1)(y - y_2) + (z - z_1)(z - z_2) = 0,$$

CHAPTER 7. ANALYTIC GEOMETRY

where

$P_1(x_1, y_1, z_1)$ ,  $P_2(x_2, y_2, z_2)$  are the ends of a diameter.

**721.** Four Point Form

$$\begin{vmatrix} x^2 + y^2 + z^2 & x & y & z & 1 \\ x_1^2 + y_1^2 + z_1^2 & x_1 & y_1 & z_1 & 1 \\ x_2^2 + y_2^2 + z_2^2 & x_2 & y_2 & z_2 & 1 \\ x_3^2 + y_3^2 + z_3^2 & x_3 & y_3 & z_3 & 1 \\ x_4^2 + y_4^2 + z_4^2 & x_4 & y_4 & z_4 & 1 \end{vmatrix} = 0$$

**722.** General Form

$Ax^2 + Ay^2 + Az^2 + Dx + Ey + Fz + M = 0$  ( $A$  is nonzero).

The center of the sphere has coordinates  $(a, b, c)$ , where

$$a = -\frac{D}{2A}, \quad b = -\frac{E}{2A}, \quad c = -\frac{F}{2A}.$$

The radius of the sphere is

$$R = \frac{\sqrt{D^2 + E^2 + F^2 - 4A^2M}}{2A}.$$



## **Chapter 8**

# **Differential Calculus**

Functions:  $f, g, y, u, v$   
Argument (independent variable):  $x$   
Real numbers:  $a, b, c, d$   
Natural number:  $n$   
Angle:  $\alpha$   
Inverse function:  $f^{-1}$

### 8.1 Functions and Their Graphs

**723.** Even Function

$$f(-x) = f(x)$$

**724.** Odd Function

$$f(-x) = -f(x)$$

**725.** Periodic Function

$$f(x + nT) = f(x)$$

**726.** Inverse Function

$y = f(x)$  is any function,  $x = g(y)$  or  $y = f^{-1}(x)$  is its inverse function.

CHAPTER 8. DIFFERENTIAL CALCULUS




Figure 152.

**727.** Composite Function

$y = f(u)$ ,  $u = g(x)$ ,  $y = f(g(x))$  is a composite function.

**728.** Linear Function

$y = ax + b$ ,  $x \in \mathbb{R}$ ,  $a = \tan \alpha$  is the slope of the line,  $b$  is the y-intercept.

CHAPTER 8. DIFFERENTIAL CALCULUS




Figure 153.

**729.** Quadratic Function  
 $y = x^2, x \in \mathbb{R}$ .




Figure 154.

CHAPTER 8. DIFFERENTIAL CALCULUS

730.  $y = ax^2 + bx + c, x \in \mathbb{R}.$




Figure 155.

731. Cubic Function  
 $y = x^3, x \in \mathbb{R}.$

CHAPTER 8. DIFFERENTIAL CALCULUS




Figure 156.

732.  $y = ax^3 + bx^2 + cx + d, x \in \mathbb{R}.$

CHAPTER 8. DIFFERENTIAL CALCULUS




Figure 157.

**733.** Power Function  
 $y = x^n$ ,  $n \in \mathbb{N}$ .

CHAPTER 8. DIFFERENTIAL CALCULUS




Figure 158.




Figure 159.

CHAPTER 8. DIFFERENTIAL CALCULUS

**734. Square Root Function**

$$y = \sqrt{x}, \quad x \in [0, \infty).$$




Figure 160.

**735. Exponential Functions**

$$y = a^x, \quad a > 0, \quad a \neq 1,$$

$$y = e^x \text{ if } a = e, \quad e = 2.71828182846\dots$$




Figure 161.



CHAPTER 8. DIFFERENTIAL CALCULUS

- 736.** Logarithmic Functions  
 $y = \log_a x$ ,  $x \in (0, \infty)$ ,  $a > 0$ ,  $a \neq 1$ ,  
 $y = \ln x$  if  $a = e$ ,  $x > 0$ .




Figure 162.

- 737.** Hyperbolic Sine Function  
 $y = \sinh x$ ,  $\sinh x = \frac{e^x - e^{-x}}{2}$ ,  $x \in \mathbb{R}$ .

CHAPTER 8. DIFFERENTIAL CALCULUS




Figure 163.

**738.** Hyperbolic Cosine Function

$$y = \cosh x, \cosh x = \frac{e^x + e^{-x}}{2}, x \in \mathbb{R}.$$




Figure 164.

CHAPTER 8. DIFFERENTIAL CALCULUS

**739.** Hyperbolic Tangent Function

$$y = \tanh x, \quad y = \tanh x = \frac{\sinh x}{\cosh x} = \frac{e^x - e^{-x}}{e^x + e^{-x}}, \quad x \in \mathbb{R}.$$




Figure 165.

**740.** Hyperbolic Cotangent Function

$$y = \coth x, \quad y = \coth x = \frac{\cosh x}{\sinh x} = \frac{e^x + e^{-x}}{e^x - e^{-x}}, \quad x \in \mathbb{R}, \quad x \neq 0.$$

CHAPTER 8. DIFFERENTIAL CALCULUS




Figure 166.

**741.** Hyperbolic Secant Function

$$y = \operatorname{sech} x, \quad y = \operatorname{sech} x = \frac{1}{\cosh x} = \frac{2}{e^x + e^{-x}}, \quad x \in \mathbb{R}.$$





Figure 167.

CHAPTER 8. DIFFERENTIAL CALCULUS

**742.** Hyperbolic Cosecant Function

$$y = \operatorname{csch} x, \quad y = \operatorname{csch} x = \frac{1}{\sinh x} = \frac{2}{e^x - e^{-x}}, \quad x \in \mathbf{R}, \quad x \neq 0.$$



**Figure 168.**

**743.** Inverse Hyperbolic Sine Function

$$y = \operatorname{arcsinh} x, \quad x \in \mathbf{R}.$$

CHAPTER 8. DIFFERENTIAL CALCULUS




Figure 169.

- 744.** Inverse Hyperbolic Cosine Function  
 $y = \operatorname{arccosh} x$ ,  $x \in [1, \infty)$ .




Figure 170.

- 745.** Inverse Hyperbolic Tangent Function  
 $y = \operatorname{arctanh} x$ ,  $x \in (-1, 1)$ .

CHAPTER 8. DIFFERENTIAL CALCULUS




Figure 171.

**746.** Inverse Hyperbolic Cotangent Function  
 $y = \operatorname{arccoth} x$ ,  $x \in (-\infty, -1) \cup (1, \infty)$ .

CHAPTER 8. DIFFERENTIAL CALCULUS




Figure 172.

**747.** Inverse Hyperbolic Secant Function  
 $y = \operatorname{arcsech} x$ ,  $x \in (0, 1]$ .



CHAPTER 8. DIFFERENTIAL CALCULUS




Figure 173.

- 748.** Inverse Hyperbolic Cosecant Function  
 $y = \text{arccsch } x, x \in \mathbb{R}, x \neq 0.$




Figure 174.

## 8.2 Limits of Functions

Functions:  $f(x)$ ,  $g(x)$ Argument:  $x$ Real constants:  $a$ ,  $k$ 

$$749. \lim_{x \rightarrow a} [f(x) + g(x)] = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x)$$

$$750. \lim_{x \rightarrow a} [f(x) - g(x)] = \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x)$$

$$751. \lim_{x \rightarrow a} [f(x) \cdot g(x)] = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x)$$

$$752. \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}, \text{ if } \lim_{x \rightarrow a} g(x) \neq 0.$$

$$753. \lim_{x \rightarrow a} [kf(x)] = k \lim_{x \rightarrow a} f(x)$$

$$754. \lim_{x \rightarrow a} f(g(x)) = f(\lim_{x \rightarrow a} g(x))$$

$$755. \lim_{x \rightarrow a} f(x) = f(a), \text{ if the function } f(x) \text{ is continuous at } x = a.$$

$$756. \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$$

$$757. \lim_{x \rightarrow 0} \frac{\tan x}{x} = 1$$

$$758. \lim_{x \rightarrow 0} \frac{\sin^{-1} x}{x} = 1$$

CHAPTER 8. DIFFERENTIAL CALCULUS

$$759. \lim_{x \rightarrow 0} \frac{\tan^{-1} x}{x} = 1$$

$$760. \lim_{x \rightarrow 0} \frac{\ln(1+x)}{x} = 1$$

$$761. \lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x = e$$

$$762. \lim_{x \rightarrow \infty} \left(1 + \frac{k}{x}\right)^x = e^k$$

$$763. \lim_{x \rightarrow 0} a^x = 1$$

### 8.3 Definition and Properties of the Derivative

Functions:  $f, g, y, u, v$

Independent variable:  $x$

Real constant:  $k$

Angle:  $\alpha$

$$764. y'(x) = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x} = \frac{dy}{dx}$$

CHAPTER 8. DIFFERENTIAL CALCULUS




Figure 175.

765.  $\frac{dy}{dx} = \tan \alpha$

766.  $\frac{d(u + v)}{dx} = \frac{du}{dx} + \frac{dv}{dx}$

767.  $\frac{d(u - v)}{dx} = \frac{du}{dx} - \frac{dv}{dx}$

768.  $\frac{d(ku)}{dx} = k \frac{du}{dx}$

769. Product Rule  
 $\frac{d(u \cdot v)}{dx} = \frac{du}{dx} \cdot v + u \cdot \frac{dv}{dx}$

CHAPTER 8. DIFFERENTIAL CALCULUS

**770.** Quotient Rule

$$\frac{d}{dx} \left( \frac{u}{v} \right) = \frac{\frac{du}{dx} \cdot v - u \cdot \frac{dv}{dx}}{v^2}$$

**771.** Chain Rule

$$y = f(g(x)), \quad u = g(x),$$

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}.$$

**772.** Derivative of Inverse Function

$$\frac{dy}{dx} = \frac{1}{\frac{dx}{dy}},$$

where  $x(y)$  is the inverse function of  $y(x)$ .

**773.** Reciprocal Rule

$$\frac{d}{dx} \left( \frac{1}{y} \right) = -\frac{\frac{dy}{dx}}{y^2}$$

**774.** Logarithmic Differentiation

$$y = f(x), \quad \ln y = \ln f(x),$$

$$\frac{dy}{dx} = f(x) \cdot \frac{d}{dx} [\ln f(x)].$$

## 8.4 Table of Derivatives

Independent variable:  $x$

Real constants:  $C, a, b, c$

Natural number:  $n$

CHAPTER 8. DIFFERENTIAL CALCULUS

$$775. \frac{d}{dx}(C) = 0$$

$$776. \frac{d}{dx}(x) = 1$$

$$777. \frac{d}{dx}(ax + b) = a$$

$$778. \frac{d}{dx}(ax^2 + bx + c) = 2ax + b$$

$$779. \frac{d}{dx}(x^n) = nx^{n-1}$$

$$780. \frac{d}{dx}(x^{-n}) = -\frac{n}{x^{n+1}}$$

$$781. \frac{d}{dx}\left(\frac{1}{x}\right) = -\frac{1}{x^2}$$

$$782. \frac{d}{dx}(\sqrt{x}) = \frac{1}{2\sqrt{x}}$$

$$783. \frac{d}{dx}(\sqrt[n]{x}) = \frac{1}{n\sqrt[n]{x^{n-1}}}$$

$$784. \frac{d}{dx}(\ln x) = \frac{1}{x}$$

$$785. \frac{d}{dx}(\log_a x) = \frac{1}{x \ln a}, \quad a > 0, \quad a \neq 1.$$

CHAPTER 8. DIFFERENTIAL CALCULUS

$$786. \frac{d}{dx}(a^x) = a^x \ln a, \quad a > 0, \quad a \neq 1.$$

$$787. \frac{d}{dx}(e^x) = e^x$$

$$788. \frac{d}{dx}(\sin x) = \cos x$$

$$789. \frac{d}{dx}(\cos x) = -\sin x$$

$$790. \frac{d}{dx}(\tan x) = \frac{1}{\cos^2 x} = \sec^2 x$$

$$791. \frac{d}{dx}(\cot x) = -\frac{1}{\sin^2 x} = -\csc^2 x$$

$$792. \frac{d}{dx}(\sec x) = \tan x \cdot \sec x$$

$$793. \frac{d}{dx}(\csc x) = -\cot x \cdot \csc x$$

$$794. \frac{d}{dx}(\arcsin x) = \frac{1}{\sqrt{1-x^2}}$$

$$795. \frac{d}{dx}(\arccos x) = -\frac{1}{\sqrt{1-x^2}}$$

$$796. \frac{d}{dx}(\arctan x) = \frac{1}{1+x^2}$$

CHAPTER 8. DIFFERENTIAL CALCULUS

$$797. \frac{d}{dx}(\operatorname{arccot} x) = -\frac{1}{1+x^2}$$

$$798. \frac{d}{dx}(\operatorname{arcsec} x) = \frac{1}{|x|\sqrt{x^2-1}}$$

$$799. \frac{d}{dx}(\operatorname{arccsc} x) = -\frac{1}{|x|\sqrt{x^2-1}}$$

$$800. \frac{d}{dx}(\sinh x) = \cosh x$$

$$801. \frac{d}{dx}(\cosh x) = \sinh x$$

$$802. \frac{d}{dx}(\tanh x) = \frac{1}{\cosh^2 x} = \operatorname{sech}^2 x$$

$$803. \frac{d}{dx}(\operatorname{coth} x) = -\frac{1}{\sinh^2 x} = -\operatorname{csch}^2 x$$

$$804. \frac{d}{dx}(\operatorname{sech} x) = -\operatorname{sech} x \cdot \tanh x$$

$$805. \frac{d}{dx}(\operatorname{csch} x) = -\operatorname{csch} x \cdot \operatorname{coth} x$$

$$806. \frac{d}{dx}(\operatorname{arcsinh} x) = \frac{1}{\sqrt{x^2+1}}$$

$$807. \frac{d}{dx}(\operatorname{arccosh} x) = \frac{1}{\sqrt{x^2-1}}$$



## CHAPTER 8. DIFFERENTIAL CALCULUS

$$808. \frac{d}{dx}(\operatorname{arctanh} x) = \frac{1}{1-x^2}, |x| < 1.$$

$$809. \frac{d}{dx}(\operatorname{arccoth} x) = -\frac{1}{x^2-1}, |x| > 1.$$

$$810. \frac{d}{dx}(u^v) = vu^{v-1} \cdot \frac{du}{dx} + u^v \ln u \cdot \frac{dv}{dx}$$

## 8.5 Higher Order Derivatives

Functions:  $f, y, u, v$

Independent variable:  $x$

Natural number:  $n$

811. Second derivative

$$f'' = (f')' = \left(\frac{dy}{dx}\right)' = \frac{d}{dx}\left(\frac{dy}{dx}\right) = \frac{d^2y}{dx^2}$$

812. Higher-Order derivative

$$f^{(n)} = \frac{d^n y}{dx^n} = y^{(n)} = (f^{(n-1)})'$$

$$813. (u+v)^{(n)} = u^{(n)} + v^{(n)}$$

$$814. (u-v)^{(n)} = u^{(n)} - v^{(n)}$$

815. Leibnitz's Formulas

$$(uv)'' = u''v + 2u'v' + uv''$$

## CHAPTER 8. DIFFERENTIAL CALCULUS

$$(\mathbf{uv})''' = \mathbf{u}'''\mathbf{v} + 3\mathbf{u}''\mathbf{v}' + 3\mathbf{u}'\mathbf{v}'' + \mathbf{uv}'''$$

$$(\mathbf{uv})^{(n)} = \mathbf{u}^{(n)}\mathbf{v} + n\mathbf{u}^{(n-1)}\mathbf{v}' + \frac{n(n-1)}{1 \cdot 2}\mathbf{u}^{(n-2)}\mathbf{v}'' + \dots + \mathbf{uv}^{(n)}$$

$$816. \quad (\mathbf{x}^m)^{(n)} = \frac{\mathbf{m}!}{(\mathbf{m}-n)!} \mathbf{x}^{m-n}$$

$$817. \quad (\mathbf{x}^n)^{(n)} = n!$$

$$818. \quad (\log_a \mathbf{x})^{(n)} = \frac{(-1)^{n-1}(n-1)!}{\mathbf{x}^n \ln a}$$

$$819. \quad (\ln \mathbf{x})^{(n)} = \frac{(-1)^{n-1}(n-1)!}{\mathbf{x}^n}$$

$$820. \quad (\mathbf{a}^{\mathbf{x}})^{(n)} = \mathbf{a}^{\mathbf{x}} \ln^n \mathbf{a}$$

$$821. \quad (\mathbf{e}^{\mathbf{x}})^{(n)} = \mathbf{e}^{\mathbf{x}}$$

$$822. \quad (\mathbf{a}^{\mathbf{mx}})^{(n)} = \mathbf{m}^n \mathbf{a}^{\mathbf{mx}} \ln^n \mathbf{a}$$

$$823. \quad (\sin \mathbf{x})^{(n)} = \sin\left(\mathbf{x} + \frac{n\pi}{2}\right)$$

$$824. \quad (\cos \mathbf{x})^{(n)} = \cos\left(\mathbf{x} + \frac{n\pi}{2}\right)$$

## 8.6 Applications of Derivative

Functions:  $f, g, y$

Position of an object:  $s$

Velocity:  $v$

Acceleration:  $w$

Independent variable:  $x$

Time:  $t$

Natural number:  $n$

### 825. Velocity and Acceleration

$s = f(t)$  is the position of an object relative to a fixed coordinate system at a time  $t$ ,

$v = s' = f'(t)$  is the instantaneous velocity of the object,

$w = v' = s'' = f''(t)$  is the instantaneous acceleration of the object.

### 826. Tangent Line

$$y - y_0 = f'(x_0)(x - x_0)$$

CHAPTER 8. DIFFERENTIAL CALCULUS




Figure 176.

**827.** Normal Line

$$y - y_0 = -\frac{1}{f'(x_0)}(x - x_0) \text{ (Fig 176)}$$

**828.** Increasing and Decreasing Functions.

If  $f'(x_0) > 0$ , then  $f(x)$  is increasing at  $x_0$ . (Fig 177,  $x < x_1$ ,  $x_2 < x$ ),

If  $f'(x_0) < 0$ , then  $f(x)$  is decreasing at  $x_0$ . (Fig 177,  $x_1 < x < x_2$ ),

If  $f'(x_0)$  does not exist or is zero, then the test fails.

CHAPTER 8. DIFFERENTIAL CALCULUS




Figure 177.

**829.** Local extrema

A function  $f(x)$  has a **local maximum** at  $x_1$  if and only if there exists some interval containing  $x_1$  such that  $f(x_1) \geq f(x)$  for all  $x$  in the interval (Fig.177).

A function  $f(x)$  has a **local minimum** at  $x_2$  if and only if there exists some interval containing  $x_2$  such that  $f(x_2) \leq f(x)$  for all  $x$  in the interval (Fig.177).

**830.** Critical Points

A critical point on  $f(x)$  occurs at  $x_0$  if and only if either  $f'(x_0)$  is zero or the derivative doesn't exist.

**831.** First Derivative Test for Local Extrema.

If  $f(x)$  is increasing ( $f'(x) > 0$ ) for all  $x$  in some interval  $(a, x_1]$  and  $f(x)$  is decreasing ( $f'(x) < 0$ ) for all  $x$  in some interval  $[x_1, b)$ , then  $f(x)$  has a local maximum at  $x_1$  (Fig.177).

CHAPTER 8. DIFFERENTIAL CALCULUS

- 832.** If  $f(x)$  is decreasing ( $f'(x) < 0$ ) for all  $x$  in some interval  $(a, x_2]$  and  $f(x)$  is increasing ( $f'(x) > 0$ ) for all  $x$  in some interval  $[x_2, b)$ , then  $f(x)$  has a local minimum at  $x_2$ . (Fig.177).
- 833.** Second Derivative Test for Local Extrema.  
If  $f'(x_1) = 0$  and  $f''(x_1) < 0$ , then  $f(x)$  has a local maximum at  $x_1$ .  
If  $f'(x_2) = 0$  and  $f''(x_2) > 0$ , then  $f(x)$  has a local minimum at  $x_2$ . (Fig.177)
- 834.** Concavity.  
 $f(x)$  is concave upward at  $x_0$  if and only if  $f'(x)$  is increasing at  $x_0$  (Fig.177,  $x_3 < x$ ).  
 $f(x)$  is concave downward at  $x_0$  if and only if  $f'(x)$  is decreasing at  $x_0$ . (Fig.177,  $x < x_3$ ).
- 835.** Second Derivative Test for Concavity.  
If  $f''(x_0) > 0$ , then  $f(x)$  is concave upward at  $x_0$ .  
If  $f''(x_0) < 0$ , then  $f(x)$  is concave downward at  $x_0$ .  
If  $f''(x)$  does not exist or is zero, then the test fails.
- 836.** Inflection Points  
If  $f'(x_3)$  exists and  $f''(x)$  changes sign at  $x = x_3$ , then the point  $(x_3, f(x_3))$  is an **inflection point** of the graph of  $f(x)$ . If  $f''(x_3)$  exists at the inflection point, then  $f''(x_3) = 0$  (Fig.177).
- 837.** L'Hopital's Rule
- $$\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \lim_{x \rightarrow c} \frac{f'(x)}{g'(x)} \text{ if } \lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} g(x) = \begin{cases} 0 \\ \infty \end{cases} .$$

## 8.7 Differential

Functions:  $f, u, v$

Independent variable:  $x$

Derivative of a function:  $y'(x), f'(x)$

Real constant:  $C$

Differential of function  $y = f(x)$ :  $dy$

Differential of  $x$ :  $dx$

Small change in  $x$ :  $\Delta x$

Small change in  $y$ :  $\Delta y$

**838.**  $dy = y' dx$

**839.**  $f(x + \Delta x) = f(x) + f'(x)\Delta x$




Figure 178.

CHAPTER 8. DIFFERENTIAL CALCULUS

**840.** Small Change in  $y$   
 $\Delta y = f(x + \Delta x) - f(x)$

**841.**  $d(u + v) = du + dv$

**842.**  $d(u - v) = du - dv$

**843.**  $d(Cu) = Cdu$

**844.**  $d(uv) = vdu + udv$

**845.**  $d\left(\frac{u}{v}\right) = \frac{vdu - udv}{v^2}$

## 8.8 Multivariable Functions

Functions of two variables:  $z(x, y)$ ,  $f(x, y)$ ,  $g(x, y)$ ,  $h(x, y)$

Arguments:  $x, y, t$

Small changes in  $x, y, z$ , respectively:  $\Delta x, \Delta y, \Delta z$ .

**846.** First Order Partial Derivatives

The partial derivative with respect to  $x$

$$\frac{\partial f}{\partial x} = f_x \quad (\text{also } \frac{\partial z}{\partial x} = z_x),$$

The partial derivative with respect to  $y$

$$\frac{\partial f}{\partial y} = f_y \quad (\text{also } \frac{\partial z}{\partial y} = z_y).$$



## CHAPTER 8. DIFFERENTIAL CALCULUS

**847.** Second Order Partial Derivatives

$$\frac{\partial}{\partial x} \left( \frac{\partial f}{\partial x} \right) = \frac{\partial^2 f}{\partial x^2} = f_{xx},$$

$$\frac{\partial}{\partial y} \left( \frac{\partial f}{\partial y} \right) = \frac{\partial^2 f}{\partial y^2} = f_{yy},$$

$$\frac{\partial}{\partial y} \left( \frac{\partial f}{\partial x} \right) = \frac{\partial^2 f}{\partial y \partial x} = f_{xy},$$

$$\frac{\partial}{\partial x} \left( \frac{\partial f}{\partial y} \right) = \frac{\partial^2 f}{\partial x \partial y} = f_{yx}.$$

If the derivatives are continuous, then

$$\frac{\partial^2 f}{\partial y \partial x} = \frac{\partial^2 f}{\partial x \partial y}.$$

**848.** Chain Rules

If  $f(x, y) = g(h(x, y))$  ( $g$  is a function of one variable  $h$ ), then

$$\frac{\partial f}{\partial x} = g'(h(x, y)) \frac{\partial h}{\partial x}, \quad \frac{\partial f}{\partial y} = g'(h(x, y)) \frac{\partial h}{\partial y}.$$

$$\text{If } h(t) = f(x(t), y(t)), \text{ then } h'(t) = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}.$$

If  $z = f(x(u, v), y(u, v))$ , then

$$\frac{\partial z}{\partial u} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial u} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial u}, \quad \frac{\partial z}{\partial v} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial v} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial v}.$$

**849.** Small Changes

$$\Delta z \approx \frac{\partial f}{\partial x} \Delta x + \frac{\partial f}{\partial y} \Delta y$$

## CHAPTER 8. DIFFERENTIAL CALCULUS

**850.** Local Maxima and Minima

$f(x, y)$  has a **local maximum** at  $(x_0, y_0)$  if  $f(x, y) \leq f(x_0, y_0)$  for all  $(x, y)$  sufficiently close to  $(x_0, y_0)$ .

$f(x, y)$  has a **local minimum** at  $(x_0, y_0)$  if  $f(x, y) \geq f(x_0, y_0)$  for all  $(x, y)$  sufficiently close to  $(x_0, y_0)$ .

**851.** Stationary Points

$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial y} = 0.$$

Local maxima and local minima occur at stationary points.

**852.** Saddle Point

A stationary point which is neither a local maximum nor a local minimum

**853.** Second Derivative Test for Stationary Points

Let  $(x_0, y_0)$  be a stationary point ( $\frac{\partial f}{\partial x} = \frac{\partial f}{\partial y} = 0$ ).

$$D = \begin{vmatrix} f_{xx}(x_0, y_0) & f_{xy}(x_0, y_0) \\ f_{yx}(x_0, y_0) & f_{yy}(x_0, y_0) \end{vmatrix}.$$

If  $D > 0$ ,  $f_{xx}(x_0, y_0) > 0$ ,  $(x_0, y_0)$  is a point of local minima.

If  $D > 0$ ,  $f_{xx}(x_0, y_0) < 0$ ,  $(x_0, y_0)$  is a point of local maxima.

If  $D < 0$ ,  $(x_0, y_0)$  is a saddle point.

If  $D = 0$ , the test fails.

**854.** Tangent Plane

The equation of the tangent plane to the surface  $z = f(x, y)$  at  $(x_0, y_0, z_0)$  is

$$z - z_0 = f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0).$$

## CHAPTER 8. DIFFERENTIAL CALCULUS

**855.** Normal to Surface

The equation of the normal to the surface  $z = f(x, y)$  at  $(x_0, y_0, z_0)$  is

$$\frac{x - x_0}{f_x(x_0, y_0)} = \frac{y - y_0}{f_y(x_0, y_0)} = \frac{z - z_0}{-1}.$$

## 8.9 Differential Operators

Unit vectors along the coordinate axes:  $\vec{i}$ ,  $\vec{j}$ ,  $\vec{k}$

Scalar functions (scalar fields):  $f(x, y, z)$ ,  $u(x_1, x_2, \dots, x_n)$

Gradient of a scalar field:  $\text{grad } u$ ,  $\nabla u$

Directional derivative:  $\frac{\partial f}{\partial l}$

Vector function (vector field):  $\vec{F}(P, Q, R)$

Divergence of a vector field:  $\text{div } \vec{F}$ ,  $\nabla \cdot \vec{F}$

Curl of a vector field:  $\text{curl } \vec{F}$ ,  $\nabla \times \vec{F}$

Laplacian operator:  $\nabla^2$

**856.** Gradient of a Scalar Function

$$\text{grad } f = \nabla f = \left( \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right),$$

$$\text{grad } u = \nabla u = \left( \frac{\partial u}{\partial x_1}, \frac{\partial u}{\partial x_2}, \dots, \frac{\partial u}{\partial x_n} \right).$$

**857.** Directional Derivative

$$\frac{\partial f}{\partial l} = \frac{\partial f}{\partial x} \cos \alpha + \frac{\partial f}{\partial y} \cos \beta + \frac{\partial f}{\partial z} \cos \gamma,$$

## CHAPTER 8. DIFFERENTIAL CALCULUS

where the direction is defined by the vector  
 $\vec{l}(\cos \alpha, \cos \beta, \cos \gamma)$ ,  $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$ .

**858.** Divergence of a Vector Field

$$\operatorname{div} \vec{F} = \nabla \cdot \vec{F} = \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z}$$

**859.** Curl of a Vector Field

$$\operatorname{curl} \vec{F} = \nabla \times \vec{F} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ P & Q & R \end{vmatrix}$$

$$= \left( \frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z} \right) \vec{i} + \left( \frac{\partial P}{\partial z} - \frac{\partial R}{\partial x} \right) \vec{j} + \left( \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \vec{k}$$

**860.** Laplacian Operator

$$\nabla^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}$$

$$\mathbf{861.} \quad \operatorname{div}(\operatorname{curl} \vec{F}) = \nabla \cdot (\nabla \times \vec{F}) \equiv 0$$

$$\mathbf{862.} \quad \operatorname{curl}(\operatorname{grad} f) = \nabla \times (\nabla f) \equiv 0$$

$$\mathbf{863.} \quad \operatorname{div}(\operatorname{grad} f) = \nabla \cdot (\nabla f) = \nabla^2 f$$

$$\mathbf{864.} \quad \operatorname{curl}(\operatorname{curl} \vec{F}) = \operatorname{grad}(\operatorname{div} \vec{F}) - \nabla^2 \vec{F} = \nabla(\nabla \cdot \vec{F}) - \nabla^2 \vec{F}$$

## Chapter 9

# Integral Calculus

Functions:  $f, g, u, v$

Independent variables:  $x, t, \xi$

Indefinite integral of a function:  $\int f(x)dx, \int g(x)dx, \dots$

Derivative of a function:  $y'(x), f'(x), F'(x), \dots$

Real constants:  $C, a, b, c, d, k$

Natural numbers:  $m, n, i, j$

### 9.1 Indefinite Integral

$$865. \int f(x)dx = F(x) + C \text{ if } F'(x) = f(x).$$

$$866. \left( \int f(x)dx \right)' = f(x)$$

$$867. \int kf(x)dx = k \int f(x)dx$$

$$868. \int [f(x) + g(x)]dx = \int f(x)dx + \int g(x)dx$$

$$869. \int [f(x) - g(x)]dx = \int f(x)dx - \int g(x)dx$$

$$870. \int f(ax)dx = \frac{1}{a}F(ax) + C$$

CHAPTER 9. INTEGRAL CALCULUS

$$871. \int f(ax + b)dx = \frac{1}{a}F(ax + b) + C$$

$$872. \int f(x)f'(x)dx = \frac{1}{2}f^2(x) + C$$

$$873. \int \frac{f'(x)}{f(x)}dx = \ln|f(x)| + C$$

874. Method of Substitution

$$\int f(x)dx = \int f(u(t))u'(t)dt \text{ if } x = u(t).$$

875. Integration by Parts

$$\int u dv = uv - \int v du,$$

where  $u(x)$ ,  $v(x)$  are differentiable functions.

## 9.2 Integrals of Rational Functions

$$876. \int adx = ax + C$$

$$877. \int xdx = \frac{x^2}{2} + C$$

$$878. \int x^2dx = \frac{x^3}{3} + C$$

$$879. \int x^p dx = \frac{x^{p+1}}{p+1} + C, p \neq -1.$$

## CHAPTER 9. INTEGRAL CALCULUS

$$880. \int (ax + b)^n dx = \frac{(ax + b)^{n+1}}{a(n+1)} + C, n \neq -1.$$

$$881. \int \frac{dx}{x} = \ln|x| + C$$

$$882. \int \frac{dx}{ax + b} = \frac{1}{a} \ln|ax + b| + C$$

$$883. \int \frac{ax + b}{cx + d} dx = \frac{a}{c}x + \frac{bc - ad}{c^2} \ln|cx + d| + C$$

$$884. \int \frac{dx}{(x+a)(x+b)} = \frac{1}{a-b} \ln \left| \frac{x+b}{x+a} \right| + C, a \neq b.$$

$$885. \int \frac{xdx}{a + bx} = \frac{1}{b^2} (a + bx - a \ln|a + bx|) + C$$

$$886. \int \frac{x^2 dx}{a + bx} = \frac{1}{b^3} \left[ \frac{1}{2} (a + bx)^2 - 2a(a + bx) + a^2 \ln|a + bx| \right] + C$$

$$887. \int \frac{dx}{x(a + bx)} = \frac{1}{a} \ln \left| \frac{a + bx}{x} \right| + C$$

$$888. \int \frac{dx}{x^2(a + bx)} = -\frac{1}{ax} + \frac{b}{a^2} \ln \left| \frac{a + bx}{x} \right| + C$$

$$889. \int \frac{xdx}{(a + bx)^2} = \frac{1}{b^2} \left( \ln|a + bx| + \frac{a}{a + bx} \right) + C$$

## CHAPTER 9. INTEGRAL CALCULUS

$$890. \int \frac{x^2 dx}{(a+bx)^2} = \frac{1}{b^3} \left( a+bx - 2a \ln|a+bx| - \frac{a^2}{a+bx} \right) + C$$

$$891. \int \frac{dx}{x(a+bx)^2} = \frac{1}{a(a+bx)} + \frac{1}{a^2} \ln \left| \frac{a+bx}{x} \right| + C$$

$$892. \int \frac{dx}{x^2-1} = \frac{1}{2} \ln \left| \frac{x-1}{x+1} \right| + C$$

$$893. \int \frac{dx}{1-x^2} = \frac{1}{2} \ln \left| \frac{1+x}{1-x} \right| + C$$

$$894. \int \frac{dx}{a^2-x^2} = \frac{1}{2a} \ln \left| \frac{a+x}{a-x} \right| + C$$

$$895. \int \frac{dx}{x^2-a^2} = \frac{1}{2a} \ln \left| \frac{x-a}{x+a} \right| + C$$

$$896. \int \frac{dx}{1+x^2} = \tan^{-1} x + C$$

$$897. \int \frac{dx}{a^2+x^2} = \frac{1}{a} \tan^{-1} \frac{x}{a} + C$$

$$898. \int \frac{x dx}{x^2+a^2} = \frac{1}{2} \ln(x^2+a^2) + C$$

$$899. \int \frac{dx}{a+bx^2} = \frac{1}{\sqrt{ab}} \arctan \left( x \sqrt{\frac{b}{a}} \right) + C, \quad ab > 0.$$



## CHAPTER 9. INTEGRAL CALCULUS

$$900. \int \frac{x dx}{a + bx^2} = \frac{1}{2b} \ln \left| x^2 + \frac{a}{b} \right| + C$$

$$901. \int \frac{dx}{x(a + bx^2)} = \frac{1}{2a} \ln \left| \frac{x^2}{a + bx^2} \right| + C$$

$$902. \int \frac{dx}{a^2 - b^2 x^2} = \frac{1}{2ab} \ln \left| \frac{a + bx}{a - bx} \right| + C$$

$$903. \int \frac{dx}{ax^2 + bx + c} = \frac{1}{\sqrt{b^2 - 4ac}} \ln \left| \frac{2ax + b - \sqrt{b^2 - 4ac}}{2ax + b + \sqrt{b^2 - 4ac}} \right| + C,$$

$$b^2 - 4ac > 0.$$

$$904. \int \frac{dx}{ax^2 + bx + c} = \frac{2}{\sqrt{4ac - b^2}} \arctan \frac{2ax + b}{\sqrt{4ac - b^2}} + C,$$

$$b^2 - 4ac < 0.$$

## 9.3 Integrals of Irrational Functions

$$905. \int \frac{dx}{\sqrt{ax + b}} = \frac{2}{a} \sqrt{ax + b} + C$$

$$906. \int \sqrt{ax + b} dx = \frac{2}{3a} (ax + b)^{3/2} + C$$

$$907. \int \frac{x dx}{\sqrt{ax + b}} = \frac{2(ax - 2b)}{3a^2} \sqrt{ax + b} + C$$

## CHAPTER 9. INTEGRAL CALCULUS

$$908. \int x\sqrt{ax+b} \, dx = \frac{2(3ax-2b)}{15a^2}(ax+b)^{3/2} + C$$

$$909. \int \frac{dx}{(x+c)\sqrt{ax+b}} = \frac{1}{\sqrt{b-ac}} \ln \left| \frac{\sqrt{ax+b} - \sqrt{b-ac}}{\sqrt{ax+b} + \sqrt{b-ac}} \right| + C,$$

$b-ac > 0.$

$$910. \int \frac{dx}{(x+c)\sqrt{ax+b}} = \frac{1}{\sqrt{ac-b}} \arctan \sqrt{\frac{ax+b}{ac-b}} + C,$$

$b-ac < 0.$

$$911. \int \sqrt{\frac{ax+b}{cx+d}} \, dx = \frac{1}{c} \sqrt{(ax+b)(cx+d)} -$$

$$- \frac{ad-bc}{c\sqrt{ac}} \ln \left| \sqrt{a(cx+d)} + \sqrt{c(ax+b)} \right| + C, a > 0.$$

$$912. \int \sqrt{\frac{ax+b}{cx+d}} \, dx = \frac{1}{c} \sqrt{(ax+b)(cx+d)} -$$

$$- \frac{ad-bc}{c\sqrt{ac}} \arctan \sqrt{\frac{a(cx+d)}{c(ax+b)}} + C, (a < 0, c > 0).$$

$$913. \int x^2 \sqrt{a+bx} \, dx = \frac{2(8a^2 - 12abx + 15b^2x^2)}{105b^3} \sqrt{(a+bx)^3} + C$$

$$914. \int \frac{x^2 dx}{\sqrt{a+bx}} = \frac{2(8a^2 - 4abx + 3b^2x^2)}{15b^3} \sqrt{a+bx} + C$$

$$915. \int \frac{dx}{x\sqrt{a+bx}} = \frac{1}{\sqrt{a}} \ln \left| \frac{\sqrt{a+bx} - \sqrt{a}}{\sqrt{a+bx} + \sqrt{a}} \right| + C, a > 0.$$

## CHAPTER 9. INTEGRAL CALCULUS

$$916. \int \frac{dx}{x\sqrt{a+bx}} = \frac{2}{\sqrt{-a}} \arctan \left| \frac{a+bx}{-a} \right| + C, \quad a < 0.$$

$$917. \int \sqrt{\frac{a-x}{b+x}} dx = \sqrt{(a-x)(b+x)} + (a+b) \arcsin \sqrt{\frac{x+b}{a+b}} + C$$

$$918. \int \sqrt{\frac{a+x}{b-x}} dx = -\sqrt{(a+x)(b-x)} - (a+b) \arcsin \sqrt{\frac{b-x}{a+b}} + C$$

$$919. \int \sqrt{\frac{1+x}{1-x}} dx = -\sqrt{1-x^2} + \arcsin x + C$$

$$920. \int \frac{dx}{\sqrt{(x-a)(b-a)}} = 2 \arcsin \sqrt{\frac{x-a}{b-a}} + C$$

$$921. \int \sqrt{a+bx-cx^2} dx = \frac{2cx-b}{4c} \sqrt{a+bx-cx^2} + \\ + \frac{b^2-4ac}{8\sqrt{c^3}} \arcsin \frac{2cx-b}{\sqrt{b^2+4ac}} + C$$

$$922. \int \frac{dx}{\sqrt{ax^2+bx+c}} = \frac{1}{\sqrt{a}} \ln \left| 2ax+b+2\sqrt{a(ax^2+bx+c)} \right| + C, \\ a > 0.$$

$$923. \int \frac{dx}{\sqrt{ax^2+bx+c}} = -\frac{1}{\sqrt{a}} \arcsin \frac{2ax+b}{4a} \sqrt{b^2-4ac} + C, \quad a < 0.$$

$$924. \int \sqrt{x^2+a^2} dx = \frac{x}{2} \sqrt{x^2+a^2} + \frac{a^2}{2} \ln \left| x + \sqrt{x^2+a^2} \right| + C$$

## CHAPTER 9. INTEGRAL CALCULUS

$$925. \int x\sqrt{x^2+a^2} dx = \frac{1}{3}(x^2+a^2)^{3/2} + C$$

$$926. \int x^2\sqrt{x^2+a^2} dx = \frac{x}{8}(2x^2+a^2)\sqrt{x^2+a^2} - \frac{a^4}{8}\ln|x+\sqrt{x^2+a^2}| + C$$

$$927. \int \frac{\sqrt{x^2+a^2}}{x^2} dx = -\frac{\sqrt{x^2+a^2}}{x} + \ln|x+\sqrt{x^2+a^2}| + C$$

$$928. \int \frac{dx}{\sqrt{x^2+a^2}} = \ln|x+\sqrt{x^2+a^2}| + C$$

$$929. \int \frac{\sqrt{x^2+a^2}}{x} dx = \sqrt{x^2+a^2} + a \ln\left|\frac{x}{a+\sqrt{x^2+a^2}}\right| + C$$

$$930. \int \frac{xdx}{\sqrt{x^2+a^2}} = \sqrt{x^2+a^2} + C$$

$$931. \int \frac{x^2 dx}{\sqrt{x^2+a^2}} = \frac{x}{2}\sqrt{x^2+a^2} - \frac{a^2}{2}\ln|x+\sqrt{x^2+a^2}| + C$$

$$932. \int \frac{dx}{x\sqrt{x^2+a^2}} = \frac{1}{a}\ln\left|\frac{x}{a+\sqrt{x^2+a^2}}\right| + C$$

$$933. \int \sqrt{x^2-a^2} dx = \frac{x}{2}\sqrt{x^2-a^2} - \frac{a^2}{2}\ln|x+\sqrt{x^2-a^2}| + C$$

$$934. \int x\sqrt{x^2-a^2} dx = \frac{1}{3}(x^2-a^2)^{3/2} + C$$

## CHAPTER 9. INTEGRAL CALCULUS

$$935. \int \frac{\sqrt{x^2 - a^2}}{x} dx = \sqrt{x^2 - a^2} + a \arcsin \frac{a}{x} + C$$

$$936. \int \frac{\sqrt{x^2 - a^2}}{x^2} dx = -\frac{\sqrt{x^2 - a^2}}{x} + \ln|x + \sqrt{x^2 - a^2}| + C$$

$$937. \int \frac{dx}{\sqrt{x^2 - a^2}} = \ln|x + \sqrt{x^2 - a^2}| + C$$

$$938. \int \frac{x dx}{\sqrt{x^2 - a^2}} = \sqrt{x^2 - a^2} + C$$

$$939. \int \frac{x^2 dx}{\sqrt{x^2 - a^2}} = \frac{x}{2} \sqrt{x^2 - a^2} + \frac{a^2}{2} \ln|x + \sqrt{x^2 - a^2}| + C$$

$$940. \int \frac{dx}{x\sqrt{x^2 - a^2}} = -\frac{1}{a} \arcsin \frac{a}{x} + C$$

$$941. \int \frac{dx}{(x+a)\sqrt{x^2 - a^2}} = \frac{1}{a} \sqrt{\frac{x-a}{x+a}} + C$$

$$942. \int \frac{dx}{(x-a)\sqrt{x^2 - a^2}} = -\frac{1}{a} \sqrt{\frac{x+a}{x-a}} + C$$

$$943. \int \frac{dx}{x^2 \sqrt{x^2 - a^2}} = \frac{\sqrt{x^2 - a^2}}{a^2 x} + C$$

$$944. \int \frac{dx}{(x^2 - a^2)^{3/2}} = -\frac{x}{a^2 \sqrt{x^2 - a^2}} + C$$

## CHAPTER 9. INTEGRAL CALCULUS

$$945. \int (x^2 - a^2)^{3/2} dx = -\frac{x}{8}(2x^2 - 5a^2)\sqrt{x^2 - a^2} + \frac{3a^4}{8} \ln|x + \sqrt{x^2 - a^2}| + C$$

$$946. \int \sqrt{a^2 - x^2} dx = \frac{x}{2}\sqrt{a^2 - x^2} + \frac{a^2}{2} \arcsin \frac{x}{a} + C$$

$$947. \int x\sqrt{a^2 - x^2} dx = -\frac{1}{3}(a^2 - x^2)^{3/2} + C$$

$$948. \int x^2\sqrt{a^2 - x^2} dx = \frac{x}{8}(2x^2 - a^2)\sqrt{a^2 - x^2} + \frac{a^4}{8} \arcsin \frac{x}{a} + C$$

$$949. \int \frac{\sqrt{a^2 - x^2}}{x} dx = \sqrt{a^2 - x^2} + a \ln \left| \frac{x}{a + \sqrt{a^2 - x^2}} \right| + C$$

$$950. \int \frac{\sqrt{a^2 - x^2}}{x^2} dx = -\frac{\sqrt{a^2 - x^2}}{x} - \arcsin \frac{x}{a} + C$$

$$951. \int \frac{dx}{\sqrt{1 - x^2}} = \arcsin x + C$$

$$952. \int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1} \frac{x}{a} + C$$

$$953. \int \frac{xdx}{\sqrt{a^2 - x^2}} = -\sqrt{a^2 - x^2} + C$$

$$954. \int \frac{x^2 dx}{\sqrt{a^2 - x^2}} = -\frac{x}{2}\sqrt{a^2 - x^2} + \frac{a^2}{2} \arcsin \frac{x}{a} + C$$

## CHAPTER 9. INTEGRAL CALCULUS

$$955. \int \frac{dx}{(x+a)\sqrt{a^2-x^2}} = -\frac{1}{2} \sqrt{\frac{a-x}{a+x}} + C$$

$$956. \int \frac{dx}{(x-a)\sqrt{a^2-x^2}} = -\frac{1}{2} \sqrt{\frac{a+x}{a-x}} + C$$

$$957. \int \frac{dx}{(x+b)\sqrt{a^2-x^2}} = \frac{1}{\sqrt{b^2-a^2}} \arcsin \frac{bx+a^2}{a(x+b)} + C, \quad b > a.$$

$$958. \int \frac{dx}{(x+b)\sqrt{a^2-x^2}} = \frac{1}{\sqrt{a^2-b^2}} \ln \left| \frac{x+b}{\sqrt{a^2-b^2}\sqrt{a^2-x^2} + a^2+bx} \right| + C,$$

$b < a.$

$$959. \int \frac{dx}{x^2\sqrt{a^2-x^2}} = -\frac{\sqrt{a^2-x^2}}{a^2x} + C$$

$$960. \int (a^2-x^2)^{3/2} dx = \frac{x}{8}(5a^2-2x^2)\sqrt{a^2-x^2} + \frac{3a^4}{8} \arcsin \frac{x}{a} + C$$

$$961. \int \frac{dx}{(a^2-x^2)^{3/2}} = \frac{x}{a^2\sqrt{a^2-x^2}} + C$$

## 9.4 Integrals of Trigonometric Functions

$$962. \int \sin x dx = -\cos x + C$$

$$963. \int \cos x dx = \sin x + C$$

## CHAPTER 9. INTEGRAL CALCULUS

$$964. \int \sin^2 x \, dx = \frac{x}{2} - \frac{1}{4} \sin 2x + C$$

$$965. \int \cos^2 x \, dx = \frac{x}{2} + \frac{1}{4} \sin 2x + C$$

$$966. \int \sin^3 x \, dx = \frac{1}{3} \cos^3 x - \cos x + C = \frac{1}{12} \cos 3x - \frac{3}{4} \cos x + C$$

$$967. \int \cos^3 x \, dx = \sin x - \frac{1}{3} \sin^3 x + C = \frac{1}{12} \sin 3x + \frac{3}{4} \sin x + C$$

$$968. \int \frac{dx}{\sin x} = \int \csc x \, dx = \ln \left| \tan \frac{x}{2} \right| + C$$

$$969. \int \frac{dx}{\cos x} = \int \sec x \, dx = \ln \left| \tan \left( \frac{x}{2} + \frac{\pi}{4} \right) \right| + C$$

$$970. \int \frac{dx}{\sin^2 x} = \int \csc^2 x \, dx = -\cot x + C$$

$$971. \int \frac{dx}{\cos^2 x} = \int \sec^2 x \, dx = \tan x + C$$

$$972. \int \frac{dx}{\sin^3 x} = \int \csc^3 x \, dx = -\frac{\cos x}{2 \sin^2 x} + \frac{1}{2} \ln \left| \tan \frac{x}{2} \right| + C$$

$$973. \int \frac{dx}{\cos^3 x} = \int \sec^3 x \, dx = \frac{\sin x}{2 \cos^2 x} + \frac{1}{2} \ln \left| \tan \left( \frac{x}{2} + \frac{\pi}{4} \right) \right| + C$$

$$974. \int \sin x \cos x \, dx = -\frac{1}{4} \cos 2x + C$$



CHAPTER 9. INTEGRAL CALCULUS

$$975. \int \sin^2 x \cos x \, dx = \frac{1}{3} \sin^3 x + C$$

$$976. \int \sin x \cos^2 x \, dx = -\frac{1}{3} \cos^3 x + C$$

$$977. \int \sin^2 x \cos^2 x \, dx = \frac{x}{8} - \frac{1}{32} \sin 4x + C$$

$$978. \int \tan x \, dx = -\ln|\cos x| + C$$

$$979. \int \frac{\sin x}{\cos^2 x} \, dx = \frac{1}{\cos x} + C = \sec x + C$$

$$980. \int \frac{\sin^2 x}{\cos x} \, dx = \ln \left| \tan \left( \frac{x}{2} + \frac{\pi}{4} \right) \right| - \sin x + C$$

$$981. \int \tan^2 x \, dx = \tan x - x + C$$

$$982. \int \cot x \, dx = \ln|\sin x| + C$$

$$983. \int \frac{\cos x}{\sin^2 x} \, dx = -\frac{1}{\sin x} + C = -\csc x + C$$

$$984. \int \frac{\cos^2 x}{\sin x} \, dx = \ln \left| \tan \frac{x}{2} \right| + \cos x + C$$

$$985. \int \cot^2 x \, dx = -\cot x - x + C$$

$$986. \int \frac{dx}{\cos x \sin x} = \ln|\tan x| + C$$

## CHAPTER 9. INTEGRAL CALCULUS

$$987. \int \frac{dx}{\sin^2 x \cos x} = -\frac{1}{\sin x} + \ln \left| \tan \left( \frac{x}{2} + \frac{\pi}{4} \right) \right| + C$$

$$988. \int \frac{dx}{\sin x \cos^2 x} = \frac{1}{\cos x} + \ln \left| \tan \frac{x}{2} \right| + C$$

$$989. \int \frac{dx}{\sin^2 x \cos^2 x} = \tan x - \cot x + C$$

$$990. \int \sin mx \sin nx \, dx = -\frac{\sin(m+n)x}{2(m+n)} + \frac{\sin(m-n)x}{2(m-n)} + C, \\ m^2 \neq n^2.$$

$$991. \int \sin mx \cos nx \, dx = -\frac{\cos(m+n)x}{2(m+n)} - \frac{\cos(m-n)x}{2(m-n)} + C, \\ m^2 \neq n^2.$$

$$992. \int \cos mx \cos nx \, dx = \frac{\sin(m+n)x}{2(m+n)} + \frac{\sin(m-n)x}{2(m-n)} + C, \\ m^2 \neq n^2.$$

$$993. \int \sec x \tan x \, dx = \sec x + C$$

$$994. \int \csc x \cot x \, dx = -\csc x + C$$

$$995. \int \sin x \cos^n x \, dx = -\frac{\cos^{n+1} x}{n+1} + C$$

$$996. \int \sin^n x \cos x \, dx = \frac{\sin^{n+1} x}{n+1} + C$$

CHAPTER 9. INTEGRAL CALCULUS

$$997. \int \arcsin x \, dx = x \arcsin x + \sqrt{1-x^2} + C$$

$$998. \int \arccos x \, dx = x \arccos x - \sqrt{1-x^2} + C$$

$$999. \int \arctan x \, dx = x \arctan x - \frac{1}{2} \ln(x^2 + 1) + C$$

$$1000. \int \operatorname{arc cot} x \, dx = x \operatorname{arc cot} x + \frac{1}{2} \ln(x^2 + 1) + C$$

## 9.5 Integrals of Hyperbolic Functions

$$1001. \int \sinh x \, dx = \cosh x + C$$

$$1002. \int \cosh x \, dx = \sinh x + C$$

$$1003. \int \tanh x \, dx = \ln \cosh x + C$$

$$1004. \int \operatorname{coth} x \, dx = \ln |\sinh x| + C$$

$$1005. \int \operatorname{sech}^2 x \, dx = \tanh x + C$$

$$1006. \int \operatorname{csch}^2 x \, dx = -\operatorname{coth} x + C$$

$$1007. \int \operatorname{sech} x \tanh x \, dx = -\operatorname{sech} x + C$$

CHAPTER 9. INTEGRAL CALCULUS

$$1008. \int \operatorname{csch} x \operatorname{coth} x dx = -\operatorname{csch} x + C$$

## 9.6 Integrals of Exponential and Logarithmic Functions

$$1009. \int e^x dx = e^x + C$$

$$1010. \int a^x dx = \frac{a^x}{\ln a} + C$$

$$1011. \int e^{ax} dx = \frac{e^{ax}}{a} + C$$

$$1012. \int x e^{ax} dx = \frac{e^{ax}}{a^2} (ax - 1) + C$$

$$1013. \int \ln x dx = x \ln x - x + C$$

$$1014. \int \frac{dx}{x \ln x} = \ln |\ln x| + C$$

$$1015. \int x^n \ln x dx = x^{n+1} \left[ \frac{\ln x}{n+1} - \frac{1}{(n+1)^2} \right] + C$$

$$1016. \int e^{ax} \sin bx dx = \frac{a \sin bx - b \cos bx}{a^2 + b^2} e^{ax} + C$$

## CHAPTER 9. INTEGRAL CALCULUS

$$1017. \int e^{ax} \cos bx \, dx = \frac{a \cos bx + b \sin bx}{a^2 + b^2} e^{ax} + C$$

## 9.7 Reduction Formulas

$$1018. \int x^n e^{mx} \, dx = \frac{1}{m} x^n e^{mx} - \frac{n}{m} \int x^{n-1} e^{mx} \, dx$$

$$1019. \int \frac{e^{mx}}{x^n} \, dx = -\frac{e^{mx}}{(n-1)x^{n-1}} + \frac{m}{n-1} \int \frac{e^{mx}}{x^{n-1}} \, dx, \quad n \neq 1.$$

$$1020. \int \sinh^n x \, dx = \frac{1}{n} \sinh^{n-1} x \cosh x - \frac{n-1}{n} \int \sinh^{n-2} x \, dx$$

$$1021. \int \frac{dx}{\sinh^n x} = -\frac{\cosh x}{(n-1)\sinh^{n-1} x} - \frac{n-2}{n-1} \int \frac{dx}{\sinh^{n-2} x}, \quad n \neq 1.$$

$$1022. \int \cosh^n x \, dx = \frac{1}{n} \sinh x \cosh^{n-1} x \cosh x + \frac{n-1}{n} \int \cosh^{n-2} x \, dx$$

$$1023. \int \frac{dx}{\cosh^n x} = -\frac{\sinh x}{(n-1)\cosh^{n-1} x} + \frac{n-2}{n-1} \int \frac{dx}{\cosh^{n-2} x}, \quad n \neq 1.$$

$$1024. \int \sinh^n x \cosh^m x \, dx = \frac{\sinh^{n+1} x \cosh^{m-1} x}{n+m} \\ + \frac{m-1}{n+m} \int \sinh^n x \cosh^{m-2} x \, dx$$

$$1025. \int \sinh^n x \cosh^m x \, dx = \frac{\sinh^{n-1} x \cosh^{m+1} x}{n+m}$$

## CHAPTER 9. INTEGRAL CALCULUS

$$-\frac{n-1}{n+m} \int \sinh^{n-2} x \cosh^m x dx$$

$$1026. \int \tanh^n x dx = -\frac{1}{n-1} \tanh^{n-1} x + \int \tanh^{n-2} x dx, n \neq 1.$$

$$1027. \int \coth^n x dx = -\frac{1}{n-1} \coth^{n-1} x + \int \coth^{n-2} x dx, n \neq 1.$$

$$1028. \int \operatorname{sech}^n x dx = \frac{\operatorname{sech}^{n-2} x \tanh x}{n-1} + \frac{n-2}{n-1} \int \operatorname{sech}^{n-2} x dx, n \neq 1.$$

$$1029. \int \sin^n x dx = -\frac{1}{n} \sin^{n-1} x \cos x + \frac{n-1}{n} \int \sin^{n-2} x dx$$

$$1030. \int \frac{dx}{\sin^n x} = -\frac{\cos x}{(n-1)\sin^{n-1} x} + \frac{n-2}{n-1} \int \frac{dx}{\sin^{n-2} x}, n \neq 1.$$

$$1031. \int \cos^n x dx = \frac{1}{n} \sin x \cos^{n-1} x + \frac{n-1}{n} \int \cos^{n-2} x dx$$

$$1032. \int \frac{dx}{\cos^n x} = \frac{\sin x}{(n-1)\cos^{n-1} x} + \frac{n-2}{n-1} \int \frac{dx}{\cos^{n-2} x}, n \neq 1.$$

$$1033. \int \sin^n x \cos^m x dx = \frac{\sin^{n+1} x \cos^{m-1} x}{n+m} \\ + \frac{m-1}{n+m} \int \sin^n x \cos^{m-2} x dx$$

$$1034. \int \sin^n x \cos^m x dx = -\frac{\sin^{n-1} x \cos^{m+1} x}{n+m}$$

## CHAPTER 9. INTEGRAL CALCULUS

$$+ \frac{n-1}{n+m} \int \sin^{n-2} x \cos^m x dx$$

$$1035. \int \tan^n x dx = \frac{1}{n-1} \tan^{n-1} x - \int \tan^{n-2} x dx, n \neq 1.$$

$$1036. \int \cot^n x dx = -\frac{1}{n-1} \cot^{n-1} x - \int \cot^{n-2} x dx, n \neq 1.$$

$$1037. \int \sec^n x dx = \frac{\sec^{n-2} x \tan x}{n-1} + \frac{n-2}{n-1} \int \sec^{n-2} x dx, n \neq 1.$$

$$1038. \int \csc^n x dx = -\frac{\csc^{n-2} x \cot x}{n-1} + \frac{n-2}{n-1} \int \csc^{n-2} x dx, n \neq 1.$$

$$1039. \int x^n \ln^m x dx = \frac{x^{n+1} \ln^m x}{n+1} - \frac{m}{n+1} \int x^n \ln^{m-1} x dx$$

$$1040. \int \frac{\ln^m x}{x^n} dx = -\frac{\ln^m x}{(n-1)x^{n-1}} + \frac{m}{n-1} \int \frac{\ln^{m-1} x}{x^n} dx, n \neq 1.$$

$$1041. \int \ln^n x dx = x \ln^n x - n \int \ln^{n-1} x dx$$

$$1042. \int x^n \sinh x dx = x^n \cosh x - n \int x^{n-1} \cosh x dx$$

$$1043. \int x^n \cosh x dx = x^n \sinh x - n \int x^{n-1} \sinh x dx$$

$$1044. \int x^n \sin x dx = -x^n \cos x + n \int x^{n-1} \cos x dx$$

$$1045. \int x^n \cos x dx = x^n \sin x - n \int x^{n-1} \sin x dx$$

## CHAPTER 9. INTEGRAL CALCULUS

$$1046. \int x^n \sin^{-1} x dx = \frac{x^{n+1}}{n+1} \sin^{-1} x - \frac{1}{n+1} \int \frac{x^{n+1}}{\sqrt{1-x^2}} dx$$

$$1047. \int x^n \cos^{-1} x dx = \frac{x^{n+1}}{n+1} \cos^{-1} x + \frac{1}{n+1} \int \frac{x^{n+1}}{\sqrt{1-x^2}} dx$$

$$1048. \int x^n \tan^{-1} x dx = \frac{x^{n+1}}{n+1} \tan^{-1} x - \frac{1}{n+1} \int \frac{x^{n+1}}{1+x^2} dx$$

$$1049. \int \frac{x^n dx}{ax^n + b} = \frac{x}{a} - \frac{b}{a} \int \frac{dx}{ax^n + b}$$

$$1050. \int \frac{dx}{(ax^2 + bx + c)^n} = \frac{-2ax - b}{(n-1)(b^2 - 4ac)(ax^2 + bx + c)^{n-1}} - \frac{2(2n-3)a}{(n-1)(b^2 - 4ac)} \int \frac{dx}{(ax^2 + bx + c)^{n-1}}, \quad n \neq 1.$$

$$1051. \int \frac{dx}{(x^2 + a^2)^n} = \frac{x}{2(n-1)a^2(x^2 + a^2)^{n-1}} + \frac{2n-3}{2(n-1)a^2} \int \frac{dx}{(x^2 + a^2)^{n-1}}, \quad n \neq 1.$$

$$1052. \int \frac{dx}{(x^2 - a^2)^n} = -\frac{x}{2(n-1)a^2(x^2 - a^2)^{n-1}} - \frac{2n-3}{2(n-1)a^2} \int \frac{dx}{(x^2 - a^2)^{n-1}}, \quad n \neq 1.$$



CHAPTER 9. INTEGRAL CALCULUS

## 9.8 Definite Integral

Definite integral of a function:  $\int_a^b f(x)dx, \int_a^b g(x)dx, \dots$

Riemann sum:  $\sum_{i=1}^n f(\xi_i)\Delta x_i$

Small changes:  $\Delta x_i$

Antiderivatives:  $F(x), G(x)$

Limits of integrations:  $a, b, c, d$

$$1053. \int_a^b f(x)dx = \lim_{\substack{n \rightarrow \infty \\ \max \Delta x_i \rightarrow 0}} \sum_{i=1}^n f(\xi_i)\Delta x_i,$$

where  $\Delta x_i = x_i - x_{i-1}, x_{i-1} \leq \xi_i \leq x_i$ .




Figure 179.

CHAPTER 9. INTEGRAL CALCULUS

$$1054. \int_a^b 1 dx = b - a$$

$$1055. \int_a^b kf(x) dx = k \int_a^b f(x) dx$$

$$1056. \int_a^b [f(x) + g(x)] dx = \int_a^b f(x) dx + \int_a^b g(x) dx$$

$$1057. \int_a^b [f(x) - g(x)] dx = \int_a^b f(x) dx - \int_a^b g(x) dx$$

$$1058. \int_a^a f(x) dx = 0$$

$$1059. \int_a^b f(x) dx = - \int_b^a f(x) dx$$

$$1060. \int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx \text{ for } a < c < b.$$

$$1061. \int_a^b f(x) dx \geq 0 \text{ if } f(x) \geq 0 \text{ on } [a, b].$$

$$1062. \int_a^b f(x) dx \leq 0 \text{ if } f(x) \leq 0 \text{ on } [a, b].$$

CHAPTER 9. INTEGRAL CALCULUS

**1063.** Fundamental Theorem of Calculus

$$\int_a^b f(x)dx = F(x)\Big|_a^b = F(b) - F(a) \text{ if } F'(x) = f(x).$$

**1064.** Method of Substitution

If  $x = g(t)$ , then

$$\int_a^b f(x)dx = \int_c^d f(g(t))g'(t)dt,$$

where

$$c = g^{-1}(a), d = g^{-1}(b).$$

**1065.** Integration by Parts

$$\int_a^b u dv = (uv)\Big|_a^b - \int_a^b v du$$

**1066.** Trapezoidal Rule

$$\int_a^b f(x)dx = \frac{b-a}{2n} \left[ f(x_0) + f(x_n) + 2 \sum_{i=1}^{n-1} f(x_i) \right]$$

CHAPTER 9. INTEGRAL CALCULUS




Figure 180.

**1067.** Simpson's Rule

$$\int_a^b f(x)dx = \frac{b-a}{3n} [f(x_0) + 4f(x_1) + 2f(x_2) + 4f(x_3) + 2f(x_4) + \dots + 4f(x_{n-1}) + f(x_n)],$$

where

$$x_i = a + \frac{b-a}{n}i, \quad i = 0, 1, 2, \dots, n.$$

CHAPTER 9. INTEGRAL CALCULUS




Figure 181.

**1068.** Area Under a Curve

$$S = \int_a^b f(x) dx = F(b) - F(a),$$

where  $F'(x) = f(x)$ .

CHAPTER 9. INTEGRAL CALCULUS




Figure 182.

**1069.** Area Between Two Curves

$$S = \int_a^b [f(x) - g(x)] dx = F(b) - G(b) - F(a) + G(a),$$

where  $F'(x) = f(x)$ ,  $G'(x) = g(x)$ .

CHAPTER 9. INTEGRAL CALCULUS




Figure 183.

## 9.9 Improper Integral

**1070.** The definite integral  $\int_a^b f(x)dx$  is called an **improper integral**

if

- a or b is infinite,
- $f(x)$  has one or more points of discontinuity in the interval  $[a, b]$ .

**1071.** If  $f(x)$  is a continuous function on  $[a, \infty)$ , then

$$\int_a^{\infty} f(x)dx = \lim_{n \rightarrow \infty} \int_a^n f(x)dx.$$

CHAPTER 9. INTEGRAL CALCULUS




Figure 184.

**1072.** If  $f(x)$  is a continuous function on  $(-\infty, b]$ , then

$$\int_{-\infty}^b f(x) dx = \lim_{n \rightarrow -\infty} \int_n^b f(x) dx.$$




Figure 185.



CHAPTER 9. INTEGRAL CALCULUS

Note : The improper integrals in 1071, 1072 are **convergent** if the limits exist and are finite; otherwise the integrals are **divergent**.

$$1073. \int_{-\infty}^{\infty} f(x)dx = \int_{-\infty}^c f(x)dx + \int_c^{\infty} f(x)dx$$




Figure 186.

If for some real number  $c$ , both of the integrals in the right side are convergent, then the integral  $\int_{-\infty}^{\infty} f(x)dx$  is also **convergent**; otherwise it is **divergent**.

**1074. Comparison Theorems**

Let  $f(x)$  and  $g(x)$  be continuous functions on the closed interval  $[a, \infty)$ . Suppose that  $0 \leq g(x) \leq f(x)$  for all  $x$  in  $[a, \infty)$ .

CHAPTER 9. INTEGRAL CALCULUS

- If  $\int_a^{\infty} f(x)dx$  is convergent, then  $\int_a^{\infty} g(x)dx$  is also convergent,
- If  $\int_a^{\infty} g(x)dx$  is divergent, then  $\int_a^{\infty} f(x)dx$  is also divergent.

**1075.** Absolute Convergence

If  $\int_a^{\infty} |f(x)|dx$  is convergent, then the integral  $\int_a^{\infty} f(x)dx$  is absolutely convergent.

**1076.** Discontinuous Integrand

Let  $f(x)$  be a function which is continuous on the interval  $[a, b)$  but is discontinuous at  $x = b$ . Then

$$\int_a^b f(x)dx = \lim_{\varepsilon \rightarrow 0^+} \int_a^{b-\varepsilon} f(x)dx$$




Figure 187.

## CHAPTER 9. INTEGRAL CALCULUS

**1077.** Let  $f(x)$  be a continuous function for all real numbers  $x$  in the interval  $[a, b]$  except for some point  $c$  in  $(a, b)$ . Then

$$\int_a^b f(x) dx = \lim_{\varepsilon \rightarrow 0^+} \int_a^{c-\varepsilon} f(x) dx + \lim_{\delta \rightarrow 0^+} \int_{c+\delta}^b f(x) dx.$$




Figure 188.

## 9.10 Double Integral

Functions of two variables:  $f(x, y)$ ,  $f(u, v)$ , ...

Double integrals:  $\iint_R f(x, y) dx dy$ ,  $\iint_R g(x, y) dx dy$ , ...

Riemann sum:  $\sum_{i=1}^m \sum_{j=1}^n f(u_i, v_j) \Delta x_i \Delta y_j$

Small changes:  $\Delta x_i$ ,  $\Delta y_j$

Regions of integration:  $R$ ,  $S$

Polar coordinates:  $r$ ,  $\theta$

CHAPTER 9. INTEGRAL CALCULUS

- Area:  $A$
- Surface area:  $S$
- Volume of a solid:  $V$
- Mass of a lamina:  $m$
- Density:  $\rho(x, y)$
- First moments:  $M_x, M_y$
- Moments of inertia:  $I_x, I_y, I_0$
- Charge of a plate:  $Q$
- Charge density:  $\sigma(x, y)$
- Coordinates of center of mass:  $\bar{x}, \bar{y}$
- Average of a function:  $\mu$

**1078.** Definition of Double Integral

The double integral over a rectangle  $[a, b] \times [c, d]$  is defined to be

$$\iint_{[a, b] \times [c, d]} f(x, y) dA = \lim_{\substack{\max \Delta x_i \rightarrow 0 \\ \max \Delta y_j \rightarrow 0}} \sum_{i=1}^m \sum_{j=1}^n f(u_i, v_j) \Delta x_i \Delta y_j,$$

where  $(u_i, v_j)$  is some point in the rectangle

$(x_{i-1}, x_i) \times (y_{j-1}, y_j)$ , and  $\Delta x_i = x_i - x_{i-1}$ ,  $\Delta y_j = y_j - y_{j-1}$ .




Figure 189.

## CHAPTER 9. INTEGRAL CALCULUS

The double integral over a general region  $R$  is

$$\iint_R f(x, y) dA = \iint_{[a, b] \times [c, d]} g(x, y) dA,$$

where rectangle  $[a, b] \times [c, d]$  contains  $R$ ,

$g(x, y) = f(x, y)$  if  $f(x, y)$  is in  $R$  and  $g(x, y) = 0$  otherwise.




Figure 190.

$$1079. \iint_R [f(x, y) + g(x, y)] dA = \iint_R f(x, y) dA + \iint_R g(x, y) dA$$

$$1080. \iint_R [f(x, y) - g(x, y)] dA = \iint_R f(x, y) dA - \iint_R g(x, y) dA$$

$$1081. \iint_R kf(x, y) dA = k \iint_R f(x, y) dA,$$

where  $k$  is a constant.

$$1082. \text{ If } f(x, y) \leq g(x, y) \text{ on } R, \text{ then } \iint_R f(x, y) dA \leq \iint_R g(x, y) dA.$$

1083. If  $f(x, y) \geq 0$  on  $R$  and  $S \subset R$ , then

CHAPTER 9. INTEGRAL CALCULUS

$$\iint_S f(x,y)dA \leq \iint_R f(x,y)dA.$$




Figure 191.

- 1084.** If  $f(x,y) \geq 0$  on  $R$  and  $R$  and  $S$  are non-overlapping regions, then  $\iint_{R \cup S} f(x,y)dA = \iint_R f(x,y)dA + \iint_S f(x,y)dA$ . Here  $R \cup S$  is the union of the regions  $R$  and  $S$ .




Figure 192.


CHAPTER 9. INTEGRAL CALCULUS

**1085.** Iterated Integrals and Fubini's Theorem

$$\iint_R f(x, y) dA = \int_a^b \int_{p(x)}^{q(x)} f(x, y) dy dx$$

for a region of type I,

$$R = \{(x, y) \mid a \leq x \leq b, p(x) \leq y \leq q(x)\}.$$



**Figure 193.**

$$\iint_R f(x, y) dA = \int_c^d \int_{u(y)}^{v(y)} f(x, y) dx dy$$

for a region of type II,

$$R = \{(x, y) \mid u(y) \leq x \leq v(y), c \leq y \leq d\}.$$

## CHAPTER 9. INTEGRAL CALCULUS




Figure 194.

**1086.** Double Integrals over Rectangular Regions

If  $R$  is the rectangular region  $[a, b] \times [c, d]$ , then

$$\iint_R f(x, y) dx dy = \int_a^b \left( \int_c^d f(x, y) dy \right) dx = \int_c^d \left( \int_a^b f(x, y) dx \right) dy.$$

In the special case where the integrand  $f(x, y)$  can be written as  $g(x)h(y)$  we have

$$\iint_R f(x, y) dx dy = \iint_R g(x)h(y) dx dy = \left( \int_a^b g(x) dx \right) \left( \int_c^d h(y) dy \right).$$

**1087.** Change of Variables

$$\iint_R f(x, y) dx dy = \iint_S f[x(u, v), y(u, v)] \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv,$$

where  $\left| \frac{\partial(x, y)}{\partial(u, v)} \right| = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} \neq 0$  is the **jacobian** of the transformations

$(x, y) \rightarrow (u, v)$ , and  $S$  is the pullback of  $R$  which



CHAPTER 9. INTEGRAL CALCULUS

can be computed by  $x = x(u, v)$ ,  $y = y(u, v)$  into the definition of  $R$ .

**1088.** Polar Coordinates  
 $x = r \cos \theta$ ,  $y = r \sin \theta$ .




Figure 195.

**1089.** Double Integrals in Polar Coordinates

The Differential  $dx dy$  for Polar Coordinates is

$$dx dy = \left| \frac{\partial(x, y)}{\partial(r, \theta)} \right| dr d\theta = r dr d\theta.$$

Let the region  $R$  is determined as follows:  
 $0 \leq g(\theta) \leq r \leq h(\theta)$ ,  $\alpha \leq \theta \leq \beta$ , where  $\beta - \alpha \leq 2\pi$ .

Then

$$\iint_R f(x, y) dx dy = \int_{\alpha}^{\beta} \int_{g(\theta)}^{h(\theta)} f(r \cos \theta, r \sin \theta) r dr d\theta.$$

CHAPTER 9. INTEGRAL CALCULUS




Figure 196.

If the region  $R$  is the **polar rectangle** given by  $0 \leq a \leq r \leq b$ ,  $\alpha \leq \theta \leq \beta$ , where  $\beta - \alpha \leq 2\pi$ , then

$$\iint_R f(x, y) dx dy = \int_{\alpha}^{\beta} \int_a^b f(r \cos \theta, r \sin \theta) r dr d\theta.$$





Figure 197.

CHAPTER 9. INTEGRAL CALCULUS


**1090.** Area of a Region

$$A = \int_a^b \int_{g(x)}^{f(x)} dy dx \quad (\text{for a type I region}).$$



**Figure 198.**

$$A = \int_c^d \int_{p(y)}^{q(y)} dx dy \quad (\text{for a type II region}).$$




**Figure 199.**

CHAPTER 9. INTEGRAL CALCULUS

**1091.** Volume of a Solid

$$V = \iint_R f(x,y) dA.$$



**Figure 200.**

If  $R$  is a type I region bounded by  $x = a$ ,  $x = b$ ,  $y = h(x)$ ,  $y = g(x)$ , then

$$V = \iint_R f(x,y) dA = \int_a^b \int_{h(x)}^{g(x)} f(x,y) dy dx.$$

If  $R$  is a type II region bounded by  $y = c$ ,  $y = d$ ,  $x = q(y)$ ,  $x = p(y)$ , then

$$V = \iint_R f(x,y) dA = \int_c^d \int_{p(y)}^{q(y)} f(x,y) dx dy.$$

## CHAPTER 9. INTEGRAL CALCULUS

If  $f(x,y) \geq g(x,y)$  over a region  $R$ , then the volume of the solid between  $z_1 = f(x,y)$  and  $z_2 = g(x,y)$  over  $R$  is given by

$$V = \iint_R [f(x,y) - g(x,y)] dA.$$

**1092.** Area and Volume in Polar Coordinates

If  $S$  is a region in the  $xy$ -plane bounded by  $\theta = \alpha$ ,  $\theta = \beta$ ,  $r = h(\theta)$ ,  $r = g(\theta)$ ,

then

$$A = \iint_S dA = \int_{\alpha}^{\beta} \int_{h(\theta)}^{g(\theta)} r dr d\theta,$$

$$V = \iint_S f(r,\theta) r dr d\theta.$$




Figure 201.

**1093.** Surface Area

$$S = \iint_R \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2} dx dy$$

## CHAPTER 9. INTEGRAL CALCULUS

**1094.** Mass of a Lamina

$$m = \iint_R \rho(x, y) dA,$$

where the lamina occupies a region  $R$  and its density at a point  $(x, y)$  is  $\rho(x, y)$ .

**1095.** Moments

The moment of the lamina about the  $x$ -axis is given by formula

$$M_x = \iint_R y\rho(x, y) dA.$$

The moment of the lamina about the  $y$ -axis is

$$M_y = \iint_R x\rho(x, y) dA.$$

The moment of inertia about the  $x$ -axis is

$$I_x = \iint_R y^2\rho(x, y) dA.$$

The moment of inertia about the  $y$ -axis is

$$I_y = \iint_R x^2\rho(x, y) dA.$$

The polar moment of inertia is

$$I_0 = \iint_R (x^2 + y^2)\rho(x, y) dA.$$

**1096.** Center of Mass

$$\bar{x} = \frac{M_y}{m} = \frac{1}{m} \iint_R x\rho(x, y) dA = \frac{\iint_R x\rho(x, y) dA}{\iint_R \rho(x, y) dA},$$

## CHAPTER 9. INTEGRAL CALCULUS

$$\bar{y} = \frac{M_x}{m} = \frac{1}{m} \iint_R y\rho(x, y) dA = \frac{\iint_R y\rho(x, y) dA}{\iint_R \rho(x, y) dA}.$$

**1097.** Charge of a Plate

$$Q = \iint_R \sigma(x, y) dA,$$

where electrical charge is distributed over a region  $R$  and its charge density at a point  $(x, y)$  is  $\sigma(x, y)$ .

**1098.** Average of a Function

$$\mu = \frac{1}{S} \iint_R f(x, y) dA,$$

$$\text{where } S = \iint_R dA.$$

## 9.11 Triple Integral

Functions of three variables:  $f(x, y, z)$ ,  $g(x, y, z)$ , ...

Triple integrals:  $\iiint_G f(x, y, z) dV$ ,  $\iiint_G g(x, y, z) dV$ , ...

Riemann sum:  $\sum_{i=1}^m \sum_{j=1}^n \sum_{k=1}^p f(u_i, v_j, w_k) \Delta x_i \Delta y_j \Delta z_k$

Small changes:  $\Delta x_i$ ,  $\Delta y_j$ ,  $\Delta z_k$

Limits of integration:  $a$ ,  $b$ ,  $c$ ,  $d$ ,  $r$ ,  $s$

Regions of integration:  $G$ ,  $T$ ,  $S$

Cylindrical coordinates:  $r$ ,  $\theta$ ,  $z$

Spherical coordinates:  $r$ ,  $\theta$ ,  $\phi$

Volume of a solid:  $V$

## CHAPTER 9. INTEGRAL CALCULUS

Mass of a solid:  $m$

Density:  $\mu(x, y, z)$

Coordinates of center of mass:  $\bar{x}$ ,  $\bar{y}$ ,  $\bar{z}$

First moments:  $M_{xy}$ ,  $M_{yz}$ ,  $M_{xz}$

Moments of inertia:  $I_{xy}$ ,  $I_{yz}$ ,  $I_{xz}$ ,  $I_x$ ,  $I_y$ ,  $I_z$ ,  $I_0$

**1099.** Definition of Triple Integral

The triple integral over a parallelepiped  $[a, b] \times [c, d] \times [r, s]$  is defined to be

$$\iiint_{[a, b] \times [c, d] \times [r, s]} f(x, y, z) dV = \lim_{\substack{\max \Delta x_i \rightarrow 0 \\ \max \Delta y_j \rightarrow 0 \\ \max \Delta z_k \rightarrow 0}} \sum_{i=1}^m \sum_{j=1}^n \sum_{k=1}^p f(u_i, v_j, w_k) \Delta x_i \Delta y_j \Delta z_k,$$

where  $(u_i, v_j, w_k)$  is some point in the parallelepiped

$(x_{i-1}, x_i) \times (y_{j-1}, y_j) \times (z_{k-1}, z_k)$ , and  $\Delta x_i = x_i - x_{i-1}$ ,

$\Delta y_j = y_j - y_{j-1}$ ,  $\Delta z_k = z_k - z_{k-1}$ .

$$\mathbf{1100.} \quad \iiint_G [f(x, y, z) + g(x, y, z)] dV = \iiint_G f(x, y, z) dV + \iiint_G g(x, y, z) dV$$

$$\mathbf{1101.} \quad \iiint_G [f(x, y, z) - g(x, y, z)] dV = \iiint_G f(x, y, z) dV - \iiint_G g(x, y, z) dV$$

$$\mathbf{1102.} \quad \iiint_G kf(x, y, z) dV = k \iiint_G f(x, y, z) dV,$$

where  $k$  is a constant.

**1103.** If  $f(x, y, z) \geq 0$  and  $G$  and  $T$  are nonoverlapping basic regions, then

$$\iiint_{G \cup T} f(x, y, z) dV = \iiint_G f(x, y, z) dV + \iiint_T f(x, y, z) dV.$$

Here  $G \cup T$  is the union of the regions  $G$  and  $T$ .



## CHAPTER 9. INTEGRAL CALCULUS

**1104.** Evaluation of Triple Integrals by Repeated Integrals

If the solid  $G$  is the set of points  $(x, y, z)$  such that  $(x, y) \in R$ ,  $\chi_1(x, y) \leq z \leq \chi_2(x, y)$ , then

$$\iiint_G f(x, y, z) dx dy dz = \iint_R \left[ \int_{\chi_1(x, y)}^{\chi_2(x, y)} f(x, y, z) dz \right] dx dy,$$

where  $R$  is projection of  $G$  onto the  $xy$ -plane.

If the solid  $G$  is the set of points  $(x, y, z)$  such that  $a \leq x \leq b$ ,  $\varphi_1(x) \leq y \leq \varphi_2(x)$ ,  $\chi_1(x, y) \leq z \leq \chi_2(x, y)$ , then

$$\iiint_G f(x, y, z) dx dy dz = \int_a^b \left[ \int_{\varphi_1(x)}^{\varphi_2(x)} \left( \int_{\chi_1(x, y)}^{\chi_2(x, y)} f(x, y, z) dz \right) dy \right] dx$$

**1105.** Triple Integrals over Parallelepiped

If  $G$  is a parallelepiped  $[a, b] \times [c, d] \times [r, s]$ , then

$$\iiint_G f(x, y, z) dx dy dz = \int_a^b \left[ \int_c^d \left( \int_r^s f(x, y, z) dz \right) dy \right] dx.$$

In the special case where the integrand  $f(x, y, z)$  can be written as  $g(x)h(y)k(z)$  we have

$$\iiint_G f(x, y, z) dx dy dz = \left( \int_a^b g(x) dx \right) \left( \int_c^d h(y) dy \right) \left( \int_r^s k(z) dz \right).$$

**1106.** Change of Variables

$$\iiint_G f(x, y, z) dx dy dz =$$

$$= \iiint_S f[x(u, v, w), y(u, v, w), z(u, v, w)] \left| \frac{\partial(x, y, z)}{\partial(u, v, w)} \right| dx dy dz,$$

## CHAPTER 9. INTEGRAL CALCULUS

where  $\left| \frac{\partial(x, y, z)}{\partial(u, v, w)} \right| = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial w} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial w} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial w} \end{vmatrix} \neq 0$  is the **jacobian** of

the transformations  $(x, y, z) \rightarrow (u, v, w)$ , and  $S$  is the pullback of  $G$  which can be computed by  $x = x(u, v, w)$ ,  
 $y = y(u, v, w)$   
 $z = z(u, v, w)$  into the definition of  $G$ .

**1107.** Triple Integrals in Cylindrical Coordinates

The differential  $dx dy dz$  for cylindrical coordinates is

$$dx dy dz = \left| \frac{\partial(x, y, z)}{\partial(r, \theta, z)} \right| dr d\theta dz = r dr d\theta dz.$$

Let the solid  $G$  is determined as follows:

$$(x, y) \in R, \chi_1(x, y) \leq z \leq \chi_2(x, y),$$

where  $R$  is projection of  $G$  onto the  $xy$ -plane. Then

$$\begin{aligned} \iiint_G f(x, y, z) dx dy dz &= \iiint_S f(r \cos \theta, r \sin \theta, z) r dr d\theta dz \\ &= \iint_{R(r, \theta)} \left[ \int_{\chi_1(r \cos \theta, r \sin \theta)}^{\chi_2(r \cos \theta, r \sin \theta)} f(r \cos \theta, r \sin \theta, z) dz \right] r dr d\theta. \end{aligned}$$

Here  $S$  is the pullback of  $G$  in cylindrical coordinates.

**1108.** Triple Integrals in Spherical Coordinates

The Differential  $dx dy dz$  for Spherical Coordinates is

$$dx dy dz = \left| \frac{\partial(x, y, z)}{\partial(r, \theta, \varphi)} \right| dr d\theta d\varphi = r^2 \sin \theta dr d\theta d\varphi$$

$$\iiint_G f(x, y, z) dx dy dz =$$

CHAPTER 9. INTEGRAL CALCULUS

$$= \iiint_S f(r \sin \theta \cos \varphi, r \sin \theta \sin \varphi, r \cos \theta) r^2 \sin \theta dr d\theta d\varphi,$$

where the solid  $S$  is the pullback of  $G$  in spherical coordinates. The angle  $\theta$  ranges from  $0$  to  $2\pi$ , the angle  $\varphi$  ranges from  $0$  to  $\pi$ .




Figure 202.

**1109.** Volume of a Solid

$$V = \iiint_G dx dy dz$$

**1110.** Volume in Cylindrical Coordinates

$$V = \iiint_{S(r,\theta,z)} r dr d\theta dz$$

**1111.** Volume in Spherical Coordinates

$$V = \iiint_{S(r,\theta,\varphi)} r^2 \sin \theta dr d\theta d\varphi$$

## CHAPTER 9. INTEGRAL CALCULUS

**1112.** Mass of a Solid

$$m = \iiint_G \mu(x, y, z) dV,$$

where the solid occupies a region  $G$  and its density at a point  $(x, y, z)$  is  $\mu(x, y, z)$ .

**1113.** Center of Mass of a Solid

$$\bar{x} = \frac{M_{yz}}{m}, \quad \bar{y} = \frac{M_{xz}}{m}, \quad \bar{z} = \frac{M_{xy}}{m},$$

where

$$M_{yz} = \iiint_G x\mu(x, y, z) dV,$$

$$M_{xz} = \iiint_G y\mu(x, y, z) dV,$$

$$M_{xy} = \iiint_G z\mu(x, y, z) dV$$

are the first moments about the coordinate planes  $x=0$ ,  $y=0$ ,  $z=0$ , respectively,  $\mu(x, y, z)$  is the density function.

**1114.** Moments of Inertia about the  $xy$ -plane (or  $z=0$ ),  $yz$ -plane ( $x=0$ ), and  $xz$ -plane ( $y=0$ )

$$I_{xy} = \iiint_G z^2 \mu(x, y, z) dV,$$

$$I_{yz} = \iiint_G x^2 \mu(x, y, z) dV,$$

$$I_{xz} = \iiint_G y^2 \mu(x, y, z) dV.$$

**1115.** Moments of Inertia about the  $x$ -axis,  $y$ -axis, and  $z$ -axis

$$I_x = I_{xy} + I_{xz} = \iiint_G (z^2 + y^2) \mu(x, y, z) dV,$$

$$I_y = I_{xy} + I_{yz} = \iiint_G (z^2 + x^2) \mu(x, y, z) dV,$$

## CHAPTER 9. INTEGRAL CALCULUS

$$I_z = I_{xz} + I_{yz} = \iiint_G (y^2 + x^2) \mu(x, y, z) dV.$$

**1116.** Polar Moment of Inertia

$$I_0 = I_{xy} + I_{yz} + I_{xz} = \iiint_G (x^2 + y^2 + z^2) \mu(x, y, z) dV$$

## 9.12 Line Integral

Scalar functions:  $F(x, y, z)$ ,  $F(x, y)$ ,  $f(x)$

Scalar potential:  $u(x, y, z)$

Curves:  $C$ ,  $C_1$ ,  $C_2$

Limits of integrations:  $a$ ,  $b$ ,  $\alpha$ ,  $\beta$

Parameters:  $t$ ,  $s$

Polar coordinates:  $r$ ,  $\theta$

Vector field:  $\vec{F}(P, Q, R)$

Position vector:  $\vec{r}(s)$

Unit vectors:  $\vec{i}$ ,  $\vec{j}$ ,  $\vec{k}$ ,  $\vec{\tau}$

Area of region:  $S$

Length of a curve:  $L$

Mass of a wire:  $m$

Density:  $\rho(x, y, z)$ ,  $\rho(x, y)$

Coordinates of center of mass:  $\bar{x}$ ,  $\bar{y}$ ,  $\bar{z}$

First moments:  $M_{xy}$ ,  $M_{yz}$ ,  $M_{xz}$

Moments of inertia:  $I_x$ ,  $I_y$ ,  $I_z$

Volume of a solid:  $V$

Work:  $W$

Magnetic field:  $\vec{B}$

Current:  $I$

Electromotive force:  $\varepsilon$

Magnetic flux:  $\psi$

## CHAPTER 9. INTEGRAL CALCULUS

**1117.** Line Integral of a Scalar Function

Let a curve  $C$  be given by the vector function  $\vec{r} = \vec{r}(s)$ ,  
 $0 \leq s \leq S$ , and a **scalar function**  $F$  is defined over the curve  $C$ .

Then

$$\int_0^S F(\vec{r}(s)) ds = \int_C F(x, y, z) ds = \int_C F ds,$$

where  $ds$  is the arc length differential.

$$\mathbf{1118.} \quad \int_{C_1 \cup C_2} F ds = \int_{C_1} F ds + \int_{C_2} F ds$$




Figure 203.

**1119.** If the smooth curve  $C$  is parametrized by  $\vec{r} = \vec{r}(t)$ ,

$\alpha \leq t \leq \beta$ , then

$$\int_C F(x, y, z) ds = \int_{\alpha}^{\beta} F(x(t), y(t), z(t)) \sqrt{(x'(t))^2 + (y'(t))^2 + (z'(t))^2} dt.$$

**1120.** If  $C$  is a smooth curve in the  $xy$ -plane given by the equation

$y = f(x)$ ,  $a \leq x \leq b$ , then

$$\int_C F(x, y) ds = \int_a^b F(x, f(x)) \sqrt{1 + (f'(x))^2} dx.$$

**1121.** Line Integral of Scalar Function in Polar Coordinates

## CHAPTER 9. INTEGRAL CALCULUS

$$\int_C \mathbf{F}(x, y) ds = \int_{\alpha}^{\beta} \mathbf{F}(r \cos \theta, r \sin \theta) \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta,$$

where the curve  $C$  is defined by the polar function  $r(\theta)$ .

**1122.** Line Integral of Vector Field

Let a curve  $C$  be defined by the vector function  $\vec{r} = \vec{r}(s)$ ,  $0 \leq s \leq S$ . Then

$$\frac{d\vec{r}}{ds} = \vec{\tau} = (\cos \alpha, \cos \beta, \cos \gamma)$$

is the unit vector of the tangent line to this curve.




Figure 204.

Let a **vector field**  $\vec{F}(P, Q, R)$  is defined over the curve  $C$ . Then the line integral of the vector field  $\vec{F}$  along the curve  $C$  is

$$\int_C P dx + Q dy + R dz = \int_0^S (P \cos \alpha + Q \cos \beta + R \cos \gamma) ds.$$

## CHAPTER 9. INTEGRAL CALCULUS

**1123.** Properties of Line Integrals of Vector Fields

$$\int_{-C} (\vec{F} \cdot d\vec{r}) = - \int_C (\vec{F} \cdot d\vec{r}),$$

where  $-C$  denote the curve with the opposite orientation.

$$\int_C (\vec{F} \cdot d\vec{r}) = \int_{C_1 \cup C_2} (\vec{F} \cdot d\vec{r}) = \int_{C_1} (\vec{F} \cdot d\vec{r}) + \int_{C_2} (\vec{F} \cdot d\vec{r}),$$

where  $C$  is the union of the curves  $C_1$  and  $C_2$ .

**1124.** If the curve  $C$  is parameterized by  $\vec{r}(t) = \langle x(t), y(t), z(t) \rangle$ ,

$\alpha \leq t \leq \beta$ , then

$$\begin{aligned} \int_C Pdx + Qdy + Rdz &= \\ &= \int_{\alpha}^{\beta} \left( P(x(t), y(t), z(t)) \frac{dx}{dt} + Q(x(t), y(t), z(t)) \frac{dy}{dt} + R(x(t), y(t), z(t)) \frac{dz}{dt} \right) dt \end{aligned}$$

**1125.** If  $C$  lies in the  $xy$ -plane and given by the equation  $y = f(x)$ ,

then

$$\int_C Pdx + Qdy = \int_a^b \left( P(x, f(x)) + Q(x, f(x)) \frac{df}{dx} \right) dx.$$

**1126.** Green's Theorem

$$\iint_R \left( \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx dy = \oint_C Pdx + Qdy,$$

where  $\vec{F} = P(x, y)\vec{i} + Q(x, y)\vec{j}$  is a continuous vector function with continuous first partial derivatives  $\frac{\partial P}{\partial y}$ ,  $\frac{\partial Q}{\partial x}$  in a

some domain  $R$ , which is bounded by a closed, piecewise smooth curve  $C$ .



## CHAPTER 9. INTEGRAL CALCULUS

**1127.** Area of a Region R Bounded by the Curve C

$$S = \iint_R dx dy = \frac{1}{2} \oint_C x dy - y dx$$

**1128.** Path Independence of Line Integrals

The line integral of a vector function  $\vec{F} = P\vec{i} + Q\vec{j} + R\vec{k}$  is said to be **path independent**, if and only if P, Q, and R are continuous in a domain D, and if there exists some scalar function  $u = u(x, y, z)$  (a **scalar potential**) in D such that

$$\vec{F} = \text{grad } u, \text{ or } \frac{\partial u}{\partial x} = P, \frac{\partial u}{\partial y} = Q, \frac{\partial u}{\partial z} = R.$$

Then

$$\int_C \vec{F}(\vec{r}) \cdot d\vec{r} = \int_C P dx + Q dy + R dz = u(B) - u(A).$$

**1129.** Test for a Conservative Field

A vector field of the form  $\vec{F} = \text{grad } u$  is called a **conservative field**. The line integral of a vector function  $\vec{F} = P\vec{i} + Q\vec{j} + R\vec{k}$  is path independent if and only if

$$\text{curl } \vec{F} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ P & Q & R \end{vmatrix} = \vec{0}.$$

If the line integral is taken in xy-plane so that

$$\int_C P dx + Q dy = u(B) - u(A),$$

then the test for determining if a vector field is conservative can be written in the form

$$\frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}.$$

## CHAPTER 9. INTEGRAL CALCULUS

**1130.** Length of a Curve

$$L = \int_C ds = \int_{\alpha}^{\beta} \left| \frac{d\vec{r}}{dt}(t) \right| dt = \int_{\alpha}^{\beta} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2} dt,$$

where  $C$  is a piecewise smooth curve described by the position vector  $\vec{r}(t)$ ,  $\alpha \leq t \leq \beta$ .

If the curve  $C$  is two-dimensional, then

$$L = \int_C ds = \int_{\alpha}^{\beta} \left| \frac{d\vec{r}}{dt}(t) \right| dt = \int_{\alpha}^{\beta} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt.$$

If the curve  $C$  is the graph of a function  $y = f(x)$  in the  $xy$ -plane ( $a \leq x \leq b$ ), then

$$L = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx.$$

**1131.** Length of a Curve in Polar Coordinates

$$L = \int_{\alpha}^{\beta} \sqrt{\left(\frac{dr}{d\theta}\right)^2 + r^2} d\theta,$$

where the curve  $C$  is given by the equation  $r = r(\theta)$ ,  $\alpha \leq \theta \leq \beta$  in polar coordinates.

**1132.** Mass of a Wire

$$m = \int_C \rho(x, y, z) ds,$$

where  $\rho(x, y, z)$  is the mass per unit length of the wire.

If  $C$  is a curve parametrized by the vector function  $\vec{r}(t) = \langle x(t), y(t), z(t) \rangle$ , then the mass can be computed by the formula

## CHAPTER 9. INTEGRAL CALCULUS

$$m = \int_{\alpha}^{\beta} \rho(x(t), y(t), z(t)) \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2} dt.$$

If  $C$  is a curve in  $xy$ -plane, then the mass of the wire is given by

$$m = \int_C \rho(x, y) ds,$$

or

$$m = \int_{\alpha}^{\beta} \rho(x(t), y(t)) \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt \text{ (in parametric form).}$$

**1133. Center of Mass of a Wire**

$$\bar{x} = \frac{M_{yz}}{m}, \quad \bar{y} = \frac{M_{xz}}{m}, \quad \bar{z} = \frac{M_{xy}}{m},$$

where

$$M_{yz} = \int_C x\rho(x, y, z) ds,$$

$$M_{xz} = \int_C y\rho(x, y, z) ds,$$

$$M_{xy} = \int_C z\rho(x, y, z) ds.$$

**1134. Moments of Inertia**

The moments of inertia about the  $x$ -axis,  $y$ -axis, and  $z$ -axis are given by the formulas

$$I_x = \int_C (y^2 + z^2) \rho(x, y, z) ds,$$


$$I_y = \int_C (x^2 + z^2) \rho(x, y, z) ds,$$

$$I_z = \int_C (x^2 + y^2) \rho(x, y, z) ds.$$

## CHAPTER 9. INTEGRAL CALCULUS

**1135.** Area of a Region Bounded by a Closed Curve

$$S = \oint_C x dy = -\oint_C y dx = \frac{1}{2} \oint_C x dy - y dx.$$

**Figure 205.**

If the closed curve  $C$  is given in parametric form  $\vec{r}(t) = \langle x(t), y(t) \rangle$ , then the area can be calculated by the formula

$$S = \int_{\alpha}^{\beta} x(t) \frac{dy}{dt} dt = -\int_{\alpha}^{\beta} y(t) \frac{dx}{dt} dt = \frac{1}{2} \int_{\alpha}^{\beta} \left( x(t) \frac{dy}{dt} - y(t) \frac{dx}{dt} \right) dt.$$

**1136.** Volume of a Solid Formed by Rotating a Closed Curve about the  $x$ -axis

$$V = -\pi \oint_C y^2 dx = -2\pi \oint_C xy dy = -\frac{\pi}{2} \oint_C 2xy dy + y^2 dx$$

CHAPTER 9. INTEGRAL CALCULUS




Figure 206.

**1137.** Work

Work done by a force  $\vec{F}$  on an object moving along a curve  $C$  is given by the line integral

$$W = \int_C \vec{F} \cdot d\vec{r},$$

where  $\vec{F}$  is the vector force field acting on the object,  $d\vec{r}$  is the unit tangent vector.




Figure 207.

## CHAPTER 9. INTEGRAL CALCULUS

If the object is moved along a curve  $C$  in the  $xy$ -plane, then

$$W = \int_C \vec{F} \cdot d\vec{r} = \int_C Pdx + Qdy,$$

If a path  $C$  is specified by a parameter  $t$  ( $t$  often means time), the formula for calculating work becomes

$$W = \int_{\alpha}^{\beta} \left[ P(x(t), y(t), z(t)) \frac{dx}{dt} + Q(x(t), y(t), z(t)) \frac{dy}{dt} + R(x(t), y(t), z(t)) \frac{dz}{dt} \right] dt,$$

where  $t$  goes from  $\alpha$  to  $\beta$ .

If a vector field  $\vec{F}$  is conservative and  $u(x, y, z)$  is a scalar potential of the field, then the work on an object moving from  $A$  to  $B$  can be found by the formula

$$W = u(B) - u(A).$$

**1138.** Ampere's Law

$$\oint_C \vec{B} \cdot d\vec{r} = \mu_0 I.$$

The line integral of a magnetic field  $\vec{B}$  around a closed path  $C$  is equal to the total current  $I$  flowing through the area bounded by the path.




Figure 208.

## CHAPTER 9. INTEGRAL CALCULUS

**1139.** Faraday's Law

$$\varepsilon = \oint_C \vec{E} \cdot d\vec{r} = -\frac{d\psi}{dt}$$

The electromotive force (emf)  $\varepsilon$  induced around a closed loop  $C$  is equal to the rate of the change of magnetic flux  $\psi$  passing through the loop.




Figure 209.

## 9.13 Surface Integral

Scalar functions:  $f(x, y, z)$ ,  $z(x, y)$

Position vectors:  $\vec{r}(u, v)$ ,  $\vec{r}(x, y, z)$

Unit vectors:  $\vec{i}$ ,  $\vec{j}$ ,  $\vec{k}$

Surface:  $S$

Vector field:  $\vec{F}(P, Q, R)$

Divergence of a vector field:  $\text{div } \vec{F} = \nabla \cdot \vec{F}$

## CHAPTER 9. INTEGRAL CALCULUS

Curl of a vector field:  $\text{curl } \vec{F} = \nabla \times \vec{F}$ Vector element of a surface:  $d\vec{S}$ Normal to surface:  $\vec{n}$ Surface area:  $A$ Mass of a surface:  $m$ Density:  $\mu(x, y, z)$ Coordinates of center of mass:  $\bar{x}, \bar{y}, \bar{z}$ First moments:  $M_{xy}, M_{yz}, M_{xz}$ Moments of inertia:  $I_{xy}, I_{yz}, I_{xz}, I_x, I_y, I_z$ Volume of a solid:  $V$ Force:  $\vec{F}$ Gravitational constant:  $G$ Fluid velocity:  $\vec{v}(\vec{r})$ Fluid density:  $\rho$ Pressure:  $p(\vec{r})$ Mass flux, electric flux:  $\Phi$ Surface charge:  $Q$ Charge density:  $\sigma(x, y)$ Magnitude of the electric field:  $\vec{E}$ **1140.** Surface Integral of a Scalar FunctionLet a surface  $S$  be given by the position vector

$$\vec{r}(u, v) = x(u, v)\vec{i} + y(u, v)\vec{j} + z(u, v)\vec{k},$$

where  $(u, v)$  ranges over some domain  $D(u, v)$  of the  $uv$ -plane.The surface integral of a scalar function  $f(x, y, z)$  over the surface  $S$  is defined as

$$\iint_S f(x, y, z) dS = \iint_{D(u, v)} f(x(u, v), y(u, v), z(u, v)) \left| \frac{\partial \vec{r}}{\partial u} \times \frac{\partial \vec{r}}{\partial v} \right| du dv,$$

where the partial derivatives  $\frac{\partial \vec{r}}{\partial u}$  and  $\frac{\partial \vec{r}}{\partial v}$  are given by



## CHAPTER 9. INTEGRAL CALCULUS

$$\begin{aligned}\frac{\partial \vec{r}}{\partial u} &= \frac{\partial x}{\partial u}(\mathbf{i}) + \frac{\partial y}{\partial u}(\mathbf{j}) + \frac{\partial z}{\partial u}(\mathbf{k}), \\ \frac{\partial \vec{r}}{\partial v} &= \frac{\partial x}{\partial v}(\mathbf{i}) + \frac{\partial y}{\partial v}(\mathbf{j}) + \frac{\partial z}{\partial v}(\mathbf{k}) \\ \text{and } \frac{\partial \vec{r}}{\partial u} \times \frac{\partial \vec{r}}{\partial v} &\text{ is the cross product.}\end{aligned}$$

- 1141.** If the surface  $S$  is given by the equation  $z = z(x, y)$  where  $z(x, y)$  is a differentiable function in the domain  $D(x, y)$ , then

$$\iint_S f(x, y, z) dS = \iint_{D(x, y)} f(x, y, z(x, y)) \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2} dx dy.$$

- 1142.** Surface Integral of the Vector Field  $\vec{F}$  over the Surface  $S$

- If  $S$  is oriented **outward**, then

$$\begin{aligned}\iint_S \vec{F}(x, y, z) \cdot d\vec{S} &= \iint_S \vec{F}(x, y, z) \cdot \vec{n} dS \\ &= \iint_{D(u, v)} \vec{F}(x(u, v), y(u, v), z(u, v)) \cdot \left[ \frac{\partial \vec{r}}{\partial u} \times \frac{\partial \vec{r}}{\partial v} \right] du dv.\end{aligned}$$

- If  $S$  is oriented **inward**, then

$$\begin{aligned}\iint_S \vec{F}(x, y, z) \cdot d\vec{S} &= \iint_S \vec{F}(x, y, z) \cdot \vec{n} dS \\ &= \iint_{D(u, v)} \vec{F}(x(u, v), y(u, v), z(u, v)) \cdot \left[ \frac{\partial \vec{r}}{\partial v} \times \frac{\partial \vec{r}}{\partial u} \right] du dv.\end{aligned}$$

$d\vec{S} = \vec{n} dS$  is called the **vector element of the surface**. Dot means the scalar product of the appropriate vectors.

The partial derivatives  $\frac{\partial \vec{r}}{\partial u}$  and  $\frac{\partial \vec{r}}{\partial v}$  are given by

## CHAPTER 9. INTEGRAL CALCULUS

$$\frac{\partial \vec{r}}{\partial \mathbf{u}} = \frac{\partial \mathbf{x}}{\partial \mathbf{u}} \cdot \vec{i} + \frac{\partial \mathbf{y}}{\partial \mathbf{u}} \cdot \vec{j} + \frac{\partial \mathbf{z}}{\partial \mathbf{u}} \cdot \vec{k},$$

$$\frac{\partial \vec{r}}{\partial \mathbf{v}} = \frac{\partial \mathbf{x}}{\partial \mathbf{v}} \cdot \vec{i} + \frac{\partial \mathbf{y}}{\partial \mathbf{v}} \cdot \vec{j} + \frac{\partial \mathbf{z}}{\partial \mathbf{v}} \cdot \vec{k}.$$

**1143.** If the surface  $S$  is given by the equation  $z = z(x, y)$ , where  $z(x, y)$  is a differentiable function in the domain  $D(x, y)$ , then

- If  $S$  is oriented **upward**, i.e. the  $k$ -th component of the normal vector is positive, then

$$\begin{aligned} \iint_S \vec{F}(x, y, z) \cdot d\vec{S} &= \iint_S \vec{F}(x, y, z) \cdot \vec{n} dS \\ &= \iint_{D(x, y)} \vec{F}(x, y, z) \cdot \left( -\frac{\partial z}{\partial x} \vec{i} - \frac{\partial z}{\partial y} \vec{j} + \vec{k} \right) dx dy, \end{aligned}$$

- If  $S$  is oriented **downward**, i.e. the  $k$ -th component of the normal vector is negative, then

$$\begin{aligned} \iint_S \vec{F}(x, y, z) \cdot d\vec{S} &= \iint_S \vec{F}(x, y, z) \cdot \vec{n} dS \\ &= \iint_{D(x, y)} \vec{F}(x, y, z) \cdot \left( \frac{\partial z}{\partial x} \vec{i} + \frac{\partial z}{\partial y} \vec{j} - \vec{k} \right) dx dy. \end{aligned}$$

**1144.** 
$$\iint_S (\vec{F} \cdot \vec{n}) dS = \iint_S P dy dz + Q dz dx + R dx dy$$

$$= \iint_S (P \cos \alpha + Q \cos \beta + R \cos \gamma) dS,$$

where  $P(x, y, z)$ ,  $Q(x, y, z)$ ,  $R(x, y, z)$  are the components of the vector field  $\vec{F}$ .

$\cos \alpha$ ,  $\cos \beta$ ,  $\cos \gamma$  are the angles between the outer unit normal vector  $\vec{n}$  and the  $x$ -axis,  $y$ -axis, and  $z$ -axis, respectively.

## CHAPTER 9. INTEGRAL CALCULUS

- 1145.** If the surface  $S$  is given in parametric form by the vector  $\vec{r}(x(u,v), y(u,v), z(u,v))$ , then the latter formula can be written as

$$\iint_S (\vec{F} \cdot \vec{n}) dS = \iint_S P dydz + Q dzdx + R dx dy = \iint_{D(u,v)} \begin{vmatrix} P & Q & R \\ \frac{\partial x}{\partial u} & \frac{\partial y}{\partial u} & \frac{\partial z}{\partial u} \\ \frac{\partial x}{\partial v} & \frac{\partial y}{\partial v} & \frac{\partial z}{\partial v} \end{vmatrix} du dv,$$

where  $(u,v)$  ranges over some domain  $D(u,v)$  of the  $uv$ -plane.

- 1146.** Divergence Theorem

$$\oiint_S \vec{F} \cdot d\vec{S} = \iiint_G (\nabla \cdot \vec{F}) dV,$$

where

$$\vec{F}(x, y, z) = \langle P(x, y, z), Q(x, y, z), R(x, y, z) \rangle$$

is a vector field whose components  $P$ ,  $Q$ , and  $R$  have continuous partial derivatives,

$$\nabla \cdot \vec{F} = \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z}$$

is the **divergence** of  $\vec{F}$ , also denoted  $\text{div} \vec{F}$ . The symbol  $\oiint$  indicates that the surface integral is taken over a closed surface.

- 1147.** Divergence Theorem in Coordinate Form

$$\oiint_S P dydz + Q dx dz + R dx dy = \iiint_G \left( \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z} \right) dx dy dz.$$

- 1148.** Stoke's Theorem

$$\oint_C \vec{F} \cdot d\vec{r} = \iint_S (\nabla \times \vec{F}) \cdot d\vec{S},$$

## CHAPTER 9. INTEGRAL CALCULUS

where

$$\vec{F}(x, y, z) = \langle P(x, y, z), Q(x, y, z), R(x, y, z) \rangle$$

is a vector field whose components  $P$ ,  $Q$ , and  $R$  have continuous partial derivatives,

$$\nabla \times \vec{F} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ P & Q & R \end{vmatrix} = \left( \frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z} \right) \vec{i} + \left( \frac{\partial P}{\partial z} - \frac{\partial R}{\partial x} \right) \vec{j} + \left( \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \vec{k}$$

is the **curl** of  $\vec{F}$ , also denoted  $\text{curl } \vec{F}$ .

The symbol  $\oint$  indicates that the line integral is taken over a closed curve.

**1149.** Stoke's Theorem in Coordinate Form

$$\begin{aligned} \oint_C P dx + Q dy + R dz \\ = \iint_S \left( \frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z} \right) dy dz + \left( \frac{\partial P}{\partial z} - \frac{\partial R}{\partial x} \right) dz dx + \left( \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx dy \end{aligned}$$

**1150.** Surface Area

$$A = \iint_S dS$$

**1151.** If the surface  $S$  is parameterized by the vector

$$\vec{r}(u, v) = x(u, v)\vec{i} + y(u, v)\vec{j} + z(u, v)\vec{k},$$

then the surface area is

$$A = \iint_{D(u, v)} \left| \frac{\partial \vec{r}}{\partial u} \times \frac{\partial \vec{r}}{\partial v} \right| du dv,$$

where  $D(u, v)$  is the domain where the surface  $\vec{r}(u, v)$  is defined.

## CHAPTER 9. INTEGRAL CALCULUS

**1152.** If  $S$  is given explicitly by the function  $z(x, y)$ , then the surface area is

$$A = \iint_{D(x,y)} \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2} dx dy,$$

where  $D(x, y)$  is the projection of the surface  $S$  onto the  $xy$ -plane.

**1153.** Mass of a Surface

$$m = \iint_S \mu(x, y, z) dS,$$

where  $\mu(x, y, z)$  is the mass per unit area (density function).

**1154.** Center of Mass of a Shell

$$\bar{x} = \frac{M_{yz}}{m}, \quad \bar{y} = \frac{M_{xz}}{m}, \quad \bar{z} = \frac{M_{xy}}{m},$$

where

$$M_{yz} = \iint_S x\mu(x, y, z) dS,$$

$$M_{xz} = \iint_S y\mu(x, y, z) dS,$$

$$M_{xy} = \iint_S z\mu(x, y, z) dS$$

are the first moments about the coordinate planes  $x = 0$ ,  $y = 0$ ,  $z = 0$ , respectively.  $\mu(x, y, z)$  is the density function.

**1155.** Moments of Inertia about the  $xy$ -plane (or  $z = 0$ ),  $yz$ -plane ( $x = 0$ ), and  $xz$ -plane ( $y = 0$ )

$$I_{xy} = \iint_S z^2 \mu(x, y, z) dS,$$

$$I_{yz} = \iint_S x^2 \mu(x, y, z) dS,$$

## CHAPTER 9. INTEGRAL CALCULUS

$$I_{xz} = \iint_S y^2 \mu(x, y, z) dS.$$

**1156.** Moments of Inertia about the x-axis, y-axis, and z-axis

$$I_x = \iint_S (y^2 + z^2) \mu(x, y, z) dS,$$

$$I_y = \iint_S (x^2 + z^2) \mu(x, y, z) dS,$$

$$I_z = \iint_S (x^2 + y^2) \mu(x, y, z) dS.$$

**1157.** Volume of a Solid Bounded by a Closed Surface

$$V = \frac{1}{3} \left| \oiint_S x dy dz + y dx dz + z dx dy \right|$$

**1158.** Gravitational Force

$$\vec{F} = Gm \iint_S \mu(x, y, z) \frac{\vec{r}}{r^3} dS,$$

where  $m$  is a mass at a point  $\langle x_0, y_0, z_0 \rangle$  outside the surface,

$$\vec{r} = \langle x - x_0, y - y_0, z - z_0 \rangle,$$

$\mu(x, y, z)$  is the density function,

and  $G$  is gravitational constant.

**1159.** Pressure Force

$$\vec{F} = \iint_S p(\vec{r}) d\vec{S},$$

where the pressure  $p(\vec{r})$  acts on the surface  $S$  given by the position vector  $\vec{r}$ .

**1160.** Fluid Flux (across the surface  $S$ )

$$\Phi = \oiint_S \vec{v}(\vec{r}) \cdot d\vec{S},$$

CHAPTER 9. INTEGRAL CALCULUS

where  $\vec{v}(\vec{r})$  is the fluid velocity.

**1161.** Mass Flux (across the surface S)

$$\Phi = \iint_S \rho \vec{v}(\vec{r}) \cdot d\vec{S},$$

where  $\vec{F} = \rho \vec{v}$  is the vector field,  $\rho$  is the fluid density.

**1162.** Surface Charge

$$Q = \iint_S \sigma(x, y) dS,$$

where  $\sigma(x, y)$  is the surface charge density.

**1163.** Gauss' Law

The **electric flux** through any closed surface is proportional to the charge Q enclosed by the surface

$$\Phi = \iint_S \vec{E} \cdot d\vec{S} = \frac{Q}{\epsilon_0},$$

where

$\Phi$  is the electric flux,

$\vec{E}$  is the magnitude of the electric field strength,

$\epsilon_0 = 8,85 \times 10^{-12} \frac{F}{m}$  is permittivity of free space.

## **Chapter 10**

# **Differential Equations**

Functions of one variable:  $y, p, q, u, g, h, G, H, r, z$

Arguments (independent variables):  $x, y$

Functions of two variables:  $f(x, y), M(x, y), N(x, y)$

First order derivative:  $y', u', \dot{y}, \frac{dy}{dt}, \dots$

Second order derivatives:  $y'', \ddot{y}, \frac{d^2I}{dt^2}, \dots$

Partial derivatives:  $\frac{\partial u}{\partial t}, \frac{\partial^2 u}{\partial x^2}, \dots$

Natural number:  $n$

Particular solutions:  $y_1, y_p$

Real numbers:  $k, t, C, C_1, C_2, p, q, \alpha, \beta$

Roots of the characteristic equations:  $\lambda_1, \lambda_2$

Time:  $t$

Temperature:  $T, S$

Population function:  $P(t)$

Mass of an object:  $m$

Stiffness of a spring:  $k$

Displacement of the mass from equilibrium:  $y$

Amplitude of the displacement:  $A$

Frequency:  $\omega$

Damping coefficient:  $\gamma$

Phase angle of the displacement:  $\delta$

Angular displacement:  $\theta$

Pendulum length:  $L$



CHAPTER 10. DIFFERENTIAL EQUATIONS

Acceleration of gravity:  $g$

Current:  $I$

Resistance:  $R$

Inductance:  $L$

Capacitance:  $C$

## 10.1 First Order Ordinary Differential Equations

### 1164. Linear Equations

$$\frac{dy}{dx} + p(x)y = q(x).$$

The general solution is

$$y = \frac{\int u(x)q(x)dx + C}{u(x)},$$

where

$$u(x) = \exp\left(\int p(x)dx\right).$$

### 1165. Separable Equations

$$\frac{dy}{dx} = f(x, y) = g(x)h(y)$$

The general solution is given by

$$\int \frac{dy}{h(y)} = \int g(x)dx + C,$$

or

$$H(y) = G(x) + C.$$

## CHAPTER 10. DIFFERENTIAL EQUATIONS

**1166.** Homogeneous Equations

The differential equation  $\frac{dy}{dx} = f(x, y)$  is homogeneous, if the function  $f(x, y)$  is homogeneous, that is  $f(tx, ty) = f(x, y)$ .

The substitution  $z = \frac{y}{x}$  (then  $y = zx$ ) leads to the separable equation

$$x \frac{dz}{dx} + z = f(1, z).$$

**1167.** Bernoulli Equation

$$\frac{dy}{dx} + p(x)y = q(x)y^n.$$

The substitution  $z = y^{1-n}$  leads to the linear equation

$$\frac{dz}{dx} + (1-n)p(x)z = (1-n)q(x).$$

**1168.** Riccati Equation

$$\frac{dy}{dx} = p(x) + q(x)y + r(x)y^2$$

If a particular solution  $y_1$  is known, then the general solution can be obtained with the help of substitution

$z = \frac{1}{y - y_1}$ , which leads to the first order linear equation

$$\frac{dz}{dx} = -[q(x) + 2y_1 r(x)]z - r(x).$$

CHAPTER 10. DIFFERENTIAL EQUATIONS

**1169.** Exact and Nonexact Equations

The equation

$$M(x, y)dx + N(x, y)dy = 0$$

is called **exact** if

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x},$$

and **nonexact** otherwise.

The general solution is

$$\int M(x, y)dx + \int N(x, y)dy = C.$$

**1170.** Radioactive Decay

$$\frac{dy}{dt} = -ky,$$

where  $y(t)$  is the amount of radioactive element at time  $t$ ,  $k$  is the rate of decay.

The solution is

$$y(t) = y_0 e^{-kt}, \text{ where } y_0 = y(0) \text{ is the initial amount.}$$

**1171.** Newton's Law of Cooling

$$\frac{dT}{dt} = -k(T - S),$$

where  $T(t)$  is the temperature of an object at time  $t$ ,  $S$  is the temperature of the surrounding environment,  $k$  is a positive constant.

The solution is

$$T(t) = S + (T_0 - S)e^{-kt},$$

where  $T_0 = T(0)$  is the initial temperature of the object at time  $t = 0$ .

## CHAPTER 10. DIFFERENTIAL EQUATIONS

**1172.** Population Dynamics (Logistic Model)

$$\frac{dP}{dt} = kP \left( 1 - \frac{P}{M} \right),$$

where  $P(t)$  is population at time  $t$ ,  $k$  is a positive constant,  $M$  is a limiting size for the population.

The solution of the differential equation is

$$P(t) = \frac{MP_0}{P_0 + (M - P_0)e^{-kt}}, \text{ where } P_0 = P(0) \text{ is the initial population at time } t = 0.$$

## 10.2 Second Order Ordinary Differential Equations

**1173.** Homogeneous Linear Equations with Constant Coefficients

$$y'' + py' + qy = 0.$$

The characteristic equation is

$$\lambda^2 + p\lambda + q = 0.$$

If  $\lambda_1$  and  $\lambda_2$  are distinct real roots of the characteristic equation, then the general solution is

$$y = C_1 e^{\lambda_1 x} + C_2 e^{\lambda_2 x}, \text{ where}$$

$C_1$  and  $C_2$  are integration constants.

If  $\lambda_1 = \lambda_2 = -\frac{p}{2}$ , then the general solution is

$$y = (C_1 + C_2 x) e^{-\frac{p}{2}x}.$$

If  $\lambda_1$  and  $\lambda_2$  are complex numbers:

## CHAPTER 10. DIFFERENTIAL EQUATIONS

$\lambda_1 = \alpha + \beta i$ ,  $\lambda_2 = \alpha - \beta i$ , where

$$\alpha = -\frac{p}{2}, \beta = \frac{\sqrt{4q - p^2}}{2},$$

then the general solution is

$$y = e^{\alpha x} (C_1 \cos \beta x + C_2 \sin \beta x).$$

**1174.** Inhomogeneous Linear Equations with Constant Coefficients

$$y'' + py' + qy = f(x).$$

The general solution is given by

$$y = y_p + y_h, \text{ where}$$

$y_p$  is a particular solution of the inhomogeneous equation and  $y_h$  is the general solution of the associated homogeneous equation (see the previous topic 1173).

If the right side has the form

$$f(x) = e^{\alpha x} (P_1(x) \cos \beta x + P_2(x) \sin \beta x),$$

then the particular solution  $y_p$  is given by

$$y_p = x^k e^{\alpha x} (R_1(x) \cos \beta x + R_2(x) \sin \beta x),$$

where the polynomials  $R_1(x)$  and  $R_2(x)$  have to be found by using the **method of undetermined coefficients**.

- If  $\alpha + \beta i$  is not a root of the characteristic equation, then the power  $k = 0$ ,
- If  $\alpha + \beta i$  is a simple root, then  $k = 1$ ,
- If  $\alpha + \beta i$  is a double root, then  $k = 2$ .

**1175.** Differential Equations with  $y$  Missing

$$y'' = f(x, y').$$

Set  $u = y'$ . Then the new equation satisfied by  $v$  is

$$u' = f(x, u),$$

which is a first order differential equation.

CHAPTER 10. DIFFERENTIAL EQUATIONS

**1176.** Differential Equations with x Missing

$$y'' = f(y, y').$$

Set  $u = y'$ . Since

$$y'' = \frac{du}{dx} = \frac{du}{dy} \frac{dy}{dx} = u \frac{du}{dy},$$

we have

$$u \frac{du}{dy} = f(y, u),$$

which is a first order differential equation.

**1177.** Free Undamped Vibrations

The motion of a Mass on a Spring is described by the equation

$$m\ddot{y} + ky = 0,$$

where

$m$  is the mass of the object,

$k$  is the stiffness of the spring,

$y$  is displacement of the mass from equilibrium.

The general solution is

$$y = A \cos(\omega_0 t - \delta),$$

where

$A$  is the amplitude of the displacement,

$\omega_0$  is the fundamental frequency, the period is  $T = \frac{2\pi}{\omega_0}$ ,

$\delta$  is phase angle of the displacement.

This is an example of simple harmonic motion.

**1178.** Free Damped Vibrations

$$m\ddot{y} + \gamma\dot{y} + ky = 0, \text{ where}$$

$\gamma$  is the damping coefficient.

There are 3 cases for the general solution:

## CHAPTER 10. DIFFERENTIAL EQUATIONS

Case 1.  $\gamma^2 > 4km$  (overdamped)

$$y(t) = Ae^{\lambda_1 t} + Be^{\lambda_2 t},$$

where

$$\lambda_1 = \frac{-\gamma - \sqrt{\gamma^2 - 4km}}{2m}, \quad \lambda_2 = \frac{-\gamma + \sqrt{\gamma^2 - 4km}}{2m}.$$

Case 2.  $\gamma^2 = 4km$  (critically damped)

$$y(t) = (A + Bt)e^{\lambda t},$$

where

$$\lambda = -\frac{\gamma}{2m}.$$

Case 3.  $\gamma^2 < 4km$  (underdamped)

$$y(t) = e^{-\frac{\gamma}{2m}t} A \cos(\omega t - \delta), \text{ where}$$

$$\omega = \sqrt{4km - \gamma^2}.$$

### 1179. Simple Pendulum

$$\frac{d^2\theta}{dt^2} + \frac{g}{L}\theta = 0,$$

where  $\theta$  is the angular displacement,  $L$  is the pendulum length,  $g$  is the acceleration of gravity.

The general solution for small angles  $\theta$  is

$$\theta(t) = \theta_{\max} \sin \sqrt{\frac{g}{L}}t, \text{ the period is } T = 2\pi \sqrt{\frac{L}{g}}.$$

### 1180. RLC Circuit

$$L \frac{d^2I}{dt^2} + R \frac{dI}{dt} + \frac{1}{C}I = V'(t) = \omega E_0 \cos(\omega t),$$

## CHAPTER 10. DIFFERENTIAL EQUATIONS

where  $I$  is the current in an RLC circuit with an ac voltage source  $V(t) = E_0 \sin(\omega t)$ .

The general solution is

$$I(t) = C_1 e^{r_1 t} + C_2 e^{r_2 t} + A \sin(\omega t - \varphi),$$

where

$$r_{1,2} = \frac{-R \pm \sqrt{R^2 - \frac{4L}{C}}}{2L},$$

$$A = \frac{\omega E_0}{\sqrt{\left(L\omega^2 - \frac{1}{C}\right)^2 + R^2 \omega^2}},$$

$$\varphi = \arctan\left(\frac{L\omega}{R} - \frac{1}{RC\omega}\right),$$

$C_1, C_2$  are constants depending on initial conditions.

### 10.3. Some Partial Differential Equations

#### 1181. The Laplace Equation

$$\frac{\partial^2 \mathbf{u}}{\partial x^2} + \frac{\partial^2 \mathbf{u}}{\partial y^2} = 0$$

applies to potential energy function  $u(x, y)$  for a conservative force field in the  $xy$ -plane. Partial differential equations of this type are called **elliptic**.

#### 1182. The Heat Equation

$$\frac{\partial^2 \mathbf{u}}{\partial x^2} + \frac{\partial^2 \mathbf{u}}{\partial y^2} = \frac{\partial \mathbf{u}}{\partial t}$$



CHAPTER 10. DIFFERENTIAL EQUATIONS

applies to the temperature distribution  $u(x, y)$  in the  $xy$ -plane when heat is allowed to flow from warm areas to cool ones. The equations of this type are called **parabolic**.

**1183.** The Wave Equation

$$\frac{\partial^2 \mathbf{u}}{\partial x^2} + \frac{\partial^2 \mathbf{u}}{\partial y^2} = \frac{\partial^2 \mathbf{u}}{\partial t^2}$$

applies to the displacement  $u(x, y)$  of vibrating membranes and other wave functions. The equations of this type are called **hyperbolic**.

## Chapter 11

### Series

#### 11.1 Arithmetic Series

Initial term:  $a_1$

Nth term:  $a_n$

Difference between successive terms:  $d$

Number of terms in the series:  $n$

Sum of the first  $n$  terms:  $S_n$

$$1184. a_n = a_{n-1} + d = a_{n-2} + 2d = \dots = a_1 + (n-1)d$$

$$1185. a_1 + a_n = a_2 + a_{n-1} = \dots = a_i + a_{n+1-i}$$

$$1186. a_i = \frac{a_{i-1} + a_{i+1}}{2}$$

$$1187. S_n = \frac{a_1 + a_n}{2} \cdot n = \frac{2a_1 + (n-1)d}{2} \cdot n$$

CHAPTER 11. SERIES

## 11.2 Geometric Series

Initial term:  $a_1$

Nth term:  $a_n$

Common ratio:  $q$

Number of terms in the series:  $n$

Sum of the first  $n$  terms:  $S_n$

Sum to infinity:  $S$

$$1188. a_n = qa_{n-1} = a_1q^{n-1}$$

$$1189. a_1 \cdot a_n = a_2 \cdot a_{n-1} = \dots = a_i \cdot a_{n+1-i}$$

$$1190. a_i = \sqrt{a_{i-1} \cdot a_{i+1}}$$

$$1191. S_n = \frac{a_nq - a_1}{q-1} = \frac{a_1(q^n - 1)}{q-1}$$

$$1192. S = \lim_{n \rightarrow \infty} S_n = \frac{a_1}{1-q}$$

For  $|q| < 1$ , the sum  $S$  converges as  $n \rightarrow \infty$ .

## 11.3 Some Finite Series

Number of terms in the series:  $n$

CHAPTER 11. SERIES

$$1193. 1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$$

$$1194. 2 + 4 + 6 + \dots + 2n = n(n+1)$$

$$1195. 1 + 3 + 5 + \dots + (2n-1) = n^2$$

$$1196. k + (k+1) + (k+2) + \dots + (k+n-1) = \frac{n(2k+n-1)}{2}$$

$$1197. 1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}$$

$$1198. 1^3 + 2^3 + 3^3 + \dots + n^3 = \left[ \frac{n(n+1)}{2} \right]^2$$

$$1199. 1^2 + 3^2 + 5^2 + \dots + (2n-1)^2 = \frac{n(4n^2-1)}{3}$$

$$1200. 1^3 + 3^3 + 5^3 + \dots + (2n-1)^3 = n^2(2n^2-1)$$

$$1201. 1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots + \frac{1}{2^n} + \dots = 2$$

$$1202. \frac{1}{1 \cdot 2} + \frac{1}{2 \cdot 3} + \frac{1}{3 \cdot 4} + \dots + \frac{1}{n(n+1)} + \dots = 1$$

$$1203. 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots + \frac{1}{(n-1)!} + \dots = e$$

CHAPTER 11. SERIES

## 11.4 Infinite Series

Sequence:  $\{a_n\}$

First term:  $a_1$

Nth term:  $a_n$

### 1204. Infinite Series

$$\sum_{n=1}^{\infty} a_n = a_1 + a_2 + \dots + a_n + \dots$$

### 1205. Nth Partial Sum

$$S_n = \sum_{n=1}^n a_n = a_1 + a_2 + \dots + a_n$$

### 1206. Convergence of Infinite Series

$$\sum_{n=1}^{\infty} a_n = L, \text{ if } \lim_{n \rightarrow \infty} S_n = L$$

### 1207. Nth Term Test

- If the series  $\sum_{n=1}^{\infty} a_n$  is convergent, then  $\lim_{n \rightarrow \infty} a_n = 0$ .
- If  $\lim_{n \rightarrow \infty} a_n \neq 0$ , then the series is divergent.

## 11.5 Properties of Convergent Series

Convergent Series:  $\sum_{n=1}^{\infty} a_n = A, \sum_{n=1}^{\infty} b_n = B$

Real number:  $c$

## CHAPTER 11. SERIES

$$1208. \sum_{n=1}^{\infty} (a_n + b_n) = \sum_{n=1}^{\infty} a_n + \sum_{n=1}^{\infty} b_n = A + B$$

$$1209. \sum_{n=1}^{\infty} ca_n = c \sum_{n=1}^{\infty} a_n = cA.$$

## 11.6 Convergence Tests

## 1210. The Comparison Test

Let  $\sum_{n=1}^{\infty} a_n$  and  $\sum_{n=1}^{\infty} b_n$  be series such that  $0 < a_n \leq b_n$  for all  $n$ .

- If  $\sum_{n=1}^{\infty} b_n$  is convergent then  $\sum_{n=1}^{\infty} a_n$  is also convergent.
- If  $\sum_{n=1}^{\infty} a_n$  is divergent then  $\sum_{n=1}^{\infty} b_n$  is also divergent.

## 1211. The Limit Comparison Test

Let  $\sum_{n=1}^{\infty} a_n$  and  $\sum_{n=1}^{\infty} b_n$  be series such that  $a_n$  and  $b_n$  are positive for all  $n$ .

- If  $0 < \lim_{n \rightarrow \infty} \frac{a_n}{b_n} < \infty$  then  $\sum_{n=1}^{\infty} a_n$  and  $\sum_{n=1}^{\infty} b_n$  are either both convergent or both divergent.
- If  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 0$  then  $\sum_{n=1}^{\infty} b_n$  convergent implies that  $\sum_{n=1}^{\infty} a_n$  is also convergent.

## CHAPTER 11. SERIES

- If  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \infty$  then  $\sum_{n=1}^{\infty} b_n$  divergent implies that  $\sum_{n=1}^{\infty} a_n$  is also divergent.

**1212.** p-series

p-series  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  converges for  $p > 1$  and diverges for  $0 < p \leq 1$ .

**1213.** The Integral Test

Let  $f(x)$  be a function which is continuous, positive, and decreasing for all  $x \geq 1$ . The series

$$\sum_{n=1}^{\infty} f(n) = f(1) + f(2) + f(3) + \dots + f(n) + \dots$$

converges if  $\int_1^{\infty} f(x) dx$  converges, and diverges if

$$\int_1^n f(x) dx \rightarrow \infty \text{ as } n \rightarrow \infty.$$

**1214.** The Ratio Test

Let  $\sum_{n=1}^{\infty} a_n$  be a series with positive terms.

- If  $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} < 1$  then  $\sum_{n=1}^{\infty} a_n$  is convergent.
- If  $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} > 1$  then  $\sum_{n=1}^{\infty} a_n$  is divergent.
- If  $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = 1$  then  $\sum_{n=1}^{\infty} a_n$  may converge or diverge and the ratio test is inconclusive; some other tests must be used.

## CHAPTER 11. SERIES

**1215.** The Root Test

Let  $\sum_{n=1}^{\infty} a_n$  be a series with positive terms.

- If  $\lim_{n \rightarrow \infty} \sqrt[n]{a_n} < 1$  then  $\sum_{n=1}^{\infty} a_n$  is convergent.
- If  $\lim_{n \rightarrow \infty} \sqrt[n]{a_n} > 1$  then  $\sum_{n=1}^{\infty} a_n$  is divergent.
- If  $\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = 1$  then  $\sum_{n=1}^{\infty} a_n$  may converge or diverge, but no conclusion can be drawn from this test.

## 11.7 Alternating Series

**1216.** The Alternating Series Test (Leibniz's Theorem)

Let  $\{a_n\}$  be a sequence of positive numbers such that  $a_{n+1} < a_n$  for all  $n$ .

$$\lim_{n \rightarrow \infty} a_n = 0.$$

Then the alternating series  $\sum_{n=1}^{\infty} (-1)^n a_n$  and  $\sum_{n=1}^{\infty} (-1)^{n-1} a_n$  both converge.

**1217.** Absolute Convergence

- A series  $\sum_{n=1}^{\infty} a_n$  is **absolutely convergent** if the series  $\sum_{n=1}^{\infty} |a_n|$  is convergent.



## CHAPTER 11. SERIES

- If the series  $\sum_{n=1}^{\infty} a_n$  is absolutely convergent then it is convergent.

**1218.** Conditional Convergence

A series  $\sum_{n=1}^{\infty} a_n$  is **conditionally convergent** if the series is convergent but is not absolutely convergent.

**11.8** Power Series

Real numbers:  $x, x_0$

Power series:  $\sum_{n=0}^{\infty} a_n x^n, \sum_{n=0}^{\infty} a_n (x - x_0)^n$

Whole number:  $n$

Radius of Convergence:  $R$

**1219.** Power Series in  $x$ 

$$\sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + \dots + a_n x^n + \dots$$

**1220.** Power Series in  $(x - x_0)$ 

$$\sum_{n=0}^{\infty} a_n (x - x_0)^n = a_0 + a_1 (x - x_0) + a_2 (x - x_0)^2 + \dots + a_n (x - x_0)^n + \dots$$

**1221.** Interval of Convergence

The set of those values of  $x$  for which the function

$f(x) = \sum_{n=0}^{\infty} a_n (x - x_0)^n$  is convergent is called the **interval of convergence**.

## CHAPTER 11. SERIES

**1222.** Radius of Convergence

If the interval of convergence is  $(x_0 - R, x_0 + R)$  for some  $R \geq 0$ , the  $R$  is called the **radius of convergence**. It is given as

$$R = \lim_{n \rightarrow \infty} \frac{1}{\sqrt[n]{a_n}} \quad \text{or} \quad R = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right|.$$

## 11.9 Differentiation and Integration of Power Series

Continuous function:  $f(x)$

Power series:  $\sum_{n=0}^{\infty} a_n x^n$

Whole number:  $n$

Radius of Convergence:  $R$

**1223.** Differentiation of Power Series

Let  $f(x) = \sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + \dots$  for  $|x| < R$ .

Then, for  $|x| < R$ ,  $f(x)$  is continuous, the derivative  $f'(x)$  exists and

$$\begin{aligned} f'(x) &= \frac{d}{dx} a_0 + \frac{d}{dx} a_1 x + \frac{d}{dx} a_2 x^2 + \dots \\ &= a_1 + 2a_2 x + 3a_3 x^2 + \dots = \sum_{n=1}^{\infty} n a_n x^{n-1}. \end{aligned}$$

## CHAPTER 11. SERIES

**1224.** Integration of Power Series

Let  $f(x) = \sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + \dots$  for  $|x| < R$ .

Then, for  $|x| < R$ , the indefinite integral  $\int f(x) dx$  exists and

$$\begin{aligned} \int f(x) dx &= \int a_0 dx + \int a_1 x dx + \int a_2 x^2 dx + \dots \\ &= a_0 x + a_1 \frac{x^2}{2} + a_2 \frac{x^3}{3} + \dots = \sum_{n=0}^{\infty} a_n \frac{x^{n+1}}{n+1} + C. \end{aligned}$$

## 11.10 Taylor and Maclaurin Series

Whole number:  $n$

Differentiable function:  $f(x)$

Remainder term:  $R_n$

**1225.** Taylor Series

$$\begin{aligned} f(x) &= \sum_{n=0}^{\infty} f^{(n)}(a) \frac{(x-a)^n}{n!} = f(a) + f'(a)(x-a) + \frac{f''(a)(x-a)^2}{2!} + \dots \\ &\quad + \frac{f^{(n)}(a)(x-a)^n}{n!} + R_n. \end{aligned}$$

**1226.** The Remainder After  $n+1$  Terms is given by

$$R_n = \frac{f^{(n+1)}(\xi)(x-a)^{n+1}}{(n+1)!}, \quad a < \xi < x.$$

**1227.** Maclaurin Series

## CHAPTER 11. SERIES

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)x^n}{n!} = f(0) + f'(0)x + \frac{f''(0)x^2}{2!} + \dots + \frac{f^{(n)}(0)x^n}{n!} + R_n$$

## 11.11 Power Series Expansions for Some Functions

Whole number:  $n$ Real number:  $x$ 

$$1228. e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots + \frac{x^n}{n!} + \dots$$

$$1229. a^x = 1 + \frac{x \ln a}{1!} + \frac{(x \ln a)^2}{2!} + \frac{(x \ln a)^3}{3!} + \dots + \frac{(x \ln a)^n}{n!} + \dots$$

$$1230. \ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots + \frac{(-1)^n x^{n+1}}{n+1} \pm \dots, -1 < x \leq 1.$$

$$1231. \ln \frac{1+x}{1-x} = 2 \left( x + \frac{x^3}{3} + \frac{x^5}{5} + \frac{x^7}{7} + \dots \right), |x| < 1.$$

$$1232. \ln x = 2 \left[ \frac{x-1}{x+1} + \frac{1}{3} \left( \frac{x-1}{x+1} \right)^3 + \frac{1}{5} \left( \frac{x-1}{x+1} \right)^5 \dots \right], x > 0.$$

$$1233. \cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots + \frac{(-1)^n x^{2n}}{(2n)!} \pm \dots$$

## CHAPTER 11. SERIES

$$1234. \sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots + \frac{(-1)^n x^{2n+1}}{(2n+1)!} \pm \dots$$

$$1235. \tan x = x + \frac{x^3}{3} + \frac{2x^5}{15} + \frac{17x^7}{315} + \frac{62x^9}{2835} + \dots, |x| < \frac{\pi}{2}.$$

$$1236. \cot x = \frac{1}{x} - \left( \frac{x}{3} + \frac{x^3}{45} + \frac{2x^5}{945} + \frac{2x^7}{4725} + \dots \right), |x| < \pi.$$

$$1237. \arcsin x = x + \frac{x^3}{2 \cdot 3} + \frac{1 \cdot 3x^5}{2 \cdot 4 \cdot 5} + \dots + \frac{1 \cdot 3 \cdot 5 \dots (2n-1)x^{2n+1}}{2 \cdot 4 \cdot 6 \dots (2n)(2n+1)} + \dots,$$

$$|x| < 1.$$

$$1238. \arccos x = \frac{\pi}{2} - \left( x + \frac{x^3}{2 \cdot 3} + \frac{1 \cdot 3x^5}{2 \cdot 4 \cdot 5} + \dots + \frac{1 \cdot 3 \cdot 5 \dots (2n-1)x^{2n+1}}{2 \cdot 4 \cdot 6 \dots (2n)(2n+1)} + \dots \right),$$

$$|x| < 1.$$

$$1239. \arctan x = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots + \frac{(-1)^n x^{2n+1}}{2n+1} \pm \dots, |x| \leq 1.$$

$$1240. \cosh x = 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \frac{x^6}{6!} + \dots + \frac{x^{2n}}{(2n)!} + \dots$$

$$1241. \sinh x = x + \frac{x^3}{3!} + \frac{x^5}{5!} + \frac{x^7}{7!} + \dots + \frac{x^{2n+1}}{(2n+1)!} + \dots$$

## CHAPTER 11. SERIES

## 11.12 Binomial Series

Whole numbers:  $n, m$ Real number:  $x$ Combinations:  ${}^n C_m$ 

$$1242. (1+x)^n = 1 + {}^n C_1 x + {}^n C_2 x^2 + \dots + {}^n C_n x^n + \dots + x^n$$

$$1243. {}^n C_m = \frac{n(n-1)\dots[n-(m-1)]}{m!}, |x| < 1.$$

$$1244. \frac{1}{1+x} = 1 - x + x^2 - x^3 + \dots, |x| < 1.$$

$$1245. \frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots, |x| < 1.$$

$$1246. \sqrt{1+x} = 1 + \frac{x}{2} - \frac{x^2}{2 \cdot 4} + \frac{1 \cdot 3x^3}{2 \cdot 4 \cdot 6} - \frac{1 \cdot 3 \cdot 5x^4}{2 \cdot 4 \cdot 6 \cdot 8} + \dots, |x| \leq 1.$$

$$1247. \sqrt[3]{1+x} = 1 + \frac{x}{3} - \frac{1 \cdot 2x^2}{3 \cdot 6} + \frac{1 \cdot 2 \cdot 5x^3}{3 \cdot 6 \cdot 9} - \frac{1 \cdot 2 \cdot 5 \cdot 8x^4}{3 \cdot 6 \cdot 9 \cdot 12} + \dots, |x| \leq 1.$$

## 11.13 Fourier Series

Integrable function:  $f(x)$ Fourier coefficients:  $a_0, a_n, b_n$ Whole number:  $n$

CHAPTER 11. SERIES

$$1248. f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx)$$

$$1249. a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx \, dx$$

$$1250. b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx \, dx$$

## Chapter 12

# Probability

### 12.1 Permutations and Combinations

Permutations:  ${}^n P_m$   
Combinations:  ${}^n C_m$   
Whole numbers:  $n, m$

**1251.** Factorial  
 $n! = 1 \cdot 2 \cdot 3 \cdots (n-2)(n-1)n$   
 $0! = 1$

**1252.**  ${}^n P_n = n!$

**1253.**  ${}^n P_m = \frac{n!}{(n-m)!}$

**1254.** Binomial Coefficient  
 ${}^n C_m = \binom{n}{m} = \frac{n!}{m!(n-m)!}$

**1255.**  ${}^n C_m = {}^n C_{n-m}$

**1256.**  ${}^n C_m + {}^n C_{m+1} = {}^{n+1} C_{m+1}$



CHAPTER 12. PROBABILITY

**1257.**  ${}^n C_0 + {}^n C_1 + {}^n C_2 + \dots + {}^n C_n = 2^n$

**1258.** Pascal's Triangle

Row 0					1				
Row 1				1		1			
Row 2			1		2		1		
Row 3		1		3		3		1	
Row 4		1		4		6		4	
Row 5	1		5		10		10		5
Row 6	1	6		15		20		15	6

## 12.2 Probability Formulas

- Events: A, B
- Probability: P
- Random variables: X, Y, Z
- Values of random variables: x, y, z
- Expected value of X:  $\mu$
- Any positive real number:  $\varepsilon$
- Standard deviation:  $\sigma$
- Variance:  $\sigma^2$
- Density functions:  $f(x), f(t)$

**1259.** Probability of an Event

$$P(A) = \frac{m}{n},$$

where

m is the number of possible positive outcomes,  
n is the total number of possible outcomes.

CHAPTER 12. PROBABILITY

**1260.** Range of Probability Values

$$0 \leq P(A) \leq 1$$

**1261.** Certain Event

$$P(A) = 1$$

**1262.** Impossible Event

$$P(A) = 0$$

**1263.** Complement

$$P(\bar{A}) = 1 - P(A)$$

**1264.** Independent Events

$$P(A/B) = P(A),$$

$$P(B/A) = P(B)$$

**1265.** Addition Rule for Independent Events

$$P(A \cup B) = P(A) + P(B)$$

**1266.** Multiplication Rule for Independent Events

$$P(A \cap B) = P(A) \cdot P(B)$$

**1267.** General Addition Rule

$$P(A \cup B) = P(A) + P(B) - P(A \cap B),$$

where

$A \cup B$  is the union of events A and B,

$A \cap B$  is the intersection of events A and B.

**1268.** Conditional Probability

$$P(A/B) = \frac{P(A \cap B)}{P(B)}$$

**1269.**  $P(A \cap B) = P(B) \cdot P(A/B) = P(A) \cdot P(B/A)$

## CHAPTER 12. PROBABILITY

**1270.** Law of Total Probability

$$P(A) = \sum_{i=1}^m P(B_i)P(A/B_i),$$

where  $B_i$  is a sequence of mutually exclusive events.

**1271.** Bayes' Theorem

$$P(B/A) = \frac{P(A/B) \cdot P(B)}{P(A)}$$

**1272.** Bayes' Formula

$$P(B_i/A) = \frac{P(B_i) \cdot P(A/B_i)}{\sum_{k=1}^m P(B_k) \cdot P(A/B_k)},$$

where

$B_i$  is a set of mutually exclusive events (hypotheses),

$A$  is the final event,

$P(B_i)$  are the prior probabilities,

$P(B_i/A)$  are the posterior probabilities.

**1273.** Law of Large Numbers

$$P\left(\left|\frac{S_n}{n} - \mu\right| \geq \varepsilon\right) \rightarrow 0 \text{ as } n \rightarrow \infty,$$

$$P\left(\left|\frac{S_n}{n} - \mu\right| < \varepsilon\right) \rightarrow 1 \text{ as } n \rightarrow \infty,$$

where

$S_n$  is the sum of random variables,

$n$  is the number of possible outcomes.

**1274.** Chebyshev Inequality

$$P(|X - \mu| \geq \varepsilon) \leq \frac{V(X)}{\varepsilon^2},$$

where  $V(X)$  is the variance of  $X$ .

CHAPTER 12. PROBABILITY

**1275.** Normal Density Function

$$\varphi(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}},$$

where  $x$  is a particular outcome.

**1276.** Standard Normal Density Function

$$\varphi(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}}$$

Average value  $\mu = 0$ , deviation  $\sigma = 1$ .




Figure 210.

**1277.** Standard Z Value

$$Z = \frac{X - \mu}{\sigma}$$

**1278.** Cumulative Normal Distribution Function

$$F(x) = \frac{1}{\sigma\sqrt{2\pi}} \int_{-\infty}^x e^{-\frac{(t-\mu)^2}{2\sigma^2}} dt,$$

CHAPTER 12. PROBABILITY

where

$x$  is a particular outcome,

$t$  is a variable of integration.

$$1279. P(\alpha < X < \beta) = F\left(\frac{\alpha - \mu}{\sigma}\right) - F\left(\frac{\beta - \mu}{\sigma}\right),$$

where

$X$  is normally distributed random variable,

$F$  is cumulative normal distribution function,

$P(\alpha < X < \beta)$  is interval probability.

$$1280. P(|X - \mu| < \varepsilon) = 2F\left(\frac{\varepsilon}{\sigma}\right),$$

where

$X$  is normally distributed random variable,

$F$  is cumulative normal distribution function.

1281. Cumulative Distribution Function

$$F(x) = P(X < x) = \int_{-\infty}^x f(t) dt,$$

where  $t$  is a variable of integration.

1282. Bernoulli Trials Process

$$\mu = np, \quad \sigma^2 = npq,$$

where

$n$  is a sequence of experiments,

$p$  is the probability of success of each experiments,

$q$  is the probability of failure,  $q = 1 - p$ .

1283. Binomial Distribution Function

$$b(n, p, q) = \binom{n}{k} p^k q^{n-k},$$

CHAPTER 12. PROBABILITY

$$\mu = np, \sigma^2 = npq,$$

$$f(x) = (q + pe^x)^n,$$

where

$n$  is the number of trials of selections,

$p$  is the probability of success,

$q$  is the probability of failure,  $q = 1 - p$ .

**1284.** Geometric Distribution

$$P(T = j) = q^{j-1}p,$$

$$\mu = \frac{1}{p}, \sigma^2 = \frac{q}{p^2},$$

where

$T$  is the first successful event in the series,

$j$  is the event number,

$p$  is the probability that any one event is successful,

$q$  is the probability of failure,  $q = 1 - p$ .

**1285.** Poisson Distribution

$$P(X = k) \approx \frac{\lambda^k}{k!} e^{-\lambda}, \lambda = np,$$

$$\mu = \lambda, \sigma^2 = \lambda,$$

where

$\lambda$  is the rate of occurrence,

$k$  is the number of positive outcomes.

**1286.** Density Function

$$P(a \leq X \leq b) = \int_a^b f(x) dx$$

**1287.** Continuous Uniform Density

$$f = \frac{1}{b-a}, \mu = \frac{a+b}{2},$$

CHAPTER 12. PROBABILITY

where  $f$  is the density function.

**1288.** Exponential Density Function

$$f(t) = \lambda e^{-\lambda t}, \quad \mu = \lambda, \quad \sigma^2 = \lambda^2$$

where  $t$  is time,  $\lambda$  is the failure rate.

**1289.** Exponential Distribution Function

$$F(t) = 1 - e^{-\lambda t},$$

where  $t$  is time,  $\lambda$  is the failure rate.

**1290.** Expected Value of Discrete Random Variables

$$\mu = E(X) = \sum_{i=1}^n x_i p_i,$$

where  $x_i$  is a particular outcome,  $p_i$  is its probability.

**1291.** Expected Value of Continuous Random Variables

$$\mu = E(X) = \int_{-\infty}^{\infty} x f(x) dx$$

**1292.** Properties of Expectations

$$E(X + Y) = E(X) + E(Y),$$

$$E(X - Y) = E(X) - E(Y),$$

$$E(cX) = cE(X),$$

$$E(XY) = E(X) \cdot E(Y),$$

where  $c$  is a constant.

**1293.**  $E(X^2) = V(X) + \mu^2,$

where

$\mu = E(X)$  is the expected value,

$V(X)$  is the variance.

## CHAPTER 12. PROBABILITY

**1294.** Markov Inequality

$$P(X > k) \leq \frac{E(X)}{k},$$

where  $k$  is some constant.

**1295.** Variance of Discrete Random Variables

$$\sigma^2 = V(X) = E[(X - \mu)^2] = \sum_{i=1}^n (x_i - \mu)^2 p_i,$$

where

$x_i$  is a particular outcome,

$p_i$  is its probability.

**1296.** Variance of Continuous Random Variables

$$\sigma^2 = V(X) = E[(X - \mu)^2] = \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx$$

**1297.** Properties of Variance

$$V(X + Y) = V(X) + V(Y),$$

$$V(X - Y) = V(X) + V(Y),$$

$$V(X + c) = V(X),$$

$$V(cX) = c^2 V(X),$$

where  $c$  is a constant.

**1298.** Standard Deviation

$$D(X) = \sqrt{V(X)} = \sqrt{E[(X - \mu)^2]}$$

**1299.** Covariance

$$\text{cov}(X, Y) = E[(X - \mu(X))(Y - \mu(Y))] = E(XY) - \mu(X)\mu(Y),$$

where

$X$  is random variable,

$V(X)$  is the variance of  $X$ ,

$\mu$  is the expected value of  $X$  or  $Y$ .



CHAPTER 12. PROBABILITY

**1300.** Correlation

$$\rho(X, Y) = \frac{\text{cov}(X, Y)}{\sqrt{V(X)V(Y)}},$$

where

$V(X)$  is the variance of  $X$ ,

$V(Y)$  is the variance of  $Y$ .

Look for other handbooks and solved problem guides at  
[www.math-ebooks.com](http://www.math-ebooks.com).

**Get more e-books from [www.ketabton.com](http://www.ketabton.com)  
Ketabton.com: The Digital Library**